




TABLE 14: COMPARISON OF THE AVERAGE NUTRITIONAL VALUES BETWEEN WHITE AND YELLOW MAIZE OVER THE PAST TEN SEASONS

Season	White maize			Yellow maize		
	Fat	Protein	Starch	Fat	Protein	Starch
1996/97	3.9	8.7	-	4.2	8.7	-
1997/98	4.0	8.9	-	4.1	9.0	-
1998/99	4.1	9.2	71.8	4.2	9.5	72.1
1999/00	4.0	8.1	71.9	4.1	8.0	72.0
2000/01	4.2	8.8	70.2	4.2	8.7	70.5
2001/02	4.2	8.9	71.4	4.1	8.9	71.7
2002/03	4.1	9.2	71.4	4.1	9.2	72.0
2003/04	4.0	9.1	71.2	4.0	9.0	71.1
2004/05	4.0	8.9	71.1	3.8	8.6	71.7
2005/06	4.0	8.5	71.1	3.9	8.4	71.5

TABLE 15: AVERAGE NUTRITIONAL VALUES OF SOUTH AFRICAN MAIZE (1996/97 - 2005/06)

Season	Fat %	Protein %	Starch %
1996/97	4.1	8.7	-
1997/98	4.0	9.0	-
1998/99	4.2	9.3	71.9
1999/00	4.0	8.1	71.9
2000/01	4.2	8.8	70.3
2001/02	4.2	8.9	71.5
2002/03	4.1	9.2	71.6
2003/04	4.0	9.1	71.1
2004/05	3.9	8.8	71.3
2005/06	4.0	8.4	71.2
Weighted averages	4.1	8.8	71.3



Please note: Different starch methods have been used over years and data have been corrected accordingly.