




TABLE 14: COMPARISON OF THE AVERAGE NUTRITIONAL VALUES BETWEEN WHITE AND YELLOW MAIZE OVER THE PAST TEN SEASONS

Season	Number of samples	White maize			Number of samples	Yellow maize		
		Fat	Protein	Starch		Fat	Protein	Starch
1998/99	256	4.1	9.2	71.8	189	4.2	9.5	72.1
1999/00	493	4.0	8.1	71.9	407	4.1	8.0	72.0
2000/01	522	4.2	8.8	70.2	378	4.2	8.7	70.5
2001/02	471	4.2	8.9	71.4	429	4.1	8.9	71.7
2002/03	517	4.1	9.2	71.4	383	4.1	9.2	72.0
2003/04	599	4.0	9.1	71.2	301	4.0	9.0	71.1
2004/05	601	4.0	8.9	71.1	399	3.8	8.6	71.7
2005/06	593	4.0	8.5	71.1	307	3.9	8.4	71.5
2006/07	563	3.9	9.3	72.9	337	3.5	9.5	73.3
2007/08	483	4.0	8.6	71.9	417	3.6	8.4	72.3
Weighted averages		4.0	8.9	71.5		3.9	8.8	71.8

TABLE 15: AVERAGE NUTRITIONAL VALUES OF SOUTH AFRICAN MAIZE (1998/99 - 2007/08)

Season	Number of samples	Fat %	Protein %	Starch %
1998/99	445	4.2	9.3	71.9
1999/00	900	4.0	8.1	71.9
2000/01	900	4.2	8.8	70.3
2001/02	900	4.2	8.9	71.5
2002/03	900	4.1	9.2	71.6
2003/04	900	4.0	9.1	71.1
2004/05	1000	3.9	8.8	71.3
2005/06	900	4.0	8.4	71.2
2006/07	900	3.7	9.4	73.0
2007/08	900	3.8	8.5	72.1
Weighted averages		4.0	8.8	71.6



Please note: Different starch methods have been used over years and data have been corrected accordingly.