

Table 17: Presence of Genetically Modified Maize (2008/2009)

Region	RSA Grade	MON810, % (LOD: 0.15%)	NK603 (Roundup Ready), % (LOD: 0.25%)	Region	RSA Grade	MON810, % (LOD: 0.15%)	NK603 (Roundup Ready), % (LOD: 0.25%)
10	YM1	>2	>1.8	25	WM1	>2	>1.8
10	YM1	0.5	0.4	25	YM1	>2	>1.8
11	WM1	>2	0.3	26	WM2	>2	0.4
11	YM1	>2	<0.25	26	YM1	>2	>1.8
12	WM1	>2	>1.8	26	YM1	>2	1.4
12	WM2	>2	0.5	26	YM2	>2	>1.8
12	YM1	>2	1.2	27	WM1	>2	>1.8
12	YM1	>2	>1.8	27	YM1	>2	>1.8
13	WM1	>2	>1.8	28	WM1	>2	>1.8
13	WM1	>2	>1.8	28	WM2	>2	0.8
13	WM1	>2	>1.8	28	YM1	>2	>1.8
14	WM1	>2	1.5	28	YM1	>2	>1.8
14	WM1	>2	>1.8	28	YM1	>2	1.6
14	WM1	>2	>1.8	28	YM1	>2	1.0
14	YM1	>2	1.8	28	YM1	>2	1.5
14	YM1	>2	1.1	28	YM2	>2	>1.8
15	WM1	>2	>1.8	29	WM1	>2	>1.8
15	WM2	>2	>1.8	29	WM1	1.3	>1.8
16	WM1	>2	>1.8	29	WM1	<0.15	>1.8
16	WM1	>2	>1.8	29	WM1	>2	>1.8
16	WM1	>2	>1.8	29	WM1	>2	>1.8
16	WM2	>2	<0.25	30	WM1	>2	>1.8
16	YM1	>2	0.9	30	WM1	0.7	<0.25
17	COM (W)	>2	>1.8	30	WM1	1.7	0.5
17	WM1	>2	>1.8	30	WM2	>2	>1.8
17	WM1	>2	>1.8	30	YM1	>2	1.5
17	WM1	>2	>1.8	30	YM1	>2	>1.8
18	WM1	>2	0.9	30	YM2	>2	>1.8
18	WM1	>2	>1.8	31	WM1	1.4	1.4
19	COM (Y)	>2	>1.8	31	WM1	<0.15	0.4
19	WM1	0.2	0.3	31	WM1	<0.15	<0.25
20	YM1	>2	0.7	31	YM1	<0.15	<0.25
20	YM1	>2	>1.8	32	WM1	<0.15	<0.25
21	WM1	>2	>1.8	32	WM2	<0.15	<0.25
21	WM1	>2	>1.8	32	YM1	>2	>1.8
21	YM1	>2	>1.8	33	YM1	<0.15	<0.25
22	WM1	>2	>1.8	34	WM1	>2	>1.8
22	WM1	>2	>1.8	34	WM1	>2	>1.8
22	YM1	>2	>1.8	34	YM1	<0.15	0.4
23	WM1	>2	>1.8	35	YM1	>2	0.4
23	WM2	>2	>1.8	35	YM1	>2	0.4
24	WM1	>2	>1.8	36	WM1	>2	>1.8
24	YM1	>2	>1.8	36	WM2	>2	>1.8
24	YM1	>2	>1.8	36	YM1	>2	>1.8
25	WM1	>2	<0.25	36	YM1	>2	>1.8
n	Season	% Samples positive for MON810 (Bt)		n	Season	% Samples positive for NK603 (RUR)	
90	2008/2009	91		90	2008/2009	90	
100	2007/2008	95		100	2007/2008	69	
90	2006/2007	97		90	2006/2007	59	
90	2005/2006	91		90	2005/2006	31	
100	2004/2005	78		100	2004/2005	31	