



Limpopo province's two samples had the highest average percentage foreign matter. The weighted average percentage foreign matter in the rest of the samples ranged from 0.54 in the Northern Cape to 0.95 in North West province. Please refer to Graph 17.

Graph 17: Average percentage foreign matter per province over four seasons



North West province (18 samples) reported the highest weighted average percentage soybeans and parts of soybeans above the 1.8 mm slotted sieve which pass through the 4.75 mm round hole sieve, namely 2.15% and Limpopo (2 samples) the lowest at 0.67%. Mpumalanga province with the highest number of samples (77) reported an average of 1.78%. The Free State province averaged 1.77% (42 samples). The national weighted average percentage increased from 1.36% last season to 1.81% this season. Please see Graph 18. No samples were received from KwaZulu-Natal for the 2014/2015 season.

Graph 18: Average percentage soybeans and parts of soybeans which pass through the 4.75 mm round hole sieve per province over four seasons



*Please note that the 2014/2015 results represent soybeans and parts of soybeans above the 1.8 mm slotted sieve which pass through the 4.75 mm round hole sieve.

Eight samples were submitted from Gauteng province and the lowest weighted average percentage defective soybeans on the 4.75 mm sieve were reported on these, namely 1.51%. Mpumalanga province reported the highest percentage of 2.12, followed by the Northern Cape and North West provinces with 2.00 and 1.99 respectively. The national weighted average increased from 1.07% in the previous season to 1.95% this season. Please see Graph 19.