

South African

Commercial Maize Quality for the 2024/2025 season

Acknowledgments


With gratitude to:

- The Maize Trust for financial support in conducting this survey.
- Agbiz Grain and its members for providing maize samples for this survey.
- The Crop Estimates Committee (CEC) of the Department of Agriculture (DoA) for providing production related figures.
- South African Grain Information Service (SAGIS) for providing supply and demand figures relating to maize and maize products.



Introduction

During the 2025 harvesting season, a representative sample of each delivery of maize at the various grain intake points was taken according to the prescribed grading regulation and sampling protocol provided. The sampling procedure for the samples used in this survey is described on page 91. A total of 1 000 composite samples, representing white and yellow maize of each production region, were received and analysed to determine the quality. The samples consisted of 458 white and 542 yellow maize samples.



Graph 1: Provincial contribution to the production of the 2024/25 maize crop

Figures provided by the CEC.

The quality attributes tested, include:

RSA grading: Samples were graded according to the following factors, as defined in the South African grading regulation: defective kernels above and below the 6.35 mm sieve, total defective kernels, foreign matter, other colour kernels and combined deviations.

USA grading: Samples were graded according to the American grading regulations to determine the following factors: Test weight per bushel (pounds), heat damaged kernels, total damaged kernels, broken corn and foreign matter (BCFM) and other colour.

Nutritional values: Moisture, crude protein, crude fat, crude fibre and starch.

Physical quality factors: Test weight (kg/hl), 100 kernel mass, kernel size, breakage susceptibility, stress cracks, milling index and grit yield.

All white maize samples were milled on the Bühler MCKA maize mill, while 50% of the samples were also milled on the Roff laboratory maize mill for comparison purposes. The whiteness index of all the milled maize meal samples was determined.

Mycotoxin analyses were performed on 350 samples representative of white and yellow maize produced per region.