



**SOUTH AFRICAN MAIZE CROP**

**Quality Report  
2020/2021  
Season**



Introduction	1 - 2
Provincial contribution to the production of the 2020/21 crop (Graph 1)	1
Production	3 - 6
Maize production in RSA from 2010/11 to 2020/21 (Graph 2)	3
Total RSA area utilised for maize production from 2010/11 to 2020/21 (Graph 3)	3
RSA maize yield from 2010/11 to 2020/21 (Graph 4)	4
Maize production overview, dryland vs irrigation - 2020/21 season (Table 1)	4
Area utilised for maize production per province over three seasons (Graph 5)	5
Maize production per province over three seasons (Graph 6)	5
Maize yield per province over three seasons (Graph 7)	5
Area utilised for maize production in the Free State, Mpumalanga and North West since 2010/11 (Graphs 8, 10 and 12)	6
Maize production in the Free State, Mpumalanga and North West since 2010/11 (Graphs 9, 11 and 13)	6
Supply and Demand	7 - 8
Maize supply and demand overview 2021/22 marketing season (Graph 14)	8
SAGIS Total Maize Supply and Demand Figures	9
Total Maize: Supply and demand graphs over 10 marketing seasons (Graphs 15 - 18)	10
SAGIS White Maize Supply and Demand Figures	11
White Maize: Supply and demand graphs over 10 marketing seasons (Graphs 19 - 22)	12
SAGIS Yellow Maize Supply and Demand Figures	13
Yellow Maize: Supply and demand graphs over 10 marketing seasons (Graphs 23 - 26)	14
SAGIS White Maize Exports/Imports figures per country and harbour 2021/22 marketing season	15
Major destinations for RSA white maize exports 2021/22 season (Graph 27)	15
SAGIS Yellow Maize Exports/Imports figures per country and harbour 2021/22 marketing season	16
Major destinations for RSA yellow maize exports 2021/22 season (Graph 28)	16
SAGIS Total Whole Maize Processed per Province	17
SAGIS Whole White Maize Processed per Province	18
SAGIS Whole Yellow Maize Processed per Province	19

SAGIS Maize Product Information	20
White and yellow maize products manufactured from May 2021 to March 2022 (Graphs 29 – 32)	20
SAGIS Maize Products Manufactured per Marketing Year	21
SAGIS Maize Products Imported per Marketing Year	22
SAGIS Maize Products Exported per Marketing Year	23
Maize Crop Quality 2020/21 - summary of results	24 - 26
Percentage samples graded as Grade 1 over seasons (Graph 33)	24
South African maize crop quality 2020/21 (weighted averages) (Table 2)	27
RSA Production Regions	28
RSA Provinces (Figure 1)	28
RSA Crop Production Regions (Figure 2)	29
List of grain production regions with silo/intake stands and type of storage structure	30 - 33
Main maize producing provinces – comparison of results	34
RSA grading of white maize according to grade (Table 3)	35 - 39
RSA grading of white maize (Table 4)	40
RSA grading of yellow maize according to grade (Table 5)	41 - 44
RSA grading of yellow maize (Table 6)	45
Grading quality over 10 seasons (Table 7)	46
Percentage Defective kernels and Combined deviations over 10 seasons (Graphs 34 - 36)	47
USA grading of white maize (Table 8)	48 - 50
USA grading of yellow maize (Table 9)	51 - 53
RSA and USA Grading Regulations tables (Tables 10 and 11)	54
Physical quality characteristics of white maize according to grade (Table 12)	55 - 58
Physical quality characteristics of white maize (Table 13)	59
Physical quality characteristics of yellow maize according to grade (Table 14)	60 - 63
Physical quality characteristics of yellow maize (Table 15)	64
Physical quality characteristics over 10 seasons (Table 16)	65
Test weight (kg/hl) over 10 seasons (Graph 37)	65
100 Kernel mass and kernel size over 10 seasons (Graphs 38 - 40)	66
Roff milling and whiteness index of white maize according to grade (Table 17)	67 - 70
Roff milling and whiteness index of white maize (Table 18)	71

Roff milling fractions and total meal extraction (Graphs 41 – 46)	72
Nutritional values of white (Table 19) and yellow maize (Table 20) according to grade	73 - 76
Nutritional values of white and yellow maize (Table 21)	77 - 78
Nutritional values over 10 seasons (Table 22)	79
Fat, protein and starch content over 10 seasons (Graphs 47 - 49)	80
Presence of Genetically Modified Maize (Table 23)	81
Mycotoxin results discussion (Graphs 50 - 60)	82 - 87
International Mycotoxin Regulations	87
National Mycotoxin Regulations	88
Mycotoxin results for the 2020/21 season (Table 24)	89 - 101
Mycotoxin results summary from 2010/11 to 2020/21 (Table 25)	102
Methods	103 - 107
SANAS Certificate and Schedule of Accreditation	108 - 111
Grading Regulations for Maize, Government Notice No. R.473 of 8 May 2009	112 - 120
Industry-wide Dispensation REF NO: 20/4/14/1 of 15 April 2010	121 - 122

**Compiled and issued by:  
The Southern African Grain Laboratory NPC**

Grain Building - Agri-Hub Office  
477 Witherite Street  
The Willows  
Pretoria  
SOUTH AFRICA

PostNet Suite # 391  
Private Bag X 1  
The Willows  
0041



Tel: +27 (12) 807 4019  
Fax: +27(12) 807 4160  
E-mail: [info@sagl.co.za](mailto:info@sagl.co.za)  
Website: [www.sagl.co.za](http://www.sagl.co.za)

# South African

## COMMERCIAL MAIZE QUALITY 2020/2021



### ACKNOWLEDGMENTS


#### WITH GRATITUDE TO:

- The Maize Trust for financial support in conducting this survey.
- Agbiz Grain and its members for providing the samples to make this survey possible.
- The Crop Estimates Committee (CEC) of the Department of Agriculture, Land Reform and Rural Development (DALRRD) for providing production related figures.
- South African Grain Information Service (SAGIS) for providing supply and demand figures relating to maize and maize products.
- The Bureau for Food and Agricultural Policy (BFAP) for providing research based market analysis.

## Introduction

During the 2021 harvesting season, a representative sample of each delivery of maize at the various grain intake points was taken according to the prescribed grading regulation. The sampling procedure for the samples used in this survey is described on page 103. A total of 1 000 composite samples, representing white and yellow maize of each production region, were received and analysed to determine the quality. The samples consisted of 560 white and 440 yellow maize samples.

GRAPH 1: PROVINCIAL CONTRIBUTION TO THE PRODUCTION OF THE 2020/21 MAIZE CROP



Figures provided by the CEC.

The quality attributes which were tested for, include:

- RSA grading: Samples were graded according to the following factors, as defined in the South African grading regulation: defective kernels above and below the 6.35 mm sieve, total defective kernels, foreign matter, other colour kernels, combined deviations and pinked kernels.
- USA grading: Samples were graded according to the American Grading Regulations to determine the following factors: Test weight per bushel (pounds), heat damaged kernels, total damaged kernels, broken corn and foreign matter (BCFM) and other colour.
- Nutritional values: Moisture, crude protein, crude fat, crude fibre and starch.
- Physical Quality factors: Test weight (kg/hl), 100 kernel mass, kernel size, breakage susceptibility, stress cracks, milling index and grit yield.
- All white maize samples were milled on the Roff laboratory mill and the whiteness index of the maize meal determined.
- Mycotoxin analyses were performed on 350 samples representative of white and yellow maize produced per region.

Testing for the presence of a selection of traits present in Genetically Modified (GM) maize were performed on 100 samples representative of white and yellow maize produced per region.

Please refer to pages 103 - 107 for the methodologies followed.

The maize crop quality survey is performed annually by the Southern African Grain Laboratory NPC (SAGL). SAGL was established in 1997 on request of the Grain Industry. SAGL is an ISO 17025 accredited testing laboratory and participates in a number of proficiency testing schemes, both nationally and internationally, as part of our ongoing quality assurance procedures to demonstrate technical competency and international comparability.

The results of this, the 24<sup>th</sup> survey, as well as previous years' surveys are available on the SAGL website ([www.sagl.co.za](http://www.sagl.co.za)). The report, in an easy to page format, is also available on the website. Hard copy reports are distributed to industry stakeholders and interested parties.

In addition to the quality information, production figures (obtained from the Crop Estimates Committee (CEC)) relating to hectares planted, tons produced and yields obtained on a national as well as provincial basis, over an eleven season period, are provided in this report. SAGIS (South African Grain Information Service) supply and demand figures over several years are provided in table and graph format, also import and export data. Information on maize processed per province as well as the manufacture, import and export of maize products is also included in this report. The national grading regulations as published in the Government Gazette of 8 May 2009, are provided on pages 112 to 122.

The goal of this crop quality survey is to accumulate quality data on the commercial maize crop on a national level. This valuable data reveals general tendencies, highlights quality differences in the commercial maize produced in different local production regions and provides important information on the quality of commercial maize intended for export. During seasons when maize is imported for domestic use, the quality of the imported maize can also be compared to that of locally produced maize.

The Maize Trust investment in the annual Crop Quality Surveys, has created a unique and extremely useful database of crop quality measurements over several seasons and regions. Historically, the data has only been presented in table and graph format and has never been used for trend analyses or to assist in the development of prediction models such as the Milling Index Model.

In order to address this issue, SAGL undertook a data mining project, titled "Data Mining of past eleven years' Milling Index and Crop Survey Results", funded by the Maize Trust. A complete statistical analysis of the maize quality data from the 2001/2002 to 2011/2012 seasons were performed for the following measurements: Protein (crude), starch, fat (crude), hectolitre mass, 100 kernel mass, total deviations (grading data), Roff Milling Index, Break 1 flour yield, Break 2 flour yield, Break 3 flour yield, Grit yield and Bran yield (all Roff milling data). Data is added annually to this data set. The project outcome provides a decision-making tool to the maize industry stakeholders to assist in the identification of potential problem areas in maize quality and to focus future research activities. A number of data mining as well as quality related investigation/research projects have been based on results obtained from this valuable data base.


As part of the project, the possibility was explored to develop a Geographic Information System (GIS) map system, where grain production regions (with the boundaries illustrated) are presented on a map of South Africa. SIQ (with additional data from Agbiz Grain on the regional boundary specifications) created a software package based on an open source GIS package (QGIS). These GIS maps show mean values for a trait for a specific region as an average for all seasons combined or as individual seasons on a year to year basis. The results of the crop quality traits are represented in a colour scale format – highest to lowest values are indicated by the darkest to the lightest colour. Mean values are showed as a legend. This GIS tool provides a good starting point and can be optimised based on specific requirements.

# Production

The final figure for the 2020/21 season's commercial maize crop as overseen by the National Crop Estimates Liaison Committee (CELC) is 16 315 million tons, the second largest maize crop on record. This figure represents a year on year increase of 6.6% and is also 33.5% higher than the previous 10-year crop average (12 218 million tons). White maize's contribution to the total production was 8 600 million tons (52.7%) and that of yellow maize 7 715 million tons (47.3%).


The national Crop Estimates Committee's (CEC) estimated total production figures were revised, using the South African Grain Information Services' (SAGIS) published figures of actual deliveries as basis for the calculations. Figures from the maize utilisation survey, conducted by the Department of Agriculture, Land Reform and Rural Development (DALLRD) to determine on-farm usage and retentions as well as the telephonic survey conducted by the National Crop Statistics Consortium (NCSC), were added to the SAGIS delivery figures to calculate the final crop production figures.

GRAPH 2: MAIZE PRODUCTION IN RSA FROM 2010/11 TO 2020/21



The total area utilised for maize production in the 2020/21 season was 2 755 400 hectares, representing an increase of 5.5% compared to the previous season and which is also 10.2% higher than the previous 10-year average. White maize was planted on 1 691 900 hectares and yellow maize on 1 063 500 hectares (1 616 300 and 994 500 hectares respectively in the 2019/20 season).


GRAPH 3: TOTAL RSA AREA UTILISED FOR MAIZE PRODUCTION FROM 2010/11 TO 2020/21



The maize yield of 5.92 tons per hectare (t/ha) this season, was 1% higher than the previous season. The previous 10-year average is 4.87 t/ha. White maize yielded 5.08 t/ha and yellow maize 7.25 t/ha (5.29 t/ha and 6.79 t/ha respectively last season).



GRAPH 4: RSA MAIZE YIELD FROM 2010/11 TO 2020/21



The maize area planted in the non-commercial agricultural sector is estimated at 362 900 ha, representing a 22% increase compared to the 297 460 ha of the previous season. The expected maize crop is 636 440 tons for this sector, which is 17% more than last season. Approximately 53% of non-commercial maize is produced in the Eastern Cape, followed by KwaZulu-Natal with 23%.

Table 1: Maize production overview - 2020/21 season


Province	Type of production	White			Yellow		
		Hectares planted, ha	Production, tons	Yield, t/ha	Hectares planted, ha	Production, tons	Yield, t/ha
Western Cape	Dryland	-	-	-	-	-	-
	Irrigation	500	4 800	9.60	3 500	33 300	9.51
	Total	500	4 800	9.60	3 500	33 300	9.51
Northern Cape	Dryland	-	-	-	-	-	-
	Irrigation	3 400	40 300	11.85	41 000	627 300	15.30
	Total	3 400	40 300	11.85	41 000	627 300	15.30
Free State	Dryland	890 500	4 322 000	4.85	387 000	2 170 000	5.61
	Irrigation	17 000	170 000	10.00	33 000	372 000	11.27
	Total	907 500	4 492 000	4.95	420 000	2 542 000	6.05
Eastern Cape	Dryland	4 200	18 300	4.36	12 700	68 800	5.42
	Irrigation	1 800	20 700	11.50	5 300	64 400	12.15
	Total	6 000	39 000	6.50	18 000	133 200	7.40
KwaZulu-Natal	Dryland	41 500	219 000	5.28	30 500	203 500	6.67
	Irrigation	8 500	91 000	10.71	24 500	247 500	10.10
	Total	50 000	310 000	6.20	55 000	451 000	8.20
Mpumalanga	Dryland	156 000	998 800	6.40	340 000	2 595 000	7.63
	Irrigation	9 000	101 700	11.30	20 000	225 000	11.25
	Total	165 000	1 100 500	6.67	360 000	2 820 000	7.83
Limpopo	Dryland	7 500	22 200	2.96	8 500	28 900	3.40
	Irrigation	9 000	90 000	10.00	12 500	143 300	11.46
	Total	16 500	112 200	6.80	21 000	172 200	8.20
Gauteng	Dryland	54 500	336 000	6.17	42 500	301 000	7.08
	Irrigation	3 500	35 200	10.06	7 500	79 000	10.53
	Total	58 000	371 200	6.40	50 000	380 000	7.60
North West	Dryland	471 500	2 008 000	4.26	82 300	419 500	5.10
	Irrigation	13 500	122 000	9.04	12 700	136 500	10.75
	Total	485 000	2 130 000	4.39	95 000	556 000	5.85
RSA	Dryland	1 625 700	7 924 300	4.87	903 500	5 786 700	6.40
	Irrigation	66 200	675 700	10.21	160 000	1 928 300	12.05
	Total	1 691 900	8 600 000	5.08	1 063 500	7 715 000	7.25

Figures provided by the CEC.


The major commercial maize-producing provinces are the Free State, Mpumalanga and North West, contributing 84% of the total maize production in the RSA. The Free State produced 7 034 000 tons of maize on 1 327 500 hectares with a yield of 5.30 t/ha. Mpumalanga produced 3 920 500 tons of maize on 525 000 hectares with a yield of 7.47 t/ha and North West harvested 2 686 000 tons of maize on 580 000 hectares yielding 4.63 t/ha. Yellow maize contributed 72% of the total maize production in Mpumalanga while the majority of maize produced in the Free State (64%) and North West (79%) was white.

Please see graphs 5 to 7 for provincial figures of area planted, production and yield over the last three seasons.


GRAPH 5: AREA UTILISED FOR MAIZE PRODUCTION PER PROVINCE OVER THREE SEASONS



GRAPH 6: MAIZE PRODUCTION PER PROVINCE OVER THREE SEASONS




GRAPH 7: MAIZE YIELD PER PROVINCE OVER THREE SEASONS




Figures provided by the CEC.

Graphs 8 to 13 provide an overview of the area planted and production figures for the Free State, Mpumalanga and North West from the 2010/11 to 2020/21 seasons.


**GRAPH 8: AREA UTILISED FOR MAIZE PRODUCTION IN THE FREE STATE SINCE 2010/11**




**GRAPH 9: MAIZE PRODUCTION IN THE FREE STATE SINCE 2010/11**




**GRAPH 10: AREA UTILISED FOR MAIZE PRODUCTION IN MPUMALANGA SINCE 2010/11**




**GRAPH 11: MAIZE PRODUCTION IN MPUMALANGA SINCE 2010/11**



**GRAPH 12: AREA UTILISED FOR MAIZE PRODUCTION IN NORTH WEST SINCE 2010/11**



**GRAPH 13: MAIZE PRODUCTION IN NORTH WEST SINCE 2010/11**



Figures provided by the CEC.

## Supply and Demand

World maize production for the 2020/21 season was estimated at 1 131.5 million tons according to the *International Grains Council Grain Market Report GMR 532 – 19 May 2022*, with the major maize producing countries being the USA, China and Brazil. The USA, Argentina, Brazil and the Ukraine are the biggest exporters of maize. Maize usage figures are estimated at 132.0, 298.0 and 678.6 million tons respectively for food, industrial and feed purposes. World production for the 2021/22 season is forecasted at 1 213.8 million tons and the 2022/23 figure is projected to be 1 183.8 million tons.

According to *The Bureau for Food and Agricultural Policy (BFAP) Baseline, Agricultural Outlook 2021 – 2030*, demand growth prospects for the various summer crops diverge due to differences in use and as a result also fundamentally different drivers affecting markets. While staple grains such as white maize and sorghum is predominantly consumed as food, the bulk of yellow maize is consumed as primary energy source in most animal feed rations.

After declining over the past decade, per capita consumption of maize is projected to increase by an average of 0.5% per annum over the next 10 years. This increase, in conjunction with a growing population, supports growth of 12% in white maize for human consumption by 2030 relative to the 2018-2020 base period. Relative prices dictate that a smaller share of white maize will be consumed as animal feed by 2030, compared to the base period.

Despite slower growth demand for animal protein in South Africa, the commitments made by the poultry Masterplan, which should result in some import replacement and consequently a decline in the share of imported products in domestic consumption, combined with export led expansion in the beef sector, still imply substantial growth in the demand for animal feed over the coming decade. Consequently, yellow maize consumption as animal feed is projected to rise by 34% over the next 10 years. In excess of a million tons of white maize is set to be utilised as animal feed by 2030.

White maize area increased sharply over the last two years and is projected to remain firm in the short term, before returning to the longer term trend and decline from 2023 onwards. The short term gains are nevertheless such that by 2030, white maize area will be similar to the levels planted on average between 2018 and 2020. Yield gains of 24% over the same period are sufficient to support the projected demand growth. The area cultivated to yellow maize however, continues to increase, expanding by 8% over the 10-year period to 2030.


The percentage change in area and yield for the major summer crops, comparing 2030 to the 2018-2020 base period, reflects fairly consistent yield gains based on continuous improvements in cultivar technology, as well as consistent evolution of production practices and area dynamics. Relative to the base period, white maize yields are expected to improve by 24% by 2030, largely due to technological gains. Ample production for the local market and an exportable surplus to neighbouring countries are thus provided. Yellow maize yield gains are a bit more subdued than white maize, owing to further area expansion, but are still expected to improve by 20% over the next decade.

Maize production growth over the outlook period is projected to be sufficient to sustain domestic demand and yield a consistent exportable surplus. This surplus will fluctuate in line with weather dynamics, but in normal years, white maize exports are expected to stabilise below a million tons.

White maize is mostly exported into the Southern African region, where South Africa is facing increasing competition. Zambia for example produces non-GM white maize. Yellow maize exports are projected to increase over the outlook period since it is easier to trade in the global market. This increase in exports is however less than the increase in production.

Local Supply and Demand figures, compiled by SAGIS, are provided in the graph below and in tables and graphs on pages 9 to 14.

GRAPH 14: MAIZE SUPPLY AND DEMAND OVERVIEW 2021/22 MARKETING SEASON



Information provided by SAGIS.




**TOTAL MAIZE: SUPPLY AND DEMAND TABLE BASED ON SAGIS' INFO (TON)**

Publication date: 2022-04-28


Season	Marketing Season (May - Apr)												Current		10 Year average					
	03/04	04/05	05/06	06/07	07/08	08/09	09/10	10/11	11/12	12/13	13/14	14/15	15/16	16/17		17/18	18/19	19/20	20/21	2021/22
	***																			
CEC (Crop Estimate)	9 392 000	9 482 000	11 450 000	6 618 000	7 125 000	12 700 000	12 050 000	12 815 000	10 360 000	12 120 656	11 810 600	14 250 000	9 955 000	7 776 500	16 820 000	12 510 000	11 275 000	15 300 000	16 315 000	12 217 976
CEC (Retention)	366 000	410 000	754 000	480 000	337 000	554 000	389 000	527 000	474 000	433 000	457 810	550 000	0	0	0	0	0	0	0	181 481
<b>SUPPLY</b>																				
Opening stock (1 May)	2 710 000	2 624 000	3 148 000	3 169 000	2 070 000	1 049 000	1 581 000	2 131 000	2 336 000	994 000	1 417 393	569 028	2 073 635	2 471 067	1 094 638	3 689 476	2 663 086	1 000 601	2 116 906	1 832 892
Prod deliveries*	8 409 000	9 093 000	10 055 000	6 707 000	6 882 000	11 899 000	11 629 000	12 016 000	10 340 000	11 929 000	10 981 995	13 827 632	9 794 332	7 469 600	15 628 682	11 983 852	10 887 053	15 278 983	15 033 375	11 813 113
Imports	441 000	219 000	380 000	931 000	1 120 000	27 000	27 000	0	421 000	11 000	79 682	65 250	1 983 610	2 238 743	0	171 622	509 884	463	7 583	545 905
Surplus	40 000	0	0	32 000	29 000	30 000	68 000	77 000	54 000	42 000	122 608	26 153	52 930	44 417	46 657	22 173	22 336	20 079	42 480	45 335
<b>Total Supply</b>	<b>11 600 000</b>	<b>11 936 000</b>	<b>13 563 000</b>	<b>10 839 000</b>	<b>10 101 000</b>	<b>13 005 000</b>	<b>13 305 000</b>	<b>14 224 000</b>	<b>13 151 000</b>	<b>12 976 000</b>	<b>12 611 678</b>	<b>14 508 063</b>	<b>13 884 507</b>	<b>12 221 827</b>	<b>16 769 977</b>	<b>15 867 123</b>	<b>14 087 764</b>	<b>16 300 126</b>	<b>17 200 344</b>	<b>14 030 194</b>
<b>DEMAND</b>																				
Processed	7 243 000	7 283 000	7 462 000	7 660 000	8 025 000	8 615 000	8 658 000	8 857 000	8 941 000	8 985 000	9 348 670	9 926 519	10 248 994	9 838 709	10 289 680	10 689 977	11 106 412	11 201 202	10 216 814	10 063 716
-human	3 712 000	3 740 000	3 825 000	3 816 000	3 809 000	4 524 000	4 471 000	4 513 000	4 512 000	4 499 000	4 582 310	4 840 021	4 898 482	4 809 221	4 983 476	5 160 772	5 387 572	5 657 836	4 780 162	4 914 069
-animal/industrial	3 416 000	3 427 000	3 537 000	3 763 000	4 157 000	4 020 000	4 101 000	4 271 000	4 362 000	4 378 000	4 715 295	5 040 647	5 520 248	5 003 810	5 276 447	5 507 180	5 688 317	5 527 649	5 420 451	5 102 959
-gristing	115 000	116 000	100 000	81 000	63 000	69 000	86 000	73 000	67 000	58 000	51 065	45 851	30 264	25 678	29 757	23 025	20 523	15 717	16 211	36 888
-bio-fuel	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Withdrawn by producers	299 000	255 000	315 000	241 000	217 000	273 000	291 000	267 000	142 000	138 000	148 909	124 508	76 888	94 948	102 906	64 264	57 104	35 736	33 727	98 526
Released to end-consumers	224 000	351 000	340 000	235 000	230 000	220 000	378 000	526 000	484 000	478 000	280 432	205 577	186 296	157 460	180 544	151 643	99 815	69 329	46 705	229 310
Net receipts (-)/disp(+)	25 000	18 000	28 000	36 000	42 000	49 000	51 000	44 000	15 000	62 000	12 043	22 100	21 451	9 770	15 663	13 095	8 654	9 163	5 197	18 894
Deficit	0	49 000	12 000	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<b>Total Exports</b>	<b>1 165 000</b>	<b>832 000</b>	<b>2 237 000</b>	<b>597 000</b>	<b>534 000</b>	<b>2 269 000</b>	<b>1 796 000</b>	<b>2 194 000</b>	<b>2 575 000</b>	<b>1 946 000</b>	<b>2 232 596</b>	<b>2 155 724</b>	<b>879 611</b>	<b>1 026 302</b>	<b>2 461 708</b>	<b>2 284 056</b>	<b>1 809 573</b>	<b>2 867 790</b>	<b>3 854 215</b>	<b>2 025 856</b>
Products	89 000	100 000	94 000	49 000	62 000	107 000	126 000	128 000	129 000	133 000	176 978	188 319	166 383	189 112	192 874	219 592	360 812	320 926	363 391	210 100
African Countries	34 000	48 000	56 000	28 000	35 000	67 000	87 000	84 000	86 000	95 000	123 040	137 742	132 900	144 229	117 797	142 117	320 754	282 678	324 223	188 226
Other Countries	55 000	52 000	38 000	21 000	27 000	40 000	39 000	44 000	43 000	38 000	53 938	60 577	53 483	44 883	75 077	71 475	40 058	36 248	39 168	51 874
Whole maize	1 096 000	732 000	2 143 000	548 000	472 000	2 162 000	1 670 000	2 066 000	2 446 000	1 813 000	2 055 618	1 957 405	693 428	837 190	2 288 834	2 070 466	1 448 761	2 546 864	3 490 824	1 815 757
Border Posts	950 000	591 000	1 311 000	488 000	472 000	1 332 000	703 000	629 000	584 000	613 000	921 454	691 659	684 834	804 322	591 692	630 572	1 230 762	1 380 017	703 917	813 231
Harbours	146 000	141 000	832 000	60 000	0	830 000	967 000	1 437 000	1 862 000	1 200 000	1 134 164	1 264 326	8 594	32 868	1 697 142	1 439 894	217 959	1 166 847	2 786 907	1 002 383
<b>Total Demand</b>	<b>8 976 000</b>	<b>8 788 000</b>	<b>10 394 000</b>	<b>8 169 000</b>	<b>9 052 000</b>	<b>11 424 000</b>	<b>11 174 000</b>	<b>11 888 000</b>	<b>12 157 000</b>	<b>11 569 000</b>	<b>12 022 650</b>	<b>12 434 428</b>	<b>11 413 440</b>	<b>11 127 169</b>	<b>13 080 501</b>	<b>13 204 037</b>	<b>13 087 163</b>	<b>14 183 220</b>	<b>14 156 659</b>	<b>12 426 863</b>
Ending Stock (30 Apr)	2 824 000	3 148 000	3 169 000	2 070 000	1 849 000	1 581 000	2 151 000	2 336 000	994 000	1 417 000	589 028	2 073 635	2 871 067	1 094 638	3 689 476	2 663 086	1 000 601	2 116 906	3 043 686	1 810 944
- processed / month	603 600	606 900	621 900	638 300	669 100	717 800	721 500	738 100	745 100	744 565	779 056	827 210	854 083	819 892	866 307	889 915	925 554	935 434	928 801	837 811
- months' stock	4.3	5.2	5.1	3.2	1.6	2.2	3.0	3.2	1.3	1.9	0.8	2.5	2.9	1.3	4.3	3.0	1.1	2.3	3.3	2

Note: \*\*\* Figures for current season up to date


GRAPH 15: MAIZE: CEC ESTIMATE, RETENTIONS AND SAGIS DELIVERIES OVER 10 MARKETING SEASONS




GRAPH 16: MAIZE: IMPORTS AND EXPORTS OVER 10 MARKETING SEASONS



GRAPH 17: MAIZE: RSA CONSUMPTION OVER 10 MARKETING SEASONS



GRAPH 18: MAIZE: OPENING AND ENDING STOCKS OVER 10 MARKETING SEASONS



**WHITE MAIZE: SUPPLY AND DEMAND TABLE BASED ON SAGIS' INFO (TON)**


Publication date: 2022-04-28

Season	Marketing Season (May -Apr)												Current Season May - Mar	10 Year average							
	03/04	04/05	05/06	06/07	07/08	08/09	09/10	10/11	11/12	12/13	13/14	14/15			15/16	16/17	17/18	18/19	19/20	20/21	21/22
	***																				
CEC (Crop Estimate)	6 366 000	5 805 000	6 541 000	4 187 000	4 315 000	7 480 000	6 775 000	7 830 000	6 052 000	6 903 656	5 606 900	7 710 000	4 735 000	3 408 500	9 916 000	6 540 000	5 545 000	8 547 500	8 600 000		
CEC (Retention)	116 000	113 000	184 000	144 000	11 000	120 000	83 000	119 000	100 000	114 000	110 910	150 000	0	0	0	0	0	0	0		
<b>SUPPLY</b>																					
Opening stock (1 May)	1 718 000	2 123 000	2 402 000	2 301 000	1 630 000	618 000	762 000	1 362 000	1 609 000	519 000	757 214	274 318	1 282 551	1 307 867	597 837	2 428 653	1 799 998	473 964	1 354 953		
Prod deliveries*	5 845 000	5 647 000	6 108 000	4 392 000	4 309 000	7 190 000	6 737 000	7 518 000	6 105 000	6 860 000	5 342 204	7 592 893	4 808 279	3 551 822	9 268 583	6 308 941	5 442 474	8 606 334	8 600 297		
Imports	33 000	0	0	1 000	46 000	0	0	0	133 000	11 000	0	0	100 803	644 144	0	0	0	0	7 583		
Surplus	40 000	0	4 000	20 000	19 000	25 000	48 000	45 000	18 000	22 000	8 808	8 808	17 474	31 994	21 751	1 403	0	11 215	24 307		
<b>Total Supply</b>	<b>7 636 000</b>	<b>7 770 000</b>	<b>8 514 000</b>	<b>6 714 000</b>	<b>6 004 000</b>	<b>7 833 000</b>	<b>7 847 000</b>	<b>8 925 000</b>	<b>7 865 000</b>	<b>7 431 000</b>	<b>6 169 277</b>	<b>7 876 019</b>	<b>6 209 137</b>	<b>5 535 827</b>	<b>9 888 181</b>	<b>8 738 997</b>	<b>7 241 472</b>	<b>9 081 513</b>	<b>9 447 140</b>		
<b>DEMAND</b>																					
Processed	4 212 000	4 313 000	4 186 000	4 385 000	4 751 000	4 922 000	4 555 000	5 871 000	5 374 000	5 047 000	4 808 674	5 862 438	4 319 897	4 331 787	6 533 966	6 283 320	5 449 415	6 410 756	6 557 702		
-human	3 467 000	3 478 000	3 559 000	3 526 000	3 552 000	4 198 000	4 125 000	4 157 000	4 119 000	4 095 000	4 118 448	4 361 295	4 183 067	4 232 583	4 459 504	4 594 123	4 805 569	5 073 886	4 343 738		
-animal/industrial	641 000	733 000	543 000	787 000	1 142 000	662 000	382 000	1 698 000	1 202 000	904 000	651 925	1 469 002	118 922	86 153	2 061 649	1 677 236	629 076	1 325 959	2 202 691		
-gristing	104 000	102 000	84 000	72 000	57 000	62 000	68 000	56 000	53 000	49 000	38 301	32 141	18 106	13 051	12 813	11 961	10 770	10 911	11 273		
-bio-fuel	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
Withdrawn by producers	144 000	107 000	101 000	112 000	107 000	111 000	81 000	108 000	46 000	36 000	32 409	38 940	13 365	14 063	35 885	12 844	13 111	10 069	13 154		
Released to end-consumers	76 000	181 000	71 000	80 000	69 000	45 000	62 000	189 000	126 000	95 000	43 000	38 934	13 987	5 660	30 125	22 946	17 649	5 827	3 722		
Net receipts(-)/disps(+)	12 000	17 000	11 000	27 000	28 000	27 000	10 000	22 000	7 000	28 000	1 953	14 319	-2 862	- 963	7 583	4 238	6 282	5 413	424		
Deficit	0	38 000	0	0	0	0	0	0	0	0	0	0	0	0	0	0	5 605	0	0		
Total Exports	1 069 000	712 000	1 844 000	480 000	451 000	1 965 000	1 477 000	1 126 000	1 794 000	1 468 000	1 008 923	640 807	557 063	587 423	851 969	616 651	1 275 446	1 304 475	855 778		
Products	65 000	44 000	58 000	20 000	31 000	69 000	69 000	77 000	60 000	68 000	82 877	93 307	83 636	41 042	42 038	72 280	236 537	182 824	167 591		
African Countries	22 000	23 000	51 000	14 000	24 000	57 000	58 000	62 000	47 000	56 000	72 032	77 890	73 061	36 573	40 695	70 726	236 162	182 297	165 102		
Other Countries	43 000	21 000	7 000	6 000	7 000	12 000	11 000	15 000	13 000	12 000	10 845	15 377	10 575	4 469	1 343	1 554	375	527	2 489		
Whole maize	1 004 000	688 000	1 786 000	460 000	400 000	1 897 000	1 409 000	1 049 000	1 734 000	1 400 000	926 046	547 500	473 427	546 381	809 931	544 371	1 039 909	1 121 651	688 187		
Border Posts	881 000	527 000	1 210 000	400 000	400 000	1 241 000	566 000	509 000	439 000	462 000	538 128	473 427	473 427	520 200	417 327	397 657	836 596	953 736	513 805		
Harbours	123 000	141 000	576 000	60 000	60 000	656 000	842 000	540 000	1 295 000	938 000	198 957	9 372	0	26 181	392 604	146 714	202 313	167 915	174 382		
<b>Total Demand</b>	<b>5 613 000</b>	<b>5 368 000</b>	<b>6 215 000</b>	<b>5 084 000</b>	<b>5 386 000</b>	<b>7 071 000</b>	<b>6 185 000</b>	<b>7 316 000</b>	<b>7 347 000</b>	<b>6 674 000</b>	<b>6 693 438</b>	<b>4 901 270</b>	<b>4 937 990</b>	<b>4 937 990</b>	<b>7 459 528</b>	<b>6 939 999</b>	<b>6 767 508</b>	<b>7 736 560</b>	<b>7 430 780</b>		
Ending Stock (30 Apr)	2 123 000	2 402 000	2 301 000	1 630 000	618 000	762 000	1 362 000	1 609 000	519 000	757 214	274 318	1 282 551	1 307 867	597 837	2 428 653	1 799 998	473 964	1 354 953	2 216 360		
- processed	351 000	359 400	348 800	365 400	395 900	410 200	379 600	489 300	447 800	420 583	400 723	488 537	359 975	360 982	544 497	523 610	454 118	534 200	596 155		
- months' stock	6.0	6.7	6.6	4.5	6.6	1.9	3.6	3.3	1.2	1.8	0.7	2.6	3.6	1.7	4.5	3.4	1.0	2.5	3.4		


Note: \*\*\* Figures for current season up to date




GRAPH 19: WHITE MAIZE: CEC ESTIMATE, RETENTIONS AND SAGIS DELIVERIES OVER 10 MARKETING SEASONS




GRAPH 20: WHITE MAIZE: IMPORTS AND EXPORTS OVER 10 MARKETING SEASONS



GRAPH 21: WHITE MAIZE: RSA CONSUMPTION OVER 10 MARKETING SEASONS



GRAPH 22: WHITE MAIZE: OPENING AND ENDING STOCKS OVER 10 MARKETING SEASONS



Information provided by SAGIS.


**YELLOW MAIZE: SUPPLY AND DEMAND TABLE BASED ON SAGIS' INFO (TON)**

Publication date: 2022-04-28


Season	Marketing Season (May - Apr)												Current		10 Year average					
	03/04	04/05	05/06	06/07	07/08	08/09	09/10	10/11	11/12	12/13	13/14	14/15	15/16	16/17		17/18	18/19	19/20	20/21	21/22
	May-Mar																			
CEC (Crop Estimate)	3 026 000	3 677 000	4 909 000	2 431 000	2 810 000	5 220 000	5 275 000	4 985 000	4 308 000	5 217 000	6 203 800	6 540 000	5 220 000	4 370 000	6 904 000	5 970 000	5 730 000	6 752 500	7 715 000	
CEC (Retention)	250 000	297 000	570 000	336 000	326 000	434 000	306 000	408 000	374 000	319 000	346 800	400 000	0	0	0	0	0	0	0	
<b>SUPPLY</b>																				
Opening stock (1 May)	992 000	501 000	746 000	868 000	440 000	431 000	819 000	769 000	727 000	476 000	660 179	314 710	791 054	1 163 200	496 801	1 260 823	864 088	526 637	761 953	
Prod deliveries*	2 564 000	3 446 000	3 947 000	2 315 000	2 573 000	4 709 000	4 892 000	4 498 000	4 235 000	5 049 000	5 649 791	6 234 739	4 986 053	3 917 778	6 360 089	5 674 911	5 444 579	6 672 649	6 973 078	
Imports	408 000	219 000	360 000	830 000	1 074 000	27 000	27 000	0	288 000	0	79 682	65 250	1 862 807	1 592 599	0	171 622	509 684	463	0	
Surplus	0	0	0	12 000	10 000	5 000	20 000	32 000	36 000	20 000	52 749	17 345	35 456	12 423	24 906	20 770	27 941	8 864	18 173	
<b>Total Supply</b>	<b>3 964 000</b>	<b>4 166 000</b>	<b>4 125 000</b>	<b>4 053 000</b>	<b>4 087 000</b>	<b>5 172 000</b>	<b>5 756 000</b>	<b>5 299 000</b>	<b>5 286 000</b>	<b>5 545 000</b>	<b>6 442 401</b>	<b>6 652 044</b>	<b>7 675 370</b>	<b>6 686 000</b>	<b>6 887 796</b>	<b>7 128 126</b>	<b>6 646 292</b>	<b>7 208 613</b>	<b>7 753 204</b>	
<b>DEMAND</b>																				
Processed	3 031 000	2 970 000	3 276 000	3 275 000	3 278 000	3 691 000	4 103 000	2 985 000	3 567 000	3 888 000	4 539 996	4 064 081	5 929 297	5 506 922	3 765 714	4 407 657	5 656 997	4 790 446	3 659 112	
-human	245 000	262 000	266 000	290 000	257 000	326 000	346 000	356 000	393 000	404 000	463 862	478 726	515 415	576 638	533 972	566 649	576 003	583 950	436 414	
-animal/industrial	2 775 000	2 684 000	2 976 000	2 976 000	3 015 000	3 365 000	3 739 000	2 613 000	3 160 000	3 474 000	4 069 370	3 571 645	5 401 726	4 917 657	3 214 798	3 829 944	5 069 241	4 201 690	3 217 760	
-gristing	11 000	14 000	16 000	9 000	6 000	7 000	18 000	17 000	14 000	10 000	12 764	13 710	12 156	12 627	16 944	11 064	9 753	4 806	4 938	
-bio-fuel	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Withdrawn by producers	155 000	148 000	214 000	129 000	110 000	162 000	210 000	159 000	96 000	102 000	116 500	87 568	63 503	80 865	67 021	51 420	43 993	25 647	20 573	
Released to end-consumers	148 000	170 000	269 000	155 000	161 000	175 000	316 000	337 000	358 000	363 000	237 432	166 643	172 309	151 800	150 419	128 697	82 166	63 502	42 963	
Net receipts(-)/disp(+)	13 000	1 000	17 000	9 000	14 000	22 000	41 000	22 000	8 000	34 000	10 090	7 761	24 313	10 733	8 060	8 857	2 372	3 750	4 773	
Deficit	0	11 000	16 000	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Total Exports	116 000	120 000	393 000	117 000	103 000	303 000	319 000	1 068 000	761 000	478 000	1 225 673	1 514 917	322 748	438 679	1 629 739	1 667 407	534 127	1 563 315	2 998 437	
Products	24 000	56 000	36 000	29 000	31 000	38 000	57 000	51 000	69 000	65 000	94 101	105 912	102 747	148 070	150 836	141 312	124 275	138 102	195 800	
African Countries	12 000	25 000	5 000	14 000	11 000	10 000	29 000	22 000	39 000	39 000	51 008	59 812	59 839	40 414	77 102	71 391	84 592	100 381	159 121	
Other Countries	12 000	31 000	31 000	15 000	20 000	28 000	28 000	29 000	30 000	26 000	43 093	45 200	42 908	40 414	73 734	69 921	39 693	37 721	36 679	
Whole maize	92 000	64 000	357 000	88 000	72 000	265 000	262 000	1 017 000	712 000	413 000	1 129 572	1 409 905	220 001	290 909	1 478 903	1 526 085	409 852	1 425 213	2 802 637	
Border Posts	66 000	64 000	101 000	88 000	72 000	81 000	137 000	120 000	145 000	151 000	193 465	153 331	211 407	284 122	174 365	232 915	394 166	426 281	190 112	
Harbours	23 000	0	256 000	0	0	174 000	125 000	897 000	567 000	262 000	936 107	1 254 854	8 594	6 687	1 304 538	1 293 180	15 686	998 932	2 612 525	
<b>Total Demand</b>	<b>3 463 000</b>	<b>3 420 000</b>	<b>4 165 000</b>	<b>3 665 000</b>	<b>3 666 000</b>	<b>4 353 000</b>	<b>4 989 000</b>	<b>4 572 000</b>	<b>4 810 000</b>	<b>4 885 000</b>	<b>6 127 691</b>	<b>5 840 990</b>	<b>6 512 170</b>	<b>6 189 199</b>	<b>5 620 973</b>	<b>6 264 038</b>	<b>6 319 655</b>	<b>6 446 660</b>	<b>6 725 878</b>	
Ending Stock (30 Apr)	501 000	746 000	865 000	440 000	451 000	619 000	769 000	727 000	476 000	660 000	314 710	791 054	1 163 200	496 801	1 260 823	864 088	526 637	761 953	1 027 526	
- processed p/month	252 600	247 500	273 000	272 900	273 200	307 600	341 900	246 800	297 300	324 000	376 333	338 673	494 108	458 910	313 810	367 305	471 416	399 204	332 647	
- months' stock	2.0	3.0	3.2	1.6	1.6	2.7	2.2	2.9	1.6	2.0	0.8	2.3	2.4	1.1	4.0	2.4	1.1	1.9	3.1	

Note: \*\*\* Figures for current season up to date


GRAPH 23: YELLOW MAIZE: CEC ESTIMATE, RETENTIONS AND SAGIS DELIVERIES OVER 10 MARKETING SEASONS




GRAPH 24: YELLOW MAIZE: IMPORTS AND EXPORTS OVER 10 MARKETING SEASONS



GRAPH 25: YELLOW MAIZE: RSA CONSUMPTION OVER 10 MARKETING SEASONS



GRAPH 26: YELLOW MAIZE: OPENING AND ENDING STOCKS OVER 10 MARKETING SEASONS



Information provided by SAGIS.


### WHITE MAIZE EXPORTS/IMPORTS

2021/22 MARKETING SEASON (1 May 2021 to 29 April 2022)

RSA EXPORTS		IMPORTS FOR RSA		IMPORTS FOR OTHER COUNTRIES		EXPORTS OF IMPORTED MAIZE		IMPORTS PER HARBOUR		EXPORTS PER HARBOUR	
Angola	17 264	Zambia	7 583	-	-	-	-	-	-	Durban	174 382
Botswana	247 013	-	-	-	-	-	-	-	-	-	-
Eswatini (Swaziland)	45 180	-	-	-	-	-	-	-	-	-	-
Italy	174 382	-	-	-	-	-	-	-	-	-	-
Lesotho	51 700	-	-	-	-	-	-	-	-	-	-
Mozambique	73 588	-	-	-	-	-	-	-	-	-	-
Namibia	120 100	-	-	-	-	-	-	-	-	-	-
Zimbabwe	4 616	-	-	-	-	-	-	-	-	-	-
<b>Total</b>	<b>733 843</b>	<b>Total</b>	<b>7583</b>	<b>Total</b>	<b>0</b>	<b>Total</b>	<b>0</b>	<b>Total*</b>	<b>0</b>	<b>Total</b>	<b>174 382</b>

\* Includes imports for RSA and Other Countries

GRAPH 27: MAJOR DESTINATIONS FOR RSA WHITE MAIZE EXPORTED DURING THE 2021/22 MARKETING SEASON




**YELLOW MAIZE EXPORTS/IMPORTS**

**2021/22 MARKETING SEASON (1 May 2021 to 29 April 2022)**

RSA EXPORTS		IMPORTS FOR RSA		IMPORTS FOR OTHER COUNTRIES		EXPORTS OF IMPORTED MAIZE		IMPORTS PER HARBOUR		EXPORTS PER HARBOUR	
Angola	16 448	-	-	-	-	-	-	-	-	Durban	2 787 908
Botswana	11 502	-	-	-	-	-	-	-	-	-	-
Eswatini (Swaziland)	102 226	-	-	-	-	-	-	-	-	-	-
Ghana	9 505	-	-	-	-	-	-	-	-	-	-
Italy	27 150	-	-	-	-	-	-	-	-	-	-
Japan	824 823	-	-	-	-	-	-	-	-	-	-
Korea, Rep of	430 395	-	-	-	-	-	-	-	-	-	-
Lesotho	325	-	-	-	-	-	-	-	-	-	-
Mozambique	43 409	-	-	-	-	-	-	-	-	-	-
Namibia	44 373	-	-	-	-	-	-	-	-	-	-
Saudi Arabia	1 665	-	-	-	-	-	-	-	-	-	-
Seychelles	769	-	-	-	-	-	-	-	-	-	-
Spain	103 410	-	-	-	-	-	-	-	-	-	-
Taiwan, Prov of China	880 319	-	-	-	-	-	-	-	-	-	-
Vietnam	494 783	-	-	-	-	-	-	-	-	-	-
Zimbabwe	27	-	-	-	-	-	-	-	-	-	-
<b>Total</b>	<b>2 991 129</b>	<b>Total</b>	<b>0</b>	<b>Total</b>	<b>0</b>	<b>Total</b>	<b>0</b>	<b>Total*</b>	<b>0</b>	<b>Total</b>	<b>2 787 908</b>

\* Includes imports for RSA and Other Countries

**GRAPH 28: MAJOR DESTINATIONS FOR RSA YELLOW MAIZE EXPORTED DURING THE 2021/22 MARKETING SEASON**



**TOTAL WHOLE MAIZE PROCESSED PER PROVINCE**

**PROGRESSIVE: May 2017 to April 2018 (Full 2017/18 Marketing Year)**

Processed For	Cape Provinces	Free State	Kwazulu-Natal	Mpumalanga	Limpopo	Gauteng	Northwest	Total
Human Consumption and Gristing	285 475	1 614 961	356 881	591 932	426 137	857 317	967 932	5 100 635
Animal Feed and Industrial	1 445 836	712 710	456 876	722 148	127 893	1 390 634	535 822	5 391 919
	<b>1 731 311</b>	<b>2 327 671</b>	<b>813 757</b>	<b>1 314 080</b>	<b>554 030</b>	<b>2 247 951</b>	<b>1 503 754</b>	<b>10 492 554</b>

**PROGRESSIVE: May 2018 to April 2019 (Full 2018/19 Marketing Year)**

Processed For	Cape Provinces	Free State	Kwazulu-Natal	Mpumalanga	Limpopo	Gauteng	Northwest	Total
Human Consumption and Gristing	252 646	1 756 439	319 708	641 477	395 583	891 486	1 030 218	5 287 557
Animal Feed and Industrial	1 461 593	736 532	473 837	780 575	141 674	1 446 615	576 186	5 617 012
	<b>1 714 239</b>	<b>2 492 971</b>	<b>793 545</b>	<b>1 422 052</b>	<b>537 257</b>	<b>2 338 101</b>	<b>1 606 404</b>	<b>10 904 569</b>

**PROGRESSIVE: May 2019 to April 2020 (Full 2019/20 Marketing Year)**

Processed For	Cape Provinces	Free State	Kwazulu-Natal	Mpumalanga	Limpopo	Gauteng	Northwest	Total
Human Consumption and Gristing	257 441	1 872 605	338 758	748 908	399 058	911 924	1 161 924	5 690 618
Animal Feed and Industrial	1 533 217	760 770	498 222	792 017	131 238	1 437 789	623 353	5 776 606
	<b>1 790 658</b>	<b>2 633 375</b>	<b>836 980</b>	<b>1 540 925</b>	<b>530 296</b>	<b>2 349 713</b>	<b>1 785 277</b>	<b>11 467 224</b>

**PROGRESSIVE: May 2020 to April 2021 (Full 2020/21 Marketing Year)**

Processed For	Cape Provinces	Free State	Kwazulu-Natal	Mpumalanga	Limpopo	Gauteng	Northwest	Total
Human Consumption and Gristing	259 782	2 013 461	402 528	777 393	399 818	920 505	1 160 019	5 933 506
Animal Feed and Industrial	1 387 012	728 135	503 310	849 138	139 147	1 392 472	589 408	5 588 622
	<b>1 646 794</b>	<b>2 741 596</b>	<b>905 838</b>	<b>1 626 531</b>	<b>538 965</b>	<b>2 312 977</b>	<b>1 749 427</b>	<b>11 522 128</b>

**PROGRESSIVE: May 2021 to April 2022 (2021/22 Marketing Year)**

Processed For	Cape Provinces	Free State	Kwazulu-Natal	Mpumalanga	Limpopo	Gauteng	Northwest	Total
Human Consumption and Gristing	187 871	1 419 589	247 734	571 977	271 890	762 432	731 880	4 193 373
Animal Feed and Industrial	1 045 273	641 112	496 211	620 766	156 412	1 072 572	476 457	4 508 803
	<b>1 233 144</b>	<b>2 060 701</b>	<b>743 945</b>	<b>1 192 743</b>	<b>428 302</b>	<b>1 835 004</b>	<b>1 208 337</b>	<b>8 702 176</b>

Publication Date: 2022/04/28

\* Please note that included are the products destined for exports

**WHOLE WHITE MAIZE PROCESSED PER PROVINCE**

**PROGRESSIVE: May 2017 to April 2018 (Full 2017/18 Marketing Year)**

Processed For	Cape Provinces	Free State	Kwazulu-Natal	Mpumalanga	Limpopo	Gauteng	Northwest	Total
Human Consumption and Gristing	273 153	1 354 218	334 987	473 802	426 011	754 160	898 024	4 514 355
Animal Feed and Industrial	870 796	237 831	75 440	43 510	10 689	387 213	436 170	2 061 649
	<b>1 143 949</b>	<b>1 592 049</b>	<b>410 427</b>	<b>517 312</b>	<b>436 700</b>	<b>1 141 373</b>	<b>1 334 194</b>	<b>6 576 004</b>

**PROGRESSIVE: May 2018 to April 2019 (Full 2018/19 Marketing Year)**

Processed For	Cape Provinces	Free State	Kwazulu-Natal	Mpumalanga	Limpopo	Gauteng	Northwest	Total
Human Consumption and Gristing	232 450	1 474 013	278 228	524 341	395 583	794 170	979 579	4 678 364
Animal Feed and Industrial	782 873	197 307	80 064	6 857	7 797	261 040	341 298	1 677 236
	<b>1 015 323</b>	<b>1 671 320</b>	<b>358 292</b>	<b>531 198</b>	<b>403 380</b>	<b>1 055 210</b>	<b>1 320 877</b>	<b>6 355 600</b>

**PROGRESSIVE: May 2019 to April 2020 (Full 2019/20 Marketing Year)**

Processed For	Cape Provinces	Free State	Kwazulu-Natal	Mpumalanga	Limpopo	Gauteng	Northwest	Total
Human Consumption and Gristing	235 250	1 589 290	292 091	602 276	399 058	806 144	1 132 767	5 056 876
Animal Feed and Industrial	266 750	124 731	1 514	2 498	1 924	27 978	203 681	629 076
	<b>502 000</b>	<b>1 714 021</b>	<b>293 605</b>	<b>604 774</b>	<b>400 982</b>	<b>834 122</b>	<b>1 336 448</b>	<b>5 685 952</b>

**PROGRESSIVE: May 2020 to April 2021 (Full 2020/21 Marketing Year)**

Processed For	Cape Provinces	Free State	Kwazulu-Natal	Mpumalanga	Limpopo	Gauteng	Northwest	Total
Human Consumption and Gristing	242 155	1 734 660	355 306	604 682	399 818	789 544	1 141 456	5 267 621
Animal Feed and Industrial	629 902	179 240	10 806	10 912	3 488	192 885	298 726	1 325 959
	<b>872 057</b>	<b>1 913 900</b>	<b>366 112</b>	<b>615 594</b>	<b>403 306</b>	<b>982 429</b>	<b>1 440 182</b>	<b>6 593 580</b>

**PROGRESSIVE: May 2021 to April 2022 (2021/22 Marketing Year)**

Processed For	Cape Provinces	Free State	Kwazulu-Natal	Mpumalanga	Limpopo	Gauteng	Northwest	Total
Human Consumption and Gristing	173 892	1 240 994	219 026	450 855	271 435	644 360	721 866	3 722 428
Animal Feed and Industrial	618 589	260 862	81 772	50 226	18 592	339 316	404 494	1 773 851
	<b>792 481</b>	<b>1 501 856</b>	<b>300 798</b>	<b>501 081</b>	<b>290 027</b>	<b>983 676</b>	<b>1 126 360</b>	<b>5 496 279</b>

Publication Date: 2022/04/28

\* Please note that included are the products destined for exports

**WHOLE YELLOW MAIZE PROCESSED PER PROVINCE**

**PROGRESSIVE: May 2017 to April 2018 (2017/18 Full Marketing Year)**

Processed For	Cape Provinces	Free State	Kwazulu-Natal	Mpumalanga	Limpopo	Gauteng	Northwest	Total
Human Consumption and Gristing	12 322	260 743	21 894	118 130	126	103 157	69 908	586 280
Animal Feed and Industrial	575 040	474 879	381 436	678 638	117 204	1 003 421	99 652	3 330 270
	<b>587 362</b>	<b>735 622</b>	<b>403 330</b>	<b>796 768</b>	<b>117 330</b>	<b>1 106 578</b>	<b>169 560</b>	<b>3 916 550</b>

**PROGRESSIVE: May 2018 to April 2019 (2018/19 Full Marketing Year)**

Processed For	Cape Provinces	Free State	Kwazulu-Natal	Mpumalanga	Limpopo	Gauteng	Northwest	Total
Human Consumption and Gristing	20 196	282 426	41 480	117 136		97 316	50 639	609 193
Animal Feed and Industrial	678 720	539 225	393 773	773 718	133 877	1 185 575	234 888	3 939 776
	<b>698 916</b>	<b>821 651</b>	<b>435 253</b>	<b>890 854</b>	<b>133 877</b>	<b>1 282 891</b>	<b>285 527</b>	<b>4 548 969</b>

**PROGRESSIVE: May 2019 to April 2020 (2019/20 Full Marketing Year)**

Processed For	Cape Provinces	Free State	Kwazulu-Natal	Mpumalanga	Limpopo	Gauteng	Northwest	Total
Human Consumption and Gristing	22 191	283 315	46 667	146 632		105 780	29 157	633 742
Animal Feed and Industrial	1 266 467	636 039	496 708	789 519	129 314	1 409 811	419 672	5 147 530
	<b>1 288 658</b>	<b>919 354</b>	<b>543 375</b>	<b>936 151</b>	<b>129 314</b>	<b>1 515 591</b>	<b>448 829</b>	<b>5 781 272</b>

**PROGRESSIVE: May 2020 to April 2021 (2020/21 Full Marketing Year)**

Processed For	Cape Provinces	Free State	Kwazulu-Natal	Mpumalanga	Limpopo	Gauteng	Northwest	Total
Human Consumption and Gristing	17 627	278 801	47 222	172 711		130 961	18 563	665 885
Animal Feed and Industrial	757 110	548 895	492 504	838 226	135 659	1 199 587	290 682	4 262 663
	<b>774 737</b>	<b>827 696</b>	<b>539 726</b>	<b>1 010 937</b>	<b>135 659</b>	<b>1 330 548</b>	<b>309 245</b>	<b>4 928 548</b>

**PROGRESSIVE: May 2021 to April 2022 (2021/22 Marketing Year)**

Processed For	Cape Provinces	Free State	Kwazulu-Natal	Mpumalanga	Limpopo	Gauteng	Northwest	Total
Human Consumption and Gristing	13 979	178 595	28 708	121 122	455	118 072	10 014	470 945
Animal Feed and Industrial	426 684	380 250	414 439	570 540	137 820	733 256	71 963	2 734 952
	<b>440 663</b>	<b>558 845</b>	<b>443 147</b>	<b>691 662</b>	<b>138 275</b>	<b>851 328</b>	<b>81 977</b>	<b>3 205 897</b>

Publication Date: 2022/04/28

\* Please note that included are the products destined for exports



# SAGIS Maize Product Information


On 14 November 2014, the Minister of Agriculture, Forestry & Fisheries announced statutory measures for the manufacturing of maize & wheaten products.

To comply with the abovementioned statutory measures, manufacturers of these products have to register with SAGIS and submit information with regards to the manufacture, import and export of maize products.


The Minister approved the proposed continuation and amendment of statutory measures, namely registration and the keeping of information and submitting monthly returns in respect of maize products manufactured, processed, imported and/or exported, to SAGIS, for a further period of four years, to lapse on 14 November 2022.

Please see graphs 29 to 32 below for an overview of the white and yellow maize products as well as white and yellow maize meal manufactured for the period May 2021 to March 2022. The tables on pages 21 to 23 provide a summary of the figures for maize products manufactured, imported and exported during the previous two marketing seasons (May to April) as well as the current marketing season to date (May 2021 to March 2022).


GRAPH 29: WHITE MAIZE PRODUCTS MANUFACTURED FROM MAY 2021 TO MARCH 2022




GRAPH 30: YELLOW MAIZE PRODUCTS MANUFACTURED FROM MAY 2021 TO MARCH 2022



GRAPH 31: WHITE MAIZE MEAL MANUFACTURED FROM MAY 2021 TO MARCH 2022



GRAPH 32: YELLOW MAIZE MEAL MANUFACTURED FROM MAY 2021 TO MARCH 2022



**MAIZE PRODUCTS MANUFACTURED PER MARKETING YEAR**

Date Published: 2022/05/06

	Marketing year: May 2019 - Apr 2020 Manufactured Tons Progressive: 12 Months				Marketing year: May 2020 - Apr 2021 Manufactured Tons Progressive: 12 Months				Marketing year: May 2021 - Apr 2022 Manufactured Tons Progressive: 11 Months (May - Mar)			
	White Maize	Yellow Maize	Total Maize		White Maize	Yellow Maize	Total Maize		White Maize	Yellow Maize	Total Maize	
Maize Chop	1 635 434	188 380	1 823 814		1 764 791	205 068	1 969 859		1 515 690	172 523	1 688 213	
Maize Rice	8 476	*	8 476		7 264	*	7 264		6 589	0	6 589	
Maize Grits	68 251	*	68 251		69 446	*	69 446		72 527	0	72 527	
Samp	117 573	*	117 573		110 572	*	110 572		100 109	0	100 109	
<b>* Total Yellow Maize Rice / Maize Grits / Samp</b>		<b>290 205</b>	<b>290 205</b>			<b>301 905</b>	<b>301 905</b>		<b>0</b>	<b>265 862</b>	<b>265 862</b>	
Sifted Maize Meal	22 100	30 105	52 205		21 881	36 050	57 931		20 444	55 742	76 186	
Special Maize Meal	371 313	4 443	375 756		295 406	3 501	298 907		224 758	1 767	226 525	
Super Maize Meal	2 725 436	28 056	2 753 492		2 862 883	25 084	2 887 967		2 483 393	17 430	2 500 823	
Unsifted Maize Meal	12 540	5	12 545		9 987	10	9 997		6 187	43	6 230	
Other maize products intended for Human consumption	123 704	78 202	201 906		114 903	90 768	205 671		116 214	54 838	171 052	
<b>Total</b>	<b>5 084 827</b>	<b>619 396</b>	<b>5 704 223</b>		<b>5 257 133</b>	<b>662 386</b>	<b>5 919 519</b>		<b>4 545 911</b>	<b>568 205</b>	<b>5 114 116</b>	

\* Included total for yellow rice, grits and samp

**MAIZE PRODUCTS IMPORTED PER MARKETING YEAR**

Date Published: 2022/05/06

	Marketing year: May 2019 - Apr 2020 Progressive: 12 Months			Marketing year: May 2020 - Apr 2021 Progressive: 12 Months			Marketing year: May 2021 - Apr 2022 Progressive: 11 Months (May - Mar)		
	White Maize	Yellow Maize	Total Maize	White Maize	Yellow Maize	Total Maize	White Maize	Yellow Maize	Total Maize
Maize Chop	8 861	0	8 861	6 153	0	6 153	17 510	0	17 510
Maize Rice	0	*	0	0	*	0	0	*	0
Maize Grits	0	*	0	0	*	0	0	*	0
Samp	0	*	0	0	*	0	0	*	0
<b>* Total Yellow Maize Rice / Maize Grits / Samp</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>
Sifted Maize Meal	2 869	0	2 869	4 766	0	4 766	0	0	0
Special Maize Meal	0	0	0	0	0	0	5 820	0	5 820
Super Maize Meal	11	0	11	0	0	0	51	0	51
Unsifted Maize Meal	0	0	0	0	0	0	0	0	0
Other maize products intended for Human consumption	0	0	0	0	0	0	0	0	0
<b>Total</b>	<b>11 741</b>	<b>0</b>	<b>11 741</b>	<b>10 919</b>	<b>0</b>	<b>10 919</b>	<b>23 381</b>	<b>0</b>	<b>23 381</b>

\* Included total for yellow rice, grits and samp

**MAIZE PRODUCTS EXPORTED PER MARKETING YEAR**

Date Published: 2022/05/06

	Marketing year: May 2019 - Apr 2020 Exported Tons Progressive: 12 Months				Marketing year: May 2020 - Apr 2021 Exported Tons Progressive: 12 Months				Marketing year: May 2021 - Apr 2022 Exported Tons Progressive: 11 Months (May - Mar)			
	White Maize	Yellow Maize	Total Maize		White Maize	Yellow Maize	Total Maize		White Maize	Yellow Maize	Total Maize	
Maize Chop	631	0	631		375	0	375		163	14	177	
Maize Rice	72	*	72		8	*	8		3	*	3	
Maize Grits	253	*	253		0	*	0		180	*	180	
Samp	1 322	*	1 322		1 086	*	1 086		1 421	*	1 421	
<b>* Total Yellow Maize Rice / Maize Grits / Samp</b>		<b>16 230</b>	<b>16 230</b>			<b>27 937</b>	<b>27 937</b>			<b>35 783</b>	<b>35 783</b>	
Sifted Maize Meal	0	9 696	9 696		0	11 002	11 002		552	37 027	37 579	
Special Maize Meal	36 695	817	37 512		14 136	0	14 136		7 905	0	7 905	
Super Maize Meal	92 288	8 410	100 698		89 834	7 015	96 849		53 397	2 087	55 484	
Unsifted Maize Meal	580	0	580		0	0	0		265	0	265	
Other maize products intended for Human consumption	44 814	858	45 672		32 355	9 345	41 700		52 227	2 292	54 519	
<b>Total</b>	<b>176 655</b>	<b>36 011</b>	<b>212 666</b>		<b>137 794</b>	<b>55 299</b>	<b>193 093</b>		<b>116 113</b>	<b>77 203</b>	<b>193 316</b>	


\* Included total for yellow rice, grits and samp

# Maize Crop Quality 2020/21 - summary of results

## RSA GRADING

The maize crop, the second largest in history, was of better grading quality than the previous two seasons, comparing well with that of the 2013/14 and 2015/16 seasons. 69% of white maize samples received and graded was graded as maize grade one, last season this figure was only 43%. 78% of yellow maize samples received and graded was graded as grade one, compared to the 65% of the previous season. Please see Graph 33 for the percentages of samples (white and yellow) per season graded as grade 1, since commencement of the annual maize crop quality survey in 1998.

GRAPH 33: PERCENTAGE SAMPLES GRADED AS GRADE 1 OVER SEASONS



The percentage total defective kernels above and below the 6.35 mm sieve, 5.1% for white and 4.2% for yellow maize, was respectively 5.0% and 2.4% lower than the previous season. Defective white maize kernels above the 6.35 mm sieve made the largest contribution to the decrease in the percentage total defective kernels, decreasing from 8.1% last season to 3.0% this season. The percentage defective kernels below the 6.35 mm sieve for white maize increased slightly from 2.0% to 2.1% and that of yellow maize decreased slightly from 2.3% to 2.1%. The average percentage Diplodia infected kernels in white and yellow maize equaled the 0% of the previous season. Fusarium infected kernels of white maize equaled the 0.8% of the previous season, while Fusarium infected yellow maize decreased marginally from 0.7% to 0.6%.

The percentage of white maize samples that were downgraded to class other maize as a result of the percentage foreign matter exceeding 0.75%, decreased from 7% (38 samples) to 5% (29 samples) this season. The percentage for yellow maize increased slightly from 4% (14 samples) to 5% (21 samples) this season. One white and four yellow maize sample were downgraded as a result of other colour maize that exceeded 10% and 5% (maximum permissible deviation for grade 3) respectively. The average percentage combined deviations of white maize was 5.6% compared to the 10.7% of the 2019/20 season and that of yellow maize 4.5% compared to 6.9% previously.

Please refer to Tables 3 to 7 and Graphs 34 to 36 on pages 35 to 47.

## USA GRADING

Of the 1 000 maize samples graded according to USA grading regulations, 62% were graded US1, 21% US2, 7% US3, 3% US4, 2% US5, while sample grade and class mixed corn represented 3% and 2% respectively. The percentage samples graded as US1 varies substantially over seasons, varying from 30% to 41%, 51%, 71% and 58% over the previous five seasons. The percentage samples graded as US2 compared with the 25% and 27% of the previous two seasons respectively albeit lower. The main reason for downgrading the samples was (as in previous seasons) the percentage total damaged kernels exceeding the maximum limit per grade, followed by broken corn and foreign material. Please see Tables 8 and 9 on pages 48 to 53.

## PHYSICAL QUALITY CHARACTERISTICS

Bushel weight/Test weight is applied as a grading factor in the USA grading regulations and is also routinely

done at most intake points locally for stock verification purposes. White maize had an average test weight of 75.9 kg/hl compared to the 76.5 kg/hl of yellow maize. White and yellow maize's average test weight was respectively 0.3 kg/hl and 0.2 kg/hl higher than in the previous season. The test weight in total varied from 68.5 kg/hl to 82.6 kg/hl.

21 samples (2.1%) reported Bushel weight values below the minimum requirement (56.0 lbs or 72.1 kg/hl) for USA grade 1 maize, two of these samples were from the Vaalharts region, 10 were from the North West production regions, five from the Free State and six from Mpumalanga.

The 100 kernel mass ("as is" basis) of white maize was 33.3 g (34.9 g in 2019/20) and averaged higher than yellow maize's 31.2 g (last season 30.4 g). This trend is also observed in previous seasons. The percentage white maize kernels above the 10 mm sieve (21.6%) decreased by 6.2% compared to the previous season. The percentage yellow maize kernels above the 10 mm sieve (7.4%) was 1% lower than last season. The percentage yellow maize kernels above the 10 mm sieve was on average 14.2% lower than white kernels and the percentage yellow kernels below the 8 mm sieve 32.4% higher than that of white maize.

The percentages maize below the 6.35 mm and 4.75 mm sieves provides an indication of the breakage susceptibility. White maize was slightly less susceptible to breakage than during the previous season and the same can also be said for yellow maize. The percentage stress cracks observed varied overall from 1 to 49% and averaged 12%. White and yellow maize both also averaged 12%, previously 16% and 13% respectively. The average stress crack percentages over the last three seasons were the highest of all the seasons since 1999/00 when stress crack analyses were commenced.

Please refer to Tables 12 to 16 on pages 55 to 65 and Graphs 37 to 40 on pages 65 and 66.

The milling index obtained from the SAGL Milling Index 2021 model, varied from an average of 73 (equal to 2019/20) for white maize to an average of 76 (77 previously) for yellow maize. Grit Yield All (GYA) values averaged 63 for white maize and 64 for yellow maize, both equal to the previous season's averages.

## *ROFF MILLING AND WHITENESS INDEX (WI)*

The average % extraction of total meal in white maize obtained with the Roff mill, averaged 77.6% (1.4% higher than the previous season) and varied from 66.6% to 81.4%. Please see Graphs 41 to 46 on page 72 for a comparison of the different fractions percentages as well as the percentage total meal extraction obtained on the Roff mill since 2012/13, the season when the development of the new model for Milling Index was commenced.

The whiteness index averaged 36.3 for unsifted and 27.0 for sifted maize meal. Sieving the sample eliminates differences in the readings as a result of particle size. The whiteness index of the previous season averaged 31.8 and 21.8 for unsifted and sifted maize meal respectively.

The higher the WI value obtained, the whiter the meal sample. The main contributing factors causing differences in WI values are the presence of other colour maize like yellow maize, the presence of defective kernels, the type of cultivar as well as the soil composition. The sample with the lowest sifted whiteness index values this season, namely -5.0, also had the highest percentage other colour maize, namely 12.8%. Please see Tables 17 and 18 on pages 67 to 71.

## *NUTRITIONAL VALUES*

The maize industry requested that crude fibre be added to the scope of analysis performed on the annual maize crop quality survey. With the assistance of Foss, a calibration was developed on the Infratec 1241 Grain Analyser (NIT) during the 2017/18 season. The calibration will be updated annually with the latest season's results.

The average fat content of white maize equaled the 4.0% of the previous two seasons. Yellow maize also averaged 4.0%, 0.1% higher than the previous season. The 10-year average fat content of white maize is 4.1% and that of yellow maize 4.0%. The average protein content of yellow maize was 8.7%, while white maize averaged 8.3%, the lowest since the 2010/11 season. The 10-year average for yellow and white maize respectively is 9.1% and 8.8%.

The average starch contents of both white maize (75.5%) and yellow maize (74.7%) were 2.3% and 2.4% respectively higher than in the previous season. Ten-year averages for white and yellow maize are 73.2% and 72.9% respectively. The average crude fibre content of both white and yellow maize was 2.3%, 0.4% higher than in the previous season.

The fat, starch, protein and crude fibre nutritional components are reported as % (g/100 g) on a dry base.

Please refer to Tables 19 to 22 on pages 73 to 79 and Graphs 47 to 49 on page 80.

## GENETIC MODIFICATION (GM)

The SAGL used the EnviroLogix QuickComb kit for bulk grain, to screen 100 of the crop samples in order to quantitatively determine the presence of genetically modified maize (Cry1Ab, Cry2Ab and/or CP4 EPSPS traits). 90% of the samples tested positive for the Cry1Ab trait, 96% for Cry2Ab and 98% for the CP4 EPSPS trait.

The sensitivity of the measurements for Cry1Ab using the above-mentioned kit is 0.8%, i.e. approximately 6 GM kernels in 800 conventional maize kernels. The limit of detection (LOD) for measurements of the Cry1Ab protein is 0.4%.

The sensitivity of the measurements for Cry2Ab using the above-mentioned kit is 0.9%, i.e. approximately 8 GM kernels in 800 conventional maize kernels. The limit of detection (LOD) for measurements of the Cry1Ab protein is 0.5%.

The sensitivity of the measurements for CP EPSPS using the above-mentioned kit is 0.5%, i.e. 4 GM kernels in 800 conventional maize kernels. The limit of detection (LOD) for measurements of the Roundup Ready protein is 0.25%.

Values higher than 5%, the highest value of the detection range for all three traits, are reported as > 5%.

Important to remember is that the crop quality samples received and analysed by the SAGL are composite samples per class and grade, made up of individual deliveries to grain silos.

Please see Table 23 on page 81 for the results obtained as well as page 107 for a summary of the Events and Trade names/Brands represented by these three traits.

## MYCOTOXINS

None of the 350 samples analysed this season, tested positive for Aflatoxin, Ochratoxin A, HT-2 or T-2 toxin residues.

The average Fumonisin level (Sum of B<sub>1</sub>, B<sub>2</sub> and B<sub>3</sub>) on all 350 samples tested was 100 µg/kg (ppb), compared to the previous season's average of 143 µg/kg. Levels ranged from not detected (ND) to 5 373 µg/kg. Of the 350 samples tested, 76 samples (22%) tested positive for fumonisin levels and the average of these positive results was 459 µg/kg. The previous season, 35% of the samples tested positive, with an average of 413 µg/kg.

The highest Deoxynivalenol (DON) level detected this season was 3 256 µg/kg, compared to the 7 700 µg/kg of last season. The average level of all samples tested this season was 279 µg/kg, 656 µg/kg the previous season. Both the percentage of positive results as well as the average of the positive results decreased this season. 85% of the samples tested positive for DON last season with the average of the positive results 768 µg/kg. This season, 64% of the samples tested positive with an average of 434 µg/kg.

18% of the samples tested positive for 15-acetyl-deoxynivalenol (15-ADON) residues, compared to 34% the previous season. The average of the positive results was 176 µg/kg compared to 238 µg/kg in the previous season.

Zearalenone residues were found in 3% of the samples, 13% during the previous season. Values ranged from ND to 101 µg/kg. The average of the positive samples was 38 µg/kg compared to the 70 µg/kg of the previous season.

Mycotoxin levels lower than the limit of quantitation (< LOQ) as well as limit of detection (< LOD) were seen as having tested negative for calculation purposes. Please see mycotoxin results in Table 24 on pages 89 to 101.

**TABLE 2: SOUTH AFRICAN MAIZE CROP QUALITY 2020/21 (Weighted Averages)**

Class and grade of maize	WM1	WM2	WM3	WCOM	YM1	YM2	YM3	YCOM	Weighted Ave.
<b>RSA Grading</b>									
Defective kernels above 6.35 mm sieve, %	2.2	4.4	6.4	4.3	1.9	2.5	6.0	2.5	2.6
Defective kernels below 6.35 mm sieve, %	1.6	2.9	3.6	3.2	1.8	4.0	3.3	3.2	2.1
Total defective kernels, %	3.8	7.3	10.0	7.5	3.6	6.4	9.3	5.8	4.7
Other colour maize kernels, %	0.2	0.6	0.8	0.6	0.1	0.3	0.2	0.8	0.3
Foreign matter, %	0.1	0.2	0.4	0.9	0.0	0.2	0.4	0.7	0.2
Combined deviation, %	4.1	8.1	11.3	9.0	3.7	6.8	10.0	7.3	5.1
Pinked maize kernels, %	0.2	0.2	0.1	0.0	0.0	0.0	0.0	0.0	0.1
<b>Physical Factors</b>									
Test weight, kg/ha	76.1	76.0	74.8	75.0	76.9	75.2	75.6	75.3	76.2
100 Kernel mass, g	33.2	34.2	32.6	33.1	31.6	29.4	29.7	30.2	32.4
Stress cracks, %	11	14	15	14	11	13	15	14	12
Milling Index	73	74	71	73	77	73	75	76	75
Grit Yield	63	63	63	63	64	63	64	64	63
<b>Kernel Size</b>									
% on top 10 mm	21.0	23.1	23.4	22.6	7.8	5.2	4.6	7.3	15.4
% on top 8 mm	65.6	64.3	64.3	63.5	66.6	61.1	62.9	65.9	65.5
% through 8 mm	13.4	12.6	12.3	13.9	25.6	33.8	32.5	26.8	19.1
<b>Breakage susceptibility</b>									
% Below 6.35 mm sieve	0.7	1.0	1.1	1.0	0.7	0.9	1.3	1.1	0.8
% Below 4.75 mm sieve	0.5	0.7	0.8	0.7	0.5	0.7	0.8	0.8	0.6
<b>Nutritional Values</b>									
Fat, % (db)	4.0	4.0	3.9	4.0	4.0	3.9	3.9	4.0	4.0
Protein, %	8.3	8.3	8.1	8.2	8.7	8.5	8.9	8.8	8.5
Starch, % (db)	75.6	75.5	75.6	75.5	74.7	74.9	74.4	74.5	75.2
Crude fibre, % (db)	2.3	2.3	2.3	2.3	2.3	2.3	2.3	2.4	2.3
<b>Number of samples</b>	<b>388</b>	<b>90</b>	<b>36</b>	<b>46</b>	<b>343</b>	<b>43</b>	<b>8</b>	<b>46</b>	<b>1 000</b>
<b>Mycotoxins</b>									
Total Aflatoxin, µg/kg (ppb) [max. value]	0 [0]	0 [0]	0 [0]	0 [0]	0 [0]	0 [0]	0 [0]	0 [0]	0 [0]
Total Fumonisin, µg/kg (ppb) [max. value]	29 [561]	407 [5 373]	0 [0]	78 [706]	118 [2 648]	76 [735]	114 [687]	8 [100]	109 [5 373]
Deoxynivalenol, µg/kg (ppb) [max. value]	269 [1 606]	680 [3 256]	162 [594]	245 [862]	210 [1 248]	148 [573]	240 [474]	343 [2 169]	279 [3 256]
15-ADON, µg/kg (ppb) [max. value]	29 [403]	104 [571]	9 [105]	22 [127]	20 [309]	18 [244]	17 [101]	41 [237]	71 [571]
Ochratoxin A, µg/kg (ppb) [max. value]	0 [0]	0 [0]	0 [0]	0 [0]	0 [0]	0 [0]	0 [0]	0 [0]	0 [0]
Zearalenone, µg/kg (ppb) [max. value]	1 [27]	4 [101]	0 [<20]	1 [20]	0 [52]	3 [56]	0 [<20]	0 [0]	1 [101]
HT-2 Toxin, µg/kg (ppb) [max. value]	0 [0]	0 [0]	0 [0]	0 [0]	0 [0]	0 [0]	0 [0]	0 [0]	0 [0]
T - 2 Toxin, µg/kg (ppb) [max. value]	0 [0]	0 [0]	0 [0]	0 [0]	0 [0]	0 [0]	0 [0]	0 [0]	0 [0]
<b>Number of samples</b>	<b>124</b>	<b>33</b>	<b>12</b>	<b>17</b>	<b>119</b>	<b>20</b>	<b>6</b>	<b>19</b>	<b>350</b>
<b>GMO</b>									
Cry1Ab, % Samples positive (>LOD of 0.4%)	92	70	100	50	94	86	100	100	90
Cry2Ab, % Samples positive (>LOD of 0.5%)	100	80	100	100	94	100	100	100	96
CP4 EPSPS, % Samples positive (>LOD of 0.25%)	100	80	100	100	100	100	100	100	98
<b>Number of samples</b>	<b>38</b>	<b>10</b>	<b>3</b>	<b>2</b>	<b>36</b>	<b>7</b>	<b>1</b>	<b>3</b>	<b>100</b>


Note: Not detected mycotoxin results are reported as 0, see LOQ in Table 24 page 89.



# RSA Production Regions

The Republic of South Africa is divided into 9 provinces as illustrated in Figure 1.

FIGURE 1: RSA PROVINCES



Provincial map with gratitude to SiQ.

The 9 provinces are divided into 36 grain production regions.

The regions are distributed as follows:

- |                               |                              |
|-------------------------------|------------------------------|
| Region 1: Namakwaland         | Regions 12 to 20: North West |
| Regions 2 to 4: Swartland     | Regions 21 to 28: Free State |
| Regions 5 and 6: Rûens        | Regions 29 to 33: Mpumalanga |
| Regions 7 and 8: Eastern Cape | Region 34: Gauteng           |
| Region 9: Karoo               | Region 35: Limpopo           |
| Region 10: Griqualand West    | Region 36: KwaZulu-Natal     |
| Region 11: Vaalharts          |                              |


Please see the Crop Production Regions map on the next page.

The production regions from which maize samples have been received for the crop quality survey of the 2020/21 production season, are named and described on pages 30 to 33. All the silo/intake stands as well as the type of storage structure, situated in a particular region, are provided.

The mostly rain-fed maize production area is divided into four major maize production regions according to climatological characteristics:

- The Warm Western Region (western parts of the Free State and most of the North West)
- The Temperate Eastern Region (Gauteng and the central parts of the Free State)
- The Cold Eastern Region (Mpumalanga Highveld and eastern Free State)
- The KwaZulu-Natal Region (the western/upland and central/midland parts of KZN)

FIGURE 2: RSA CROP PRODUCTION REGIONS



Regional map with gratitude to Agbiz Grain and SiQ.

# Grain Production Regions

SILO/INTAKE STANDS PER REGION INDICATING TYPE OF STORAGE STRUCTURE

## Region 10: Griqualand West Region

GWK	Douglas (Bags/Bins)	GWK	Trans Oranje (Bags/Bins/Bunkers)
GWK	Luckhoff (Bins)	OVK	Havenga Brug (Bins)
GWK	Marydale (Bins)	OVK	Morgenzon (Bins)
GWK	Modderrivier (Bags/Bins/Bulk)	OVK	Oranjerivier (Bins/Bunkers)
GWK	Prieska (Bins/Dams)	OVK	Prieska (Bins/Bunkers)
GWK	Rietrivier (Bins)	OVK	Rietrivier (Bins)
GWK	Stoffelshoek (Bunkers)		

## Region 11: Vaalharts Region

GWK	Barkly-Wes (Bins/Bulk)	Senwes	Jan Kempdorp (Bins)
GWK	Jan Kempdorp (Bags/Bunkers)	Senwes	Magogong (Bins)
Senwes	Hartswater (Bins)	Senwes	Tadcaster (Bins)

## Region 12: North West Western Region

NWK	Blaauwbank (Bins)	NWK	Mareetsane (Bins)
NWK	Bührmannsdrif (Bins)	Senwes	Kameel (Bins)
NWK	Kameel (Bins)	Senwes	Vryburg (Bins)

## Region 13: North West Central Region (Sannieshof)

NWK	Biesiesvlei (Bins)	NWK	Oppaslaagte (Bins)
NWK	Bossies (Bins)	NWK	Sannieshof (Bins)
NWK	Gerdau (Bins)		

## Region 14: North West Southern Region

NWK	Barberspan (Bins)	NWK	Taaibospan (Bins)
NWK	Delareyville (Bins)	Senwes	Amalia (Bins)
NWK	Excelsior (Bins)	Senwes	Hallatshope (Bins)
NWK	Geysdorp (Bins)	Senwes	Migdol (Bins)
NWK	Migdol (Bins)	Senwes	Schweizer-Reneke (Bins)
NWK	Nooitgedacht (Bins)		

## Region 16: North West Central-Eastern Region

Senwes	Bamboesspruit (Bins)	Senwes	Regina (Bins)
Senwes	Klerksdorp (Bins)	Senwes	Strydpoort (Bins)
Senwes	Leeudoringstad (Bins)	Senwes	Wolmaranstad (Bins)
Senwes	Makwassie (Bins)	Senwes	Zesto (Bunkers)

## Region 17: North West Central-Northern Region (Ottosdal)

NWK	Boschpoort (Bags/Bins/Bulk)	NWK	Vermaas (Bins)
NWK	Kleinharts (Bins)	Senwes	Hartbeesfontein (Bins)
NWK	Ottosdal (Bins)	Senwes	Melliodora (Bins)
NWK	Rostrataville (Bins)	Senwes	Werda (Bins)

# Grain Production Regions

SILO/INTAKE STANDS PER REGION INDICATING TYPE OF STORAGE STRUCTURE

## Region 18: North West Central Region (Ventersdorp)

NWK	Bodenstein (Bins)	Senwes	Makokskraal (Bins)
NWK	Coligny (Bins)	Senwes	Potchefstroom (Bins)
Senwes	Buckingham (Bins)	Senwes	Ventersdorp (Bins)
Senwes	Enselspruit (Bins)		

## Region 19: North West Central Region (Lichtenburg)

Afgri	Lichtenburg (Bunkers)	NWK	Lusthof (Bins)
NWK	Grootpan (Bins)	NWK	Lichtenburg Silo 3 (Bins)
NWK	Halfpad (Bins)	NWK	Lichtenburg Silo 5 (Bins)
NWK	Hibernia (Bins)	NWK	Mafikeng
NWK	Lottie Halte (Bins)		

## Region 20: North West Eastern Region

Afgri	Battery (Bins)	NWK	Derby (Bins)
Afgri	Beestekraal (Bunkers)	NWK	Koster (Bins)
Afgri	Brits (Bins)	NWK	Swartruggens (Bins)
NWK	Boons (Bins)	NWK	Syferbult (Bins)

## Region 21: Free State North-Western Region (Viljoenskroon)

Afgri	Kommandonek (Bunkers)	Senwes	Rooiwal (Bins)
Senwes	Attie (Bins)	Senwes	Vierfontein (Bins)
Senwes	Groenebloem (Bins)	Senwes	Viljoenskroon (Bins)
Senwes	Heuningspruit (Bins)	Senwes	Vredefort (Bins)
Senwes	Koppies (Bins)	Senwes	Weiveld (Bins)

## Region 22: Free State North-Western Region (Bothaville)

Senwes	Allanrigde (Bins)	Senwes	Schoonspruit (Bins)
Senwes	Bothaville (Bins)	Senwes	Schuttendraai (Bins)
Senwes	Mirage (Bins)	Senwes	Misgunst (Bunkers)
Senwes	Odendaalsrus (Bins)		

## Region 23: Free State North-Western Region (Bultfontein)

Senwes	Bultfontein (Bins)	Senwes	Tierfontein (Bins)
Senwes	Kaalplaas (Bins)	Senwes	Wesselsbron (Bins)
Senwes	Losdoorns (Bins)	Senwes	Willemsrus (Bins)
Senwes	Protespan (Bins)		

## Region 24: Free State Central Region

Senwes	Bainsvlei (Bins)	Senwes	Kroonstad (Bins)
Senwes	Bloemfontein (Bins)	Senwes	Petrusburg (Bins)
Senwes	Brandfort (Bins)	Senwes	Theunissen (Bins)
Senwes	De Brug (Bins)	Senwes	Van Tonder (Bins)
Senwes	Geneva (Bins)	Senwes	Welgeleë (Bins)
Senwes	Hennenman (Bins)	Senwes	Winburg (Bins)

# Grain Production Regions

SILO/INTAKE STANDS PER REGION INDICATING TYPE OF STORAGE STRUCTURE

## Region 25: Free State South-Western Region

Afgri	Bethlehem (Bins)	OVK	Modderpoort (Bins)
Afgri	Slabberts (Bins)	OVK	Thaba Nchu (Bunkers)
OVK	Clocolan (Bins)	OVK	Tweespruit (Bins)
OVK	Ficksburg (Bins)	OVK	Westminster (Bins)
OVK	Fouriesburg (Bins)	Senwes	Dewetsdorp (Bins)
OVK	Marseilles (Bins)		

## Region 26: Free State South-Eastern Region

Afgri	Kaallaagte (Bins)	Afgri	Monte Video (Bins)
Afgri	Libertas (Bins)	Afgri	Senekal (Bins)
Afgri	Marquard (Bins)	Senwes	Arlington (Bins)
Afgri	Meets (Bins)	Senwes	Steynsrus (Bins)

## Region 27: Free State Northern Region

Senwes	Gottenburg (Bins)	Senwes	Mooigeleë (Bins)
Senwes	Heilbron (Bins)	Senwes	Wolwehoek (Bins)
Senwes	Hoogte Grainlink (Bins)	VKB	Petrus Steyn (Bins)

## Region 28: Free State Eastern Region

Afgri	Afrikaskop (Bins/Bunkers)	VKB	Jim Fouché (Bins)
Afgri	Eeram (Bins)	VKB	Memel (Bins)
Afgri	Harrismith (Bins)	VKB	Reitz (Bins)
Afgri	Kransfontein (Bins/Bunkers)	VKB	Tweeling (Bins)
VKB	Ascent (Bins)	VKB	Villiers (Bins/Bulk)
VKB	Cornelia (Bins)	VKB	Vrede (Bins)
VKB	Daniëlsrus (Bins)	VKB	Warden (Bins)
VKB	Frankfort (Bins)	VKB	Windfield (Bins)

## Region 29: Mpumalanga Southern Region

Afgri	Balfour (Bins)	Afgri	Leeuspruit (Bins)
Afgri	Greylingstad (Bins)	Afgri	Platrand (Bins)
Afgri	Grootvlei (Bins)	Afgri	Standerton (Bins)
Afgri	Harvard (Bins)	Afgri	Vaaldrift (Bunkers)
Afgri	Holmdene (Bins)	Afgri	Val (Bins)

## Region 30: Mpumalanga Eastern Region

Afgri	Amersfoort (Bins)	Afgri	Maizefield (Bins)
Afgri	Carolina (Bins)	Afgri	Morgenzon (Bins)
Afgri	Davel (Bins)	Afgri	Overvaal (Bins)
Afgri	Eerstelingsfontein (Bunkers)	Afgri	Sandspruit (Bunkers)
Afgri	Ermelo (Bins)	BKB	Waterval (Bunkers)
Afgri	Estancia (Bins)	TWK	Mkondo (Bins)
Afgri	Hendriksvallei (Bunkers)	TWK	Panbult (Bins)
Afgri	Lothair (Bins)	TWK	Rietspruit (Bunkers)

# Grain Production Regions

SILO/INTAKE STANDS PER REGION INDICATING TYPE OF STORAGE STRUCTURE

## Region 31: Mpumalanga Central Region

Afgri	Bakenlaagte (Bunkers)	Afgri	Leslie (Bins)
Afgri	Bethal (Bins)	Afgri	Palmietfontein (Bunkers)
Afgri	Brakfontein (Bunkers)	Afgri	Trichardt (Bins)
Afgri	Devon (Bins)	Afgri	Vaalkrantz (Bunkers)
Afgri	Kinross (Bins/Bunkers)		

## Region 32: Mpumalanga Western Region

Afgri	Argent (Bins/Bunkers)	Afgri	Hawerklip (Bins)
Afgri	Delmas (Bunkers)	Afgri	Kendal (Bins)
Afgri	Dryden (Bins)	Afgri	Ogies (Bins)
Afgri	Eloff (Bins)	Afgri	Vlakfontein (Bunkers)
Afgri	Endicott (Bins)		

## Region 33: Mpumalanga Northern Region

Afgri	Arnot (Bins)	Afgri	Middelburg (Bins)
Afgri	Driefontein (Bins)	Afgri	Pan (Bins)
Afgri	Lydenburg (Bins)	Afgri	Stoffberg (Bins)
Afgri	Marble Hall (Bins)	Afgri	Wonderfontein (Bins)

## Region 34: Gauteng Region

Afgri	Bloekomspruit (Bins)	Afgri	Nigel (Bins)
Afgri	Bronkhorstspuit (Bins)	Afgri	Pretoria Wes (Bins)
Afgri	Glenroy (Bins)	Afgri	Vogelvallei (Bunkers)
Afgri	Goeie Hoek (Bins)	Senwes	Middelvlei (Bins)
Afgri	Kaalfontein (Bins)	Senwes	Oberholzer (Bins)
Afgri	Kliprivier (Bunkers)	Senwes	Raathsvlei (Bins)
Afgri	Meyerton (Bunkers)		

## Region 35: Limpopo Region

Afgri	Northam (Bins)	VKB	Nylstroom (Modimolle) (Bins)
VKB	Alma (Bins)	VKB	Potgietersrus (Mokopane) (Bins)
VKB	Lehau (Bins)	VKB	Roedtan (Bins)
VKB	Naboomspruit (Mookgophong) (Bins)	VKB	Settlers (Bins)
VKB	Nutfield (Bins)	VKB	Warmbad (Bela-Bela) (Bins)

## Region 36: KwaZulu-Natal Region

Afgri	Bergville (Bins/Bunkers)	Afgri	Mizpah (Bins)
Afgri	Chelmsford Dam (Bunkers)	Afgri	Paulpietersburg (Bins)
Afgri	Bloedrivier (Bins)	Afgri	Pietermaritzburg (Bins)
Afgri	Dannhauser (Bins)	Afgri	Vryheid (Bins)
Afgri	Dundee (Bins)	Afgri	Winterton (Bins/Bunkers)

# Main maize producing provinces – comparison of results

The quality of the maize produced in the three main maize production provinces, namely the Free State (regions 21 to 28), Mpumalanga (regions 29 to 33) and North West (regions 12 to 20) are compared below, the values provided are all weighted averages.

Average test weights expressed in kilogram per hectoliter for white maize, ranged between 74.8 in North West, 76.3 in the Free State and 76.7 in Mpumalanga. Yellow maize varied from 75.4 kg/hl in the Free State to 76.3 kg/hl in North West and 76.8 kg/hl in Mpumalanga. The white maize 100 kernel mass values ranged from 32.2 g in North West to 34.8 g in Mpumalanga, the Free State averaged 33.5 g. Yellow maize kernels had the highest average 100 kernel mass in Mpumalanga with 32.5 g, followed by 29.0 g in North West and 28.8 g in the Free State.

Kernel sizes are indicated by the percentage of sample above a 10 mm sieve as well as the percentages above and below a 8 mm sieve. The largest white kernel size with regards to the percentage of kernels above the 10 mm sieve, was found in the Free State (24.6%), followed by North West (23.0%). Mpumalanga had the smallest white kernel sizes (19.9%) on average. Mpumalanga however had the largest yellow maize kernels, averaging 9.3% kernels above the 10 mm sieve, followed by North West with 6.6% and the Free State with 5.8%.

Mpumalanga showed the least susceptibility to breakage (lowest percentage below the sieve), with 0.6% for both white and yellow maize passing through the 6.35 mm sieve. North West averaged 0.9% and 1.0% for white and yellow maize respectively and the Free State averaged 0.8% and 0.9% for white and yellow maize respectively. The percentage stress cracks on white maize ranged from 11% in Mpumalanga to 12% in the Free State and 13% in North West. Stress cracks on yellow maize varied between 11% in Mpumalanga, 12% in North West and 14% in the Free State. These values are slightly lower than those of the previous two seasons when the highest percentages of the last 21 seasons for which stress crack results are available, were reported.

The percentage total defective kernels, is the sum of the defective kernels that remained above the 6.35 mm sieve and the defective kernels which passed through the 6.35 mm sieve. Defective kernels include amongst others, mouldy, discoloured, insect damaged and small kernels that can pass through the 6.35 mm round hole sieve. The production regions in all three of these provinces averaged 5.1% for white maize. The highest percentage total defective kernels on yellow maize (4.8%) was found in the Free State, followed by North West with 4.0% and Mpumalanga with 3.7%. Please see page 103 for the definition of Defective maize kernels as quoted from the Grading Regulations.

The average milling index on white and yellow maize (yellow maize in brackets) was as follows: Mpumalanga averaged 77 (77), the Free State 70 (74) and North West 69 (82). The highest percentage total extraction as determined on the Roff laboratory mill, was found on white maize from the Free State (78.0%), followed by Mpumalanga (77.7%) and North West with 77.1%.

The meal obtained from the white maize in North West gave an average whiteness index of 39.2 (unsifted) and 30.0 (sifted). The Free State gave an average of 37.3 (unsifted) and 27.2 (sifted) and Mpumalanga 32.6 (unsifted) and 23.3 (sifted).

The nutritional component analyses namely crude fat, crude protein, crude fibre and total starch compared well between the three provinces. All three averaged 4.0% fat on white maize. The average fat content on yellow maize ranged from 3.9% in the Free State to 4.0% in North West and 4.1% in Mpumalanga. The lowest average protein content on white maize was found in North West and the Free State, both with 8.0%, Mpumalanga averaged 8.6%. The protein content on yellow maize varied from 8.5% in the Free State to 8.6% in North West and 8.9% in Mpumalanga. Crude fibre on white and yellow maize and in all three of these provinces, averaged 2.3%. North West had the highest average starch content on white maize, namely 75.8%, followed closely by the Free State with 75.7% and Mpumalanga with 75.2%. The yellow maize starch content ranged from a low of 74.3% in Mpumalanga to a high of 75.2% in North West. The Free State averaged 74.8%. These values are all reported on a dry basis.

TABLE 3: RSA GRADING OF WHITE MAIZE ACCORDING TO GRADE (2020/21)																															
Number of samples	Region	% Defective Kernels						% Total defective	% Foreign matter	% Other Colour		% Combined Deviations	% Pinked Kernels		% Diplodia Kernels	% Fusarium Kernels		% Cobrot Kernels													
		Above 6.35 mm sieve		Below 6.35 mm sieve		ave.	min.			max.	ave.		min.	max.		ave.	min.	max.	ave.	min.	max.										
		ave.	min.	max.	ave.																	min.	max.	ave.	min.	max.	ave.	min.	max.		
<b>GRADE: WM1</b>																															
8	Region 12	2.0	1.0	3.1	2.5	1.2	4.3	4.5	2.6	6.9	0.1	0.0	0.3	0.1	0.0	0.4	4.7	2.6	6.9	0.0	0.0	0.0	0.0	0.0	0.4	0.0	1.3	0.4	0.0	1.3	
23	Region 13	2.1	0.9	4.1	2.0	0.3	5.3	4.2	1.6	6.6	0.1	0.0	0.3	0.1	0.0	1.0	4.4	1.6	6.8	0.0	0.0	0.0	0.0	0.0	0.5	0.0	1.4	0.5	0.0	1.4	
25	Region 14	2.5	0.2	5.3	1.5	0.2	2.9	4.0	0.5	6.8	0.1	0.0	0.2	0.0	0.0	0.2	4.1	0.5	7.0	0.1	0.0	1.5	0.0	0.0	0.6	0.0	1.7	0.6	0.0	1.7	
6	Region 16	2.0	0.9	3.6	1.6	1.1	3.1	3.7	2.4	5.2	0.1	0.0	0.2	0.0	0.0	0.0	3.8	2.6	5.4	0.0	0.0	0.0	0.1	0.0	0.5	0.0	1.2	0.6	0.0	1.7	
15	Region 17	1.8	0.7	3.4	1.5	0.4	2.9	3.3	1.5	4.5	0.0	0.0	0.1	0.1	0.0	0.5	3.4	1.5	4.5	0.1	0.0	0.4	0.0	0.0	0.4	0.5	0.0	1.8	0.5	0.0	1.8
24	Region 18	1.6	0.4	4.4	2.0	0.7	5.1	3.6	1.7	6.0	0.1	0.0	0.3	0.4	0.0	2.4	4.0	1.7	6.3	0.0	0.0	0.2	0.0	0.0	0.0	0.4	0.0	2.6	0.4	0.0	2.6
17	Region 19	1.8	0.5	3.9	2.2	0.6	4.2	4.0	2.0	6.9	0.1	0.0	0.3	0.1	0.0	0.9	4.2	2.3	6.9	0.3	0.0	5.4	0.0	0.0	0.0	0.3	0.0	0.7	0.3	0.0	0.7
15	Region 20	1.7	0.6	4.7	1.4	0.4	2.7	3.1	1.3	6.9	0.0	0.0	0.2	0.2	0.0	0.6	3.3	1.6	7.7	0.0	0.0	0.2	0.0	0.0	0.0	0.3	0.0	0.9	0.3	0.0	0.9
27	Region 21	1.4	0.0	4.6	1.9	0.9	6.7	3.3	1.5	6.7	0.0	0.0	0.2	0.1	0.0	0.6	3.4	1.6	7.4	0.0	0.0	0.2	0.0	0.0	0.0	0.3	0.0	1.7	0.3	0.0	1.7
9	Region 22	2.8	1.3	4.2	2.1	0.7	2.9	4.9	2.7	6.7	0.1	0.0	0.2	0.0	0.0	0.0	4.9	2.8	6.7	0.0	0.0	0.0	0.1	0.0	0.7	0.9	0.0	2.1	1.0	0.0	2.1
31	Region 23	1.7	0.7	3.9	1.8	0.7	4.7	3.5	1.7	6.9	0.1	0.0	0.2	0.0	0.0	0.2	3.6	1.7	6.9	0.0	0.0	0.3	0.0	0.0	0.1	0.4	0.0	2.6	0.4	0.0	2.6
10	Region 24	2.5	0.5	4.0	1.7	0.7	2.5	4.2	2.4	6.0	0.1	0.0	0.3	0.1	0.0	0.7	4.3	2.7	6.0	0.1	0.0	0.5	0.0	0.0	0.2	0.8	0.0	1.7	0.8	0.0	1.7
2	Region 25	1.7	1.3	2.0	1.3	1.2	1.4	3.0	2.8	3.2	0.2	0.0	0.3	0.0	0.0	0.0	3.1	3.1	3.2	0.0	0.0	0.0	0.0	0.0	0.0	0.3	0.3	0.4	0.3	0.3	0.4
5	Region 26	2.4	1.0	5.0	1.1	0.2	2.4	3.5	1.2	5.2	0.1	0.0	0.2	0.4	0.0	1.3	4.0	1.8	6.6	0.1	0.0	0.4	0.0	0.0	0.0	0.5	0.0	0.9	0.5	0.0	0.9
3	Region 27	1.3	0.3	2.8	2.9	2.2	4.1	4.1	3.0	5.0	0.0	0.0	0.0	0.3	0.0	0.7	4.4	3.2	5.7	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.0	1.2	0.5	0.0	1.2
12	Region 28	1.8	0.3	4.9	1.4	0.2	2.4	3.2	1.4	5.4	0.1	0.0	0.2	0.4	0.0	1.2	3.6	1.4	6.8	0.1	0.0	0.6	0.0	0.0	0.0	0.4	0.0	1.7	0.4	0.0	1.7
26	Region 29	2.5	0.7	4.7	1.3	0.3	3.1	3.8	1.2	6.0	0.1	0.0	0.3	0.3	0.0	1.5	4.2	2.0	6.2	0.0	0.0	0.5	0.0	0.0	0.0	0.5	0.0	1.1	0.5	0.0	1.1
23	Region 30	3.2	2.0	5.0	1.2	0.4	2.4	4.5	2.9	6.4	0.0	0.0	0.1	0.8	0.0	1.8	5.2	3.1	7.1	0.1	0.0	1.2	0.0	0.0	0.4	0.8	0.0	2.3	0.8	0.0	2.3
10	Region 31	2.5	0.2	4.0	1.5	0.8	2.3	4.0	1.3	5.9	0.0	0.0	0.2	0.6	0.0	2.2	4.6	1.5	6.8	0.2	0.0	2.1	0.0	0.0	0.0	0.7	0.1	1.2	0.7	0.1	1.2
13	Region 32	2.6	0.4	4.4	1.7	0.7	3.4	4.4	2.8	6.5	0.0	0.0	0.2	0.5	0.0	1.9	4.9	2.8	7.4	0.7	0.0	3.6	0.0	0.0	0.0	0.7	0.1	1.7	0.7	0.1	1.7
40	Region 33	2.7	0.6	5.3	1.0	0.2	2.7	3.8	0.8	6.6	0.0	0.0	0.3	0.5	0.0	2.3	4.3	1.4	8.0	0.1	0.0	4.1	0.0	0.0	0.2	0.8	0.0	2.6	0.8	0.0	2.6
21	Region 34	2.4	0.6	6.0	1.7	0.2	2.8	4.1	2.1	7.0	0.0	0.0	0.3	0.1	0.0	0.4	4.2	2.1	7.1	1.4	0.0	6.6	0.0	0.0	0.0	0.5	0.0	1.8	0.5	0.0	1.8
6	Region 35	1.6	0.1	3.8	0.4	0.0	1.0	2.0	0.6	3.9	0.0	0.0	0.2	0.2	0.0	0.7	2.2	0.6	4.2	0.0	0.0	0.2	0.0	0.0	0.0	0.1	0.0	0.3	0.1	0.0	0.3
17	Region 36	3.3	1.1	5.4	1.5	0.7	4.5	4.7	2.0	6.8	0.1	0.0	0.3	0.2	0.0	1.1	5.0	2.0	6.9	0.1	0.0	0.5	0.0	0.0	0.0	0.7	0.0	1.9	0.7	0.0	1.9
388	Ave. WM1	2.2	0.0	6.0	1.6	0.0	6.7	3.8	0.5	7.0	0.1	0.0	0.3	0.2	0.0	1.1	4.1	0.5	8.0	0.2	0.0	6.6	0.0	0.0	0.7	0.5	0.0	2.6	0.5	0.0	2.6
	Min. WM1	0.0	0.0	6.0	0.0	0.0	6.7	0.5	0.5	7.0	0.0	0.0	0.3	0.0	0.0	2.4	0.5	8.0	0.0	0.0	6.6	0.0	0.0	0.7	0.5	0.0	2.6	0.0	0.0	2.6	
	Max. WM1	6.0	6.0	6.0	6.7	6.7	6.7	7.0	7.0	7.0	0.3	0.3	0.3	2.4	2.4	2.4	8.0	8.0	8.0	6.6	6.6	6.6	0.7	0.7	0.7	2.6	2.6	2.6	2.6	2.6	2.6





TABLE 3: RSA GRADING OF WHITE MAIZE ACCORDING TO GRADE (2020/21) (continue)																															
Number of samples	Region	% Defective Kernels						% Total defective	% Foreign matter	% Other Colour		% Combined Deviations	% Pinked Kernels		% Diplodia Kernels		% Fusarium Kernels		% Cobrot Kernels												
		Above 6.35 mm sieve		Below 6.35 mm sieve		ave.	min.			max.	ave.		min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.								
		ave.	min.	max.	ave.																			min.	max.	ave.	min.	max.	ave.	min.	max.
<b>GRADE: WM3</b>																															
1	Region 13	2.7	2.7	2.7	3.7	3.7	6.3	6.3	6.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.6	0.6	0.6											
3	Region 14	3.3	2.7	4.1	3.0	1.8	4.9	6.3	4.8	7.6	0.6	0.6	0.7	0.1	0.0	0.2	7.0	5.3	8.1	0.2	0.0	0.5	0.3	0.8							
1	Region 16	11.9	11.9	11.9	1.4	1.4	1.4	13.3	13.3	13.3	0.6	0.6	0.6	0.2	0.2	0.2	14.1	14.1	14.1	0.2	0.2	0.2	1.5	1.5	1.5	8.8	8.8	8.8	10.3	10.3	10.3
8	Region 17	11.4	1.4	20.6	1.3	0.4	3.2	12.7	3.4	21.3	0.3	0.0	0.7	0.0	0.0	0.1	13.1	4.0	21.3	0.1	0.0	0.3	0.5	0.0	2.8	2.3	0.0	8.0	2.9	0.0	10.5
1	Region 18	12.3	12.3	12.3	2.3	2.3	2.3	14.6	14.6	14.6	0.3	0.3	0.3	0.2	0.2	0.2	15.2	15.2	15.2	0.2	0.2	0.2	2.8	2.8	2.8	5.8	5.8	5.8	8.6	8.6	8.6
5	Region 19	4.9	1.5	12.6	6.7	2.7	10.5	11.6	4.2	16.7	0.5	0.0	0.6	0.3	0.0	0.9	12.4	4.8	17.0	0.3	0.0	1.4	0.0	0.0	0.0	1.1	0.2	3.7	1.1	0.2	3.7
1	Region 20	3.6	3.6	3.6	1.6	1.6	1.6	5.2	5.2	5.2	0.6	0.6	0.6	0.5	0.5	0.5	6.3	6.3	6.3	0.1	0.1	0.1	0.0	0.0	0.0	0.8	0.8	0.8	0.8	0.8	0.8
3	Region 21	1.4	0.8	1.8	6.3	2.1	11.4	7.7	3.9	13.1	0.5	0.1	0.7	0.0	0.0	0.0	8.2	4.5	13.2	0.0	0.0	0.0	0.0	0.0	0.0	0.6	0.5	0.6	0.6	0.5	0.6
5	Region 23	9.5	0.2	19.3	2.8	0.4	9.5	12.3	1.4	19.8	0.4	0.0	0.7	0.0	0.0	0.2	12.8	2.1	20.1	0.0	0.0	0.0	0.3	0.0	1.4	4.9	0.0	15.5	5.2	0.0	16.9
1	Region 26	7.5	7.5	7.5	21.1	21.1	21.1	28.7	28.7	28.7	0.1	0.1	0.1	1.3	1.3	1.3	30.0	30.0	30.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.4	0.4	0.4	0.4
2	Region 30	2.2	1.2	3.2	2.0	1.2	2.8	4.2	2.5	6.0	0.3	0.0	0.6	3.4	0.0	6.7	7.9	3.0	12.7	0.0	0.0	0.0	0.0	0.0	0.0	1.1	0.5	1.8	1.1	0.5	1.8
2	Region 31	2.0	1.2	2.8	2.0	0.9	3.1	4.0	3.8	4.3	0.5	0.5	0.6	4.8	1.8	7.9	9.4	6.6	12.2	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.2	0.8	0.5	0.2	0.8
1	Region 32	4.8	4.8	4.8	2.5	2.5	2.5	7.3	7.3	7.3	0.6	0.6	0.6	0.2	0.2	0.2	8.1	8.1	8.1	0.0	0.0	0.0	0.0	0.0	0.0	2.3	2.3	2.3	2.3	2.3	2.3
1	Region 33	1.4	1.4	1.4	2.6	2.6	2.6	4.1	4.1	4.1	0.4	0.4	0.4	7.4	7.4	7.4	11.9	11.9	11.9	0.4	0.4	0.4	0.0	0.0	0.0	0.3	0.3	0.3	0.3	0.3	0.3
1	Region 34	0.6	0.6	0.6	1.2	1.2	1.2	1.8	1.8	1.8	0.7	0.7	0.7	1.2	1.2	1.2	3.6	3.6	3.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
36	Ave. WM3	6.4			3.6			10.0			0.4			0.8			11.3			0.1			0.3			2.1			2.3		
	Min. WM3	0.2			0.4			1.4			0.0			0.0			2.1			0.0			0.0			0.0			0.0		
	Max. WM3	20.6			21.1			28.7			0.7			7.9			30.0			1.4			2.8			15.5			16.9		



**TABLE 3: RSA GRADING OF WHITE MAIZE ACCORDING TO GRADE (2020/21)**  
(continue)

\*The following white maize samples were downgraded to Class Other Maize due to the presence of poisonous seeds exceeding the maximum allowance

Region	Number of Poisonous seeds (Crotalaria spp., Datura spp., Ricinis communis) Max. allowance 1 seed/1000 g	Number of Poisonous seeds (Argemone mexicana L., Convolvulus spp., Ipomoea purpurea Roth., Lolium temulentum, Xanthium spp.) Max. allowance 7 seeds/1000 g
11	18 <i>Datura</i> spp.	0
12	20 <i>Datura</i> spp.	0
12	6 <i>Datura</i> spp.	0
12	6 <i>Datura</i> spp.	0
14	6 <i>Datura</i> spp.	0
17	6 <i>Datura</i> spp.	0
21	0	24 <i>Xanthium Strumarium</i>
21	6 <i>Datura</i> spp.	0
21	6 <i>Datura</i> spp.	0
28	6 <i>Datura</i> spp.	0
32	6 <i>Datura</i> spp.	0
32	12 <i>Datura</i> spp.	0
32	0	10 <i>Xanthium Strumarium</i>
32	6 <i>Datura</i> spp.	0
34	6 <i>Datura</i> spp.	0
36	0	24 <i>Xanthium Strumarium</i>
36	0	12 <i>Xanthium Strumarium</i>
36	0	12 <i>Xanthium Strumarium</i>

**TABLE 4: RSA GRADING OF WHITE MAIZE (2020/21)**

Number of samples	Region	% Defective Kernels		% Total defective		% Foreign matter		% Other Colour		% Combined Deviations		% Pinked Kernels		% Diplodia Kernels		% Fusarium Kernels		% Cobrot Kernels													
		Above 6.35 mm sieve		Below 6.35 mm sieve		ave.   min.   max.		ave.   min.   max.		ave.   min.   max.		ave.   min.   max.		ave.   min.   max.		ave.   min.   max.		ave.   min.   max.													
		ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.									
<b>GRADE: WHITE</b>																															
2	Region 11	3.2	0.9	5.5	2.6	2.4	2.8	5.8	3.3	8.3	0.2	0.1	0.3	0.2	0.0	0.3	0.3	0.0	0.5	0.0	0.0	0.0	0.0	0.0	0.2	0.2	0.3	0.2	0.2	0.3	
14	Region 12	2.8	1.0	5.0	3.4	1.2	7.6	6.2	2.6	10.8	0.2	0.0	0.4	0.1	0.0	0.4	0.6	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.6	0.0	1.9	0.6	0.0	2.3	
29	Region 13	2.4	0.9	7.8	2.3	0.3	5.3	4.7	1.6	11.5	0.2	0.0	1.2	0.1	0.0	1.0	4.9	1.6	11.7	0.0	0.0	0.3	0.0	0.0	0.6	0.5	0.0	1.8	0.6	0.0	1.8
40	Region 14	3.0	0.2	8.0	1.9	0.2	4.9	4.9	0.5	10.3	0.2	0.0	1.4	0.0	0.0	0.2	5.2	0.5	10.3	0.1	0.0	1.5	0.0	0.0	0.0	0.8	0.0	2.4	0.8	0.0	2.4
8	Region 16	3.1	0.9	11.9	1.6	1.1	3.1	4.7	2.3	13.3	0.1	0.0	0.6	0.5	0.0	4.0	5.4	2.6	14.1	0.0	0.0	0.2	0.3	0.0	1.5	1.5	0.0	8.8	1.8	0.0	10.3
29	Region 17	4.8	0.6	20.6	1.6	0.4	3.5	6.4	1.5	21.3	0.2	0.0	1.3	0.1	0.0	0.6	6.7	1.5	21.3	0.1	0.0	0.4	0.2	0.0	2.8	1.1	0.0	8.0	1.3	0.0	10.5
29	Region 18	2.0	0.4	12.3	2.4	0.7	7.4	4.4	1.7	14.6	0.1	0.0	1.1	0.3	0.0	2.4	4.9	1.7	15.2	0.0	0.0	0.2	0.1	0.0	2.8	0.5	0.0	5.8	0.6	0.0	8.6
29	Region 19	2.4	0.4	12.6	3.5	0.6	10.5	5.9	2.0	16.7	0.3	0.0	3.1	0.1	0.0	0.9	6.4	2.3	17.0	0.3	0.0	5.4	0.0	0.0	0.0	0.5	0.0	3.7	0.5	0.0	3.7
17	Region 20	2.3	0.6	9.5	1.4	0.4	2.7	3.7	1.3	10.7	0.1	0.0	0.6	0.3	0.0	1.5	4.0	1.6	12.4	0.0	0.0	0.2	0.0	0.0	0.0	0.5	0.0	3.9	0.5	0.0	3.9
36	Region 21	1.6	0.0	6.8	3.1	0.9	13.2	4.7	1.5	13.8	0.1	0.0	0.8	0.0	0.0	0.6	4.8	1.6	13.8	0.0	0.0	0.2	0.0	0.0	0.0	0.4	0.0	2.4	0.4	0.0	2.4
13	Region 22	3.4	1.3	6.8	4.6	0.7	21.0	8.0	2.7	24.1	0.5	0.0	3.9	0.2	0.0	1.4	8.7	2.8	28.7	0.0	0.0	0.0	0.1	0.0	0.7	0.8	0.0	2.1	0.9	0.0	2.1
47	Region 23	2.9	0.2	19.3	2.0	0.4	9.5	4.9	1.4	19.8	0.2	0.0	2.1	0.0	0.0	0.2	5.1	1.7	20.1	0.0	0.0	0.3	0.0	0.0	1.4	1.1	0.0	15.5	1.1	0.0	16.9
13	Region 24	2.6	0.5	5.9	2.0	0.7	6.3	4.6	2.4	7.4	0.2	0.0	0.9	0.1	0.0	0.7	4.9	2.7	7.7	0.1	0.0	0.5	0.0	0.0	0.2	0.8	0.0	1.8	0.8	0.0	1.8
3	Region 25	1.9	1.3	2.5	2.5	1.2	4.8	4.4	2.8	7.2	0.9	0.0	2.3	0.3	0.0	1.0	5.6	3.1	10.6	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.0	0.4	0.2	0.0	0.4
8	Region 26	2.6	0.4	7.5	4.8	0.2	21.1	7.4	1.2	28.7	0.1	0.0	0.2	1.2	0.0	5.7	8.7	1.8	30.0	0.0	0.0	0.4	0.0	0.0	0.0	0.4	0.0	0.9	0.4	0.0	0.9
5	Region 27	1.0	0.2	2.8	4.5	2.2	7.8	5.5	3.0	8.0	0.0	0.0	0.1	0.3	0.0	0.7	5.8	3.2	8.4	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	1.2	0.4	0.0	1.2
13	Region 28	1.8	0.3	4.9	1.3	0.2	2.4	3.1	1.4	5.4	0.1	0.0	0.4	0.4	0.0	1.2	3.5	1.4	6.8	0.1	0.0	0.6	0.0	0.0	0.0	0.4	0.0	1.7	0.4	0.0	1.7
27	Region 29	2.9	0.7	12.3	1.2	0.1	3.1	4.1	1.2	12.4	0.1	0.0	0.3	0.3	0.0	1.5	4.5	2.0	12.4	0.0	0.0	0.5	0.0	0.0	0.0	0.6	0.0	3.8	0.6	0.0	3.8
34	Region 30	3.6	1.2	7.0	1.5	0.4	5.3	5.1	2.5	10.6	0.1	0.0	1.2	1.3	0.0	6.7	6.5	3.0	12.7	0.1	0.0	1.2	0.0	0.0	0.4	1.1	0.0	2.8	1.1	0.0	2.8
20	Region 31	6.1	0.2	55.1	1.7	0.8	3.1	7.8	1.3	56.9	0.2	0.0	1.2	1.3	0.0	7.9	9.2	1.5	58.7	0.2	0.0	2.1	0.0	0.0	0.0	0.7	0.0	1.4	0.7	0.0	1.4
26	Region 32	3.9	0.4	10.2	1.7	0.7	3.4	5.6	1.7	11.0	0.1	0.0	0.7	0.6	0.0	1.9	6.3	1.7	12.6	0.5	0.0	3.6	0.0	0.0	0.0	1.2	0.1	5.1	1.2	0.1	5.1
54	Region 33	3.2	0.6	9.1	1.3	0.2	4.2	4.5	0.8	10.7	0.1	0.0	2.0	0.9	0.0	12.8	5.5	1.4	16.4	0.1	0.0	4.1	0.1	0.0	1.6	1.0	0.0	4.5	1.1	0.0	6.1
30	Region 34	2.4	0.6	6.0	1.8	0.2	4.1	4.2	1.8	8.1	0.2	0.0	1.0	0.2	0.0	1.2	4.6	2.1	8.1	1.3	0.0	6.6	0.0	0.0	0.0	0.5	0.0	1.8	0.5	0.0	1.8
7	Region 35	2.5	0.1	7.9	0.7	0.0	2.5	3.2	0.6	10.4	0.0	0.0	0.2	0.1	0.0	0.7	3.4	0.6	10.6	0.0	0.0	0.2	0.0	0.0	0.0	0.2	0.0	1.4	0.2	0.0	1.4
27	Region 36	4.0	1.1	6.7	1.7	0.5	4.5	5.6	2.0	8.9	0.2	0.0	2.2	0.2	0.0	1.8	6.1	2.0	9.3	0.1	0.0	0.5	0.0	0.0	0.0	0.9	0.0	2.6	0.9	0.0	2.6
<b>560</b>	<b>Ave. White</b>	<b>3.0</b>	<b>0.0</b>	<b>55.1</b>	<b>2.1</b>	<b>0.0</b>	<b>21.1</b>	<b>5.1</b>	<b>0.5</b>	<b>56.9</b>	<b>0.2</b>	<b>0.0</b>	<b>3.9</b>	<b>0.4</b>	<b>0.0</b>	<b>12.8</b>	<b>5.6</b>	<b>0.5</b>	<b>58.7</b>	<b>0.2</b>	<b>0.0</b>	<b>6.6</b>	<b>0.0</b>	<b>0.0</b>	<b>2.8</b>	<b>0.8</b>	<b>0.0</b>	<b>15.5</b>	<b>0.8</b>	<b>0.0</b>	<b>16.9</b>
	<b>Min. White</b>	<b>0.0</b>	<b>0.0</b>	<b>55.1</b>	<b>0.0</b>	<b>0.5</b>	<b>21.1</b>	<b>5.1</b>	<b>0.5</b>	<b>56.9</b>	<b>0.2</b>	<b>0.0</b>	<b>3.9</b>	<b>0.4</b>	<b>0.0</b>	<b>12.8</b>	<b>5.6</b>	<b>0.5</b>	<b>58.7</b>	<b>0.2</b>	<b>0.0</b>	<b>6.6</b>	<b>0.0</b>	<b>0.0</b>	<b>2.8</b>	<b>0.8</b>	<b>0.0</b>	<b>15.5</b>	<b>0.8</b>	<b>0.0</b>	<b>16.9</b>
	<b>Max. White</b>	<b>3.0</b>	<b>0.0</b>	<b>55.1</b>	<b>2.1</b>	<b>0.5</b>	<b>21.1</b>	<b>5.1</b>	<b>0.5</b>	<b>56.9</b>	<b>0.2</b>	<b>0.0</b>	<b>3.9</b>	<b>0.4</b>	<b>0.0</b>	<b>12.8</b>	<b>5.6</b>	<b>0.5</b>	<b>58.7</b>	<b>0.2</b>	<b>0.0</b>	<b>6.6</b>	<b>0.0</b>	<b>0.0</b>	<b>2.8</b>	<b>0.8</b>	<b>0.0</b>	<b>15.5</b>	<b>0.8</b>	<b>0.0</b>	<b>16.9</b>



**TABLE 5: RSA GRADING OF YELLOW MAIZE ACCORDING TO GRADE (2020/21) (continue)**

Number of samples	Region	% Defective Kernels						% Total defective		% Foreign matter		% Other Colour		% Combined Deviations		% Pinked Kernels		% Diplodia Kernels		% Fusarium Kernels		% Cobrot Kernels				
		Above 6.35 mm sieve		Below 6.35 mm sieve		ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.			
		ave.	min.	max.	ave.																			min.	max.	
<b>GRADE: YM2</b>																										
3	Region 10	0.9	0.5	1.6	4.7	1.5	6.2	5.6	2.2	7.8	0.2	0.0	0.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.0	0.5	0.2	0.0	0.5
2	Region 11	3.0	0.7	5.2	4.9	4.8	5.1	7.9	5.5	10.3	0.2	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.0	0.8	0.4	0.0	0.8
1	Region 12	8.6	8.6	8.6	0.4	0.4	0.4	8.9	8.9	8.9	0.2	0.2	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.2	0.2	0.2	0.2	0.2
1	Region 13	1.3	1.3	1.3	2.3	2.3	2.3	3.6	3.6	3.6	0.5	0.5	0.5	0.1	0.1	0.1	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.1	0.1	0.1
1	Region 14	1.1	1.1	1.1	4.2	4.2	4.2	5.3	5.3	5.3	0.3	0.3	0.3	0.2	0.2	0.2	0.0	0.0	0.0	0.0	0.4	0.4	0.4	0.4	0.4	0.4
1	Region 16	0.9	0.9	0.9	2.8	2.8	2.8	3.7	3.7	3.7	0.4	0.4	0.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
1	Region 18	1.9	1.9	1.9	3.5	3.5	3.5	5.4	5.4	5.4	0.5	0.5	0.5	0.3	0.3	0.3	0.0	0.0	0.0	0.0	0.3	0.3	0.3	0.3	0.3	0.3
2	Region 19	1.3	0.9	1.7	5.4	4.5	6.2	6.7	6.2	7.1	0.2	0.0	0.4	0.8	0.0	1.6	0.0	0.0	0.0	0.0	0.1	0.0	0.1	0.1	0.0	0.1
1	Region 21	1.6	1.6	1.6	4.5	4.5	4.5	6.0	6.0	6.0	0.1	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.0	1.0	1.0	1.0	1.0	1.0
1	Region 22	1.4	1.4	1.4	4.9	4.9	4.9	6.2	6.2	6.2	0.0	0.0	0.0	0.1	0.1	0.1	0.0	0.0	0.0	0.0	0.7	0.7	0.7	0.7	0.7	0.7
1	Region 24	1.8	1.8	1.8	8.0	8.0	8.0	9.8	9.8	9.8	0.4	0.4	0.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
5	Region 25	2.9	1.1	5.3	4.2	1.6	5.3	7.1	3.2	10.7	0.1	0.0	0.4	0.2	0.0	0.7	0.0	0.0	0.0	0.0	0.3	0.0	1.1	0.3	0.0	1.1
3	Region 26	2.5	1.1	3.6	2.5	1.2	4.6	4.9	4.2	5.7	0.1	0.0	0.4	0.9	0.0	2.8	0.0	0.0	0.0	0.0	0.7	0.4	1.1	0.7	0.4	1.1
4	Region 27	0.7	0.2	1.1	6.2	4.1	8.6	6.8	5.0	8.9	0.1	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.2	0.1	0.0	0.2
5	Region 28	2.0	0.8	2.7	4.6	4.2	5.5	6.6	5.7	7.2	0.1	0.0	0.3	0.1	0.0	0.3	0.0	0.0	0.0	0.0	0.8	0.0	1.6	0.8	0.0	1.6
2	Region 30	1.9	1.9	2.0	2.3	0.4	4.2	4.2	2.2	6.2	0.2	0.0	0.5	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.9	0.7	1.1	0.9	0.7	1.1
1	Region 31	12.8	12.8	12.8	1.1	1.1	1.1	13.8	13.8	13.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	9.7	9.7	9.7	10.1	10.1	10.1
3	Region 32	2.3	1.2	3.8	2.8	1.1	5.2	5.1	3.0	6.3	0.1	0.0	0.4	1.4	0.0	4.1	0.0	0.0	0.0	0.0	0.6	0.2	1.2	0.6	0.2	1.2
2	Region 33	2.5	1.2	3.7	2.8	0.8	4.7	5.2	2.0	8.4	0.2	0.0	0.5	0.2	0.0	0.3	0.0	0.0	0.0	0.0	0.3	0.0	0.6	0.3	0.0	0.6
2	Region 34	1.9	1.6	2.2	4.4	4.1	4.7	6.3	5.7	6.9	0.3	0.1	0.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1.0	0.9	1.2	1.0	0.9	1.2
1	Region 36	9.5	9.5	9.5	0.1	0.1	0.1	9.6	9.6	9.6	0.4	0.4	0.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
<b>43</b>	<b>Ave. YM2</b>	<b>2.5</b>			<b>4.0</b>			<b>6.4</b>		<b>2.0</b>	<b>0.2</b>		<b>0.3</b>		<b>0.0</b>		<b>6.8</b>		<b>0.0</b>	<b>0.0</b>	<b>0.7</b>		<b>0.7</b>	<b>0.7</b>		<b>0.0</b>
	<b>Min. YM2</b>	<b>0.2</b>			<b>0.1</b>			<b>2.0</b>		<b>13.8</b>	<b>0.0</b>		<b>0.0</b>		<b>0.0</b>		<b>2.7</b>		<b>0.0</b>	<b>0.0</b>	<b>0.0</b>		<b>0.0</b>	<b>0.0</b>		<b>0.0</b>
	<b>Max. YM2</b>	<b>12.8</b>			<b>8.6</b>			<b>13.8</b>			<b>0.5</b>		<b>4.1</b>		<b>13.8</b>		<b>0.0</b>		<b>0.4</b>	<b>9.7</b>		<b>9.7</b>	<b>10.1</b>	<b>10.1</b>		<b>10.1</b>

Number of samples		TABLE 5: RSA GRADING OF YELLOW MAIZE ACCORDING TO GRADE (2020/21) (continue)																							
		% Defective Kernels		% Total defective		% Foreign matter		% Other Colour		% Combined Deviations		% Pinked Kernels		% Diplodia Kernels		% Fusarium Kernels		% Cobrot Kernels							
		Above 6.35 mm sieve		Below 6.35 mm sieve		ave.   min.   max.		ave.   min.   max.		ave.   min.   max.		ave.   min.   max.		ave.   min.   max.		ave.   min.   max.		ave.   min.   max.							
		ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.			
<b>GRADE: YM3</b>																									
1	Region 10	0.5	0.5	0.5	0.8	0.8	0.8	1.4	1.4	1.4	0.6	0.6	0.6	0.0	0.0	0.0	1.9	1.9	1.9	0.0	0.0	0.0	0.4	0.4	0.4
1	Region 21	0.2	0.2	0.2	3.8	3.8	3.8	4.1	4.1	4.1	0.6	0.6	0.6	0.0	0.0	0.0	4.7	4.7	4.7	0.0	0.0	0.0	0.0	0.0	0.0
1	Region 24	1.7	1.7	1.7	3.1	3.1	3.1	4.8	4.8	4.8	0.6	0.6	0.6	0.0	0.0	0.0	5.4	5.4	5.4	0.0	0.0	0.0	0.0	0.0	0.0
1	Region 29	2.2	2.2	2.2	1.3	1.3	1.3	3.4	3.4	3.4	0.6	0.6	0.6	0.0	0.0	0.0	4.1	4.1	4.1	0.0	0.0	0.0	0.8	0.8	0.8
1	Region 30	2.3	2.3	2.3	11.3	11.3	11.3	13.6	13.6	13.6	0.1	0.1	0.1	1.2	1.2	1.2	14.8	14.8	14.8	0.0	0.0	0.0	0.1	0.1	0.1
2	Region 33	20.4	18.4	22.4	2.3	2.1	2.5	22.7	20.5	24.9	0.3	0.0	0.5	0.4	0.0	0.8	23.3	21.0	25.6	0.0	0.0	0.0	0.5	0.0	1.0
1	Region 34	0.3	0.3	0.3	1.4	1.4	1.4	1.7	1.7	1.7	0.6	0.6	0.6	0.0	0.0	0.0	2.3	2.3	2.3	0.0	0.0	0.0	0.0	0.0	0.0
8	Ave. YM3	6.0			3.3			9.3			0.4			0.2			10.0			0.0			0.3		
	Min. YM3	0.2			0.8			1.4			0.0			0.0			1.9			0.0			0.0		
	Max. YM3	22.4			11.3			24.9			0.6			1.2			25.6			0.0			1.0		
<b>CLASS: COM</b>																									
1	Region 12	0.4	0.4	0.4	0.9	0.9	0.9	1.3	1.3	1.3	0.0	0.0	0.0	0.2	0.2	0.2	1.5	1.5	1.5	0.0	0.0	0.0	0.2	0.2	0.2
1	Region 14	3.5	3.5	3.5	1.7	1.7	1.7	5.2	5.2	5.2	0.1	0.1	0.1	8.5	8.5	8.5	13.8	13.8	13.8	0.0	0.0	0.0	1.1	1.1	1.1
1	Region 17	1.1	1.1	1.1	1.9	1.9	1.9	3.0	3.0	3.0	0.0	0.0	0.0	0.1	0.1	0.1	3.1	3.1	3.1	0.0	0.0	0.0	0.0	0.0	0.0
2	Region 18	2.7	1.5	3.9	3.9	3.5	4.2	6.6	5.0	8.1	2.1	1.4	2.8	0.1	0.0	0.2	8.8	6.5	11.1	0.0	0.0	0.0	1.3	0.9	1.7
2	Region 21	1.0	0.9	1.1	3.2	1.3	5.2	4.2	2.2	6.3	0.1	0.1	0.1	4.5	0.7	8.2	8.8	7.1	10.4	0.0	0.0	0.0	0.2	0.0	0.5
1	Region 22	1.2	1.2	1.2	5.6	5.6	5.6	6.8	6.8	6.8	0.0	0.0	0.0	0.0	0.0	0.0	6.8	6.8	6.8	0.0	0.0	0.0	0.0	0.0	0.0
1	Region 25	3.4	3.4	3.4	30.9	30.9	30.9	34.3	34.3	34.3	0.4	0.4	0.4	0.0	0.0	0.0	34.8	34.8	34.8	0.0	0.0	0.0	0.0	0.0	0.0
6	Region 26	3.5	0.9	6.7	2.9	0.8	5.3	6.4	3.7	10.5	0.5	0.0	1.2	0.9	0.0	5.4	7.9	4.6	13.0	0.0	0.0	0.0	0.7	0.2	1.4
2	Region 27	3.6	0.5	6.7	5.7	3.3	8.1	9.3	8.6	10.0	1.1	0.0	2.1	0.1	0.0	0.2	10.4	8.6	12.3	0.0	0.0	0.0	0.6	0.0	1.1
1	Region 28	2.9	2.9	2.9	6.4	6.4	6.4	9.2	9.2	9.2	1.3	1.3	1.3	0.0	0.0	0.0	10.5	10.5	10.5	0.0	0.0	0.0	0.9	0.9	0.9
4	Region 29	1.1	0.3	1.9	1.4	1.0	2.0	2.5	1.4	3.4	0.8	0.0	1.6	0.1	0.0	0.4	3.4	1.8	5.0	0.0	0.0	0.0	0.4	0.0	0.8
3	Region 30	2.1	1.0	3.1	1.7	1.6	1.7	3.7	2.6	4.8	0.2	0.0	0.4	0.0	0.0	0.0	3.9	2.6	4.8	0.0	0.0	0.0	0.7	0.2	1.7
4	Region 31	3.1	1.7	5.3	1.8	1.6	2.0	4.9	3.3	7.3	0.8	0.2	1.5	2.4	0.0	9.6	8.1	4.5	13.2	0.0	0.0	0.0	0.9	0.0	1.6
5	Region 32	3.7	2.3	6.5	2.2	0.9	4.0	5.9	4.3	7.8	0.7	0.0	1.4	0.0	0.0	0.0	6.5	4.8	7.8	0.0	0.0	0.0	1.0	0.0	2.1
5	Region 33	3.2	1.0	5.5	1.8	1.1	3.4	4.9	2.3	6.9	1.2	0.0	4.7	0.1	0.0	0.2	6.2	3.1	8.1	0.0	0.0	0.0	0.9	0.0	3.1
5	Region 34	1.7	1.2	2.5	2.5	0.6	4.7	4.2	2.2	6.8	0.7	0.0	1.5	0.2	0.0	0.6	5.0	3.2	8.3	0.0	0.0	0.0	0.3	0.2	0.5
2	Region 36	1.5	1.3	1.8	4.1	2.1	6.1	5.7	3.9	7.5	1.3	0.0	2.6	0.4	0.1	0.7	7.4	4.0	10.7	0.0	0.0	0.0	0.7	0.6	0.9



**TABLE 5: RSA GRADING OF YELLOW MAIZE ACCORDING TO GRADE (2020/21) (continue)**

Number of samples	Region	% Defective Kernels						% Total defective		% Foreign matter		% Other Colour		% Combined Deviations		% Pinked Kernels		% Diplodia Kernels		% Fusarium Kernels		% Cobrot Kernels	
		Above 6.35 mm sieve		Below 6.35 mm sieve		ave.	max.	ave.	max.	ave.	max.	ave.	max.	ave.	max.	ave.	max.	ave.	max.	ave.	max.	ave.	max.
		min.	max.	min.	max.																		
46	Ave. COM	2.5	3.2	0.3	0.6	5.8	0.7	0.0	4.7	0.8	0.0	9.6	7.3	1.5	0.0	0.0	0.0	0.0	0.0	0.7	0.0	0.7	0.0
	Min. COM			6.7	30.9	1.3	34.3																
	Max. COM																						
440	Ave. YM	2.1	2.1	0.1	0.0	4.2	0.1	0.0	4.7	0.2	0.0	9.6	4.5	0.2	0.0	0.0	0.0	0.0	0.6	0.0	0.6	0.0	
	Min. YM			22.4	30.9	0.2	34.3																
	Max. YM																						
1000	Ave. Maize	2.6	2.1	0.0	0.0	4.7	0.2	0.0	4.7	0.3	0.0	12.8	5.1	0.2	0.1	0.0	0.0	0.0	0.7	0.0	0.7	0.0	
	Min. Maize			55.1	30.9	0.2	56.9																
	Max. Maize																						

*The following yellow maize samples were downgraded to Class Other Maize due to the presence of poisonous seeds exceeding the maximum allowance	
Region	Number of Poisonous seeds (Crotonaria spp., Datura spp., Ricinus communis) Max. allowance 1 seed/1000 g
12	6 <i>Datura</i> spp.
17	6 <i>Datura</i> spp.
21	6 <i>Datura</i> spp.
22	12 <i>Datura</i> spp.
26	0
26	0
27	18 <i>Datura</i> spp.
29	6 <i>Datura</i> spp.
30	0
30	0
30	6 <i>Datura</i> spp.
31	0
31	0
32	12 <i>Datura</i> spp.
32	6 <i>Datura</i> spp.
32	0
33	6 <i>Datura</i> spp.
33	0
33	6 <i>Datura</i> spp.
34	0
34	6 <i>Datura</i> spp.
34	0
34	0
36	0


Number of Poisonous seeds (Argemone mexicana L., Convolvulus spp., Ipomoea purpurea Roth., Lolium temulentum, Xanthium spp.) Max. allowance 7 seeds/1000 g	
	0
	0
	0
	0
	12 <i>Xanthium Strumarium</i>
	12 <i>Xanthium Strumarium</i>
	0
	0
	12 <i>Xanthium Strumarium</i>
	12 <i>Xanthium Strumarium</i>
	0
	96 <i>Xanthium Strumarium</i>
	12 <i>Xanthium Strumarium</i>
	17 <i>Ipomoea purpurea</i> , 12 <i>Xanthium Strumarium</i>
	0
	12 <i>Xanthium Strumarium</i>
	6 <i>Convolvulus</i> spp.
	24 <i>Xanthium Strumarium</i>
	0
	6 <i>Ipomoea purpurea</i> , 12 <i>Xanthium Strumarium</i>
	0
	24 <i>Xanthium Strumarium</i>
	23 <i>Ipomoea purpurea</i>




**TABLE 7: GRADING QUALITY OF SOUTH AFRICAN  
WHITE AND YELLOW MAIZE 2011/12 - 2020/21**

Season	Number of samples	% Defective kernels above 6.35 mm sieve			% Defective kernels below 6.35 mm sieve			% Foreign matter			% Other colour			% Combined deviations		
		ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.
		<b>White Maize</b>														
2011/12	577	2.5	0.3	21.6	2.0	0.0	8.1	0.1	0.0	1.1	0.3	0.0	43.7	5.0	0.7	51.2
2012/13	508	2.6	0.0	20.8	2.0	0.2	11.4	0.1	0.0	1.5	0.3	0.0	6.5	4.9	1.0	22.4
2013/14	451	4.6	0.6	24.7	1.7	0.1	9.8	0.1	0.0	4.5	0.4	0.0	9.2	6.8	1.9	29.2
2014/15	485	3.1	0.0	30.0	2.2	0.0	25.5	0.1	0.0	1.2	0.4	0.0	9.6	5.8	0.0	35.3
2015/16	415	3.8	0.7	79.9	2.4	0.0	14.5	0.2	0.0	2.2	0.4	0.0	8.0	6.7	1.9	91.5
2016/17	549	2.9	0.3	25.5	1.8	0.1	12.7	0.2	0.0	6.9	0.2	0.0	7.0	5.1	1.1	36.7
2017/18	451	3.7	0.2	38.6	1.8	0.0	20.7	0.1	0.0	3.0	0.3	0.0	6.2	5.9	0.3	46.2
2018/19	404	6.8	0.2	88.5	1.8	0.0	19.0	0.2	0.0	4.2	0.5	0.0	12.9	9.3	0.6	96.8
2019/20	516	8.1	0.3	95.9	2.0	0.0	18.3	0.2	0.0	7.1	0.4	0.0	18.3	10.7	0.8	98.8
2020/21	560	3.0	0.0	55.1	2.1	0.0	21.1	0.2	0.0	3.9	0.4	0.0	12.8	5.6	0.5	58.7
<b>Weighted Average</b>		<b>4.0</b>			<b>2.0</b>			<b>0.1</b>			<b>0.4</b>			<b>6.5</b>		
<b>Minimum</b>		<b>0.0</b>			<b>0.0</b>			<b>0.0</b>			<b>0.0</b>			<b>0.0</b>		
<b>Maximum</b>				<b>95.9</b>			<b>25.5</b>			<b>7.1</b>			<b>43.7</b>			<b>98.8</b>
<b>Yellow Maize</b>																
2011/12	423	2.5	0.4	66.3	2.5	0.2	22.9	0.1	0.0	3.6	0.2	0.0	5.6	5.2	1.0	90.4
2012/13	492	2.0	0.2	23.1	2.5	0.1	14.0	0.1	0.0	1.8	0.2	0.0	8.4	4.8	0.8	25.0
2013/14	479	4.0	0.5	32.3	2.1	0.1	10.5	0.1	0.0	1.9	0.2	0.0	7.8	6.4	1.7	33.7
2014/15	515	3.1	0.6	23.0	2.7	0.0	19.0	0.1	0.0	2.5	0.3	0.0	13.6	6.2	0.6	34.4
2015/16	505	3.1	0.5	24.4	2.6	0.0	18.1	0.2	0.0	1.7	0.2	0.0	4.5	6.0	0.6	32.4
2016/17	451	2.4	0.4	24.3	2.0	0.0	27.4	0.2	0.0	2.8	0.2	0.0	6.9	4.7	1.3	33.5
2017/18	449	3.1	0.2	21.2	1.8	0.0	13.5	0.1	0.0	1.3	0.1	0.0	6.2	5.1	0.8	28.0
2018/19	404	3.7	0.0	36.9	1.7	0.0	9.5	0.1	0.0	3.4	0.2	0.0	7.4	5.7	0.7	38.9
2019/20	374	4.3	0.1	85.6	2.3	0.0	11.7	0.1	0.0	5.9	0.2	0.0	17.2	6.9	0.3	86.9
2020/21	440	2.1	0.1	22.4	2.1	0.0	30.9	0.1	0.0	4.7	0.2	0.0	9.6	4.5	0.2	34.8
<b>Weighted Average</b>		<b>3.0</b>			<b>2.2</b>			<b>0.1</b>			<b>0.2</b>			<b>5.5</b>		
<b>Minimum</b>		<b>0.0</b>			<b>0.0</b>			<b>0.0</b>			<b>0.0</b>			<b>0.2</b>		
<b>Maximum</b>				<b>85.6</b>			<b>30.9</b>			<b>5.9</b>			<b>17.2</b>			<b>90.4</b>
<b>White and Yellow Maize</b>																
2011/12	1000	2.5	0.3	66.3	2.2	0.0	22.9	0.1	0.0	3.6	0.3	0.0	43.7	5.1	0.7	90.4
2012/13	1000	2.0	0.0	23.1	2.3	0.1	14.0	0.1	0.0	1.8	0.3	0.0	8.4	4.9	0.8	25.0
2013/14	930	4.3	0.5	32.3	1.9	0.1	10.5	0.1	0.0	4.5	0.3	0.0	9.2	6.6	1.7	33.7
2014/15	1000	3.1	0.0	30.0	2.5	0.0	25.5	0.1	0.0	2.5	0.3	0.0	13.6	6.0	0.0	35.3
2015/16	920	3.4	0.5	79.9	2.5	0.0	18.1	0.2	0.0	2.2	0.3	0.0	8.0	6.3	0.6	91.5
2016/17	1000	2.6	0.3	25.5	1.9	0.0	27.4	0.2	0.0	6.9	0.2	0.0	7.0	4.9	1.1	36.7
2017/18	900	3.4	0.2	38.6	1.8	0.0	20.7	0.1	0.0	3.0	0.2	0.0	6.2	5.5	0.3	46.2
2018/19	808	5.3	0.0	88.5	1.8	0.0	19.0	0.1	0.0	4.2	0.4	0.0	12.9	7.5	0.6	96.8
2019/20	890	6.5	0.1	95.9	2.1	0.0	18.3	0.2	0.0	7.1	0.3	0.0	18.3	9.1	0.3	98.8
2020/21	1000	2.6	0.0	55.1	2.1	0.0	30.9	0.2	0.0	4.7	0.3	0.0	12.8	5.1	0.2	58.7
<b>Weighted Average</b>		<b>3.5</b>			<b>2.1</b>			<b>0.1</b>			<b>0.3</b>			<b>6.0</b>		
<b>Minimum</b>		<b>0.0</b>			<b>0.0</b>			<b>0.0</b>			<b>0.0</b>			<b>0.0</b>		
<b>Maximum</b>				<b>95.9</b>			<b>30.9</b>			<b>7.1</b>			<b>43.7</b>			<b>98.8</b>


GRAPH 34: PERCENTAGE DEFECTIVE KERNELS ABOVE THE 6.35 MM SIEVE OVER 10 SEASONS



GRAPH 35: PERCENTAGE DEFECTIVE KERNELS BELOW THE 6.35 MM SIEVE OVER 10 SEASONS



GRAPH 36: PERCENTAGE COMBINED DEVIATIONS OVER 10 SEASONS



**TABLE 8: USA GRADING OF WHITE MAIZE (2020/21)**

Number of samples	Region	Damaged kernels						% Broken corn and foreign material			Bushel weight (lbs)			% Other colour		
		% Heat damaged			% Total damaged			ave.	min.	max.	ave.	min.	max.	ave.	min.	max.
		ave.	min.	max.	ave.	min.	max.									
<b>GRADE: US No.1</b>																
5	Region 12	0.0	0.0	0.0	2.1	1.6	3.0	0.8	0.4	1.1	58.0	57.7	58.5	0.1	0.0	0.4
20	Region 13	0.0	0.0	0.0	2.1	1.2	3.0	0.7	0.1	1.4	57.6	56.7	59.6	0.1	0.0	1.0
21	Region 14	0.0	0.0	0.0	1.9	0.2	3.0	0.7	0.0	1.9	58.9	56.7	60.7	0.0	0.0	0.2
4	Region 16	0.0	0.0	0.0	1.9	1.0	2.6	0.7	0.5	1.4	58.6	58.4	58.9	0.0	0.0	0.0
18	Region 17	0.0	0.0	0.0	2.0	0.9	3.0	0.7	0.1	2.0	58.4	57.3	59.6	0.1	0.0	0.6
19	Region 18	0.0	0.0	0.0	1.6	0.5	3.0	0.6	0.1	1.9	58.1	56.1	60.1	0.1	0.0	0.4
15	Region 19	0.0	0.0	0.0	1.7	0.5	3.0	0.8	0.1	2.0	58.2	57.2	59.9	0.1	0.0	1.0
13	Region 20	0.0	0.0	0.0	1.5	0.7	2.8	0.3	0.2	0.6	58.7	57.7	59.9	0.1	0.0	0.5
26	Region 21	0.0	0.0	0.0	1.7	0.6	2.9	0.6	0.1	1.7	59.2	57.7	60.7	0.0	0.0	0.2
5	Region 22	0.0	0.0	0.0	2.2	1.3	3.0	0.9	0.4	1.5	59.3	57.3	60.8	0.0	0.0	0.0
32	Region 23	0.0	0.0	0.0	1.8	0.3	3.0	0.7	0.1	1.6	59.6	57.2	60.9	0.0	0.0	0.4
7	Region 24	0.0	0.0	0.0	2.0	0.7	3.0	0.7	0.3	1.3	59.2	57.1	60.0	0.1	0.0	0.7
2	Region 25	0.0	0.0	0.0	2.0	1.7	2.2	0.4	0.2	0.7	58.1	56.5	59.6	0.0	0.0	0.0
3	Region 26	0.0	0.0	0.0	1.4	1.0	1.6	0.6	0.0	1.4	59.4	58.6	60.1	0.3	0.1	0.5
3	Region 27	0.0	0.0	0.0	1.5	0.5	2.9	1.0	0.3	2.0	58.9	58.1	59.8	0.3	0.0	0.7
11	Region 28	0.0	0.0	0.0	1.7	0.6	2.9	0.3	0.0	0.5	60.5	59.0	63.0	0.3	0.0	0.8
15	Region 29	0.0	0.0	0.0	1.9	0.8	3.0	0.5	0.0	1.4	59.2	57.9	61.6	0.5	0.0	1.5
10	Region 30	0.0	0.0	0.0	2.5	1.5	3.0	0.4	0.1	1.2	59.6	57.5	60.7	0.6	0.0	1.8
5	Region 31	0.0	0.0	0.0	1.2	0.6	1.6	0.7	0.2	2.0	59.0	57.1	61.1	0.5	0.0	1.8
8	Region 32	0.0	0.0	0.0	1.7	0.7	2.5	0.6	0.2	1.6	59.2	57.2	61.4	0.3	0.0	1.1
23	Region 33	0.0	0.0	0.0	2.0	0.6	3.0	0.3	0.0	2.0	59.6	57.3	61.1	0.5	0.0	1.8
19	Region 34	0.0	0.0	0.0	2.0	0.9	3.0	0.6	0.1	1.7	59.3	56.5	61.3	0.2	0.0	1.2
4	Region 35	0.0	0.0	0.0	0.5	0.1	0.7	0.1	0.0	0.2	60.8	60.2	61.3	0.0	0.0	0.0
7	Region 36	0.0	0.0	0.0	1.8	1.1	2.9	0.5	0.0	1.6	59.7	58.3	61.3	0.1	0.0	0.2
<b>295</b>	<b>Ave. US No.1</b>	<b>0.0</b>			<b>1.8</b>			<b>0.6</b>			<b>59.0</b>			<b>0.2</b>		
	<b>Min. US No.1</b>	<b>0.0</b>			<b>0.1</b>			<b>0.0</b>			<b>56.1</b>			<b>0.0</b>		
	<b>Max. US No.1</b>	<b>0.0</b>			<b>3.0</b>			<b>2.0</b>			<b>63.0</b>			<b>1.8</b>		
<b>GRADE: US No.2</b>																
1	Region 11	0.0	-	-	1.4	-	-	0.6	-	-	54.3	-	-	0.3	-	-
6	Region 12	0.0	0.0	0.0	3.2	1.3	4.5	1.8	1.3	2.7	57.6	55.9	58.7	0.0	0.0	0.0
8	Region 13	0.0	0.0	0.0	3.7	3.2	4.4	1.1	0.1	1.6	57.5	55.5	59.2	0.1	0.0	0.4
12	Region 14	0.0	0.0	0.0	3.9	3.1	5.0	1.3	0.3	2.9	58.6	58.2	59.1	0.0	0.0	0.2
2	Region 16	0.1	0.0	0.2	3.0	2.0	4.0	0.5	0.4	0.6	58.7	58.6	58.7	0.0	0.0	0.0
2	Region 17	0.0	0.0	0.0	2.3	1.1	3.5	1.2	0.3	2.2	57.3	56.1	58.4	0.0	0.0	0.0
4	Region 18	0.0	0.0	0.0	3.5	3.1	4.4	0.7	0.2	2.0	58.5	57.9	59.1	0.5	0.0	0.8
6	Region 19	0.0	0.0	0.0	2.8	0.7	4.8	1.4	0.8	2.3	56.9	55.0	60.5	0.0	0.0	0.2
3	Region 20	0.0	0.0	0.0	4.0	3.3	5.0	0.9	0.6	1.3	58.8	58.6	58.9	0.4	0.0	0.6
1	Region 21	0.0	-	-	1.3	-	-	2.3	-	-	58.2	-	-	0.2	-	-
5	Region 22	0.0	0.0	0.0	3.7	3.3	4.4	1.5	1.1	2.4	58.9	58.2	59.4	0.0	0.0	0.0
6	Region 23	0.0	0.0	0.2	2.9	1.2	4.0	1.1	0.1	2.5	59.8	57.9	60.4	0.0	0.0	0.0
3	Region 24	0.0	0.0	0.0	4.0	3.8	4.4	0.4	0.3	0.5	59.7	59.1	60.7	0.0	0.0	0.0
2	Region 26	0.0	0.0	0.0	4.3	3.6	5.0	0.2	0.1	0.4	58.5	57.2	59.8	0.6	0.0	1.3
1	Region 27	0.0	-	-	1.4	-	-	2.4	-	-	58.8	-	-	0.2	-	-
2	Region 28	0.1	0.0	0.2	3.0	1.1	4.9	0.5	0.4	0.6	60.1	59.1	61.0	0.9	0.4	1.3
11	Region 29	0.0	0.0	0.0	3.9	3.1	4.8	0.5	0.1	1.2	59.2	57.4	60.7	0.2	0.0	0.9
12	Region 30	0.0	0.0	0.0	3.8	3.1	4.6	0.4	0.0	0.8	60.1	59.1	61.3	0.7	0.0	1.7
6	Region 31	0.0	0.0	0.0	3.8	3.1	4.5	0.5	0.1	0.9	60.0	58.9	61.0	0.6	0.0	1.4
9	Region 32	0.0	0.0	0.0	3.8	3.1	4.8	0.8	0.3	1.7	60.1	57.2	61.3	0.9	0.0	1.9
16	Region 33	0.0	0.0	0.0	3.7	3.1	4.9	0.3	0.0	1.3	60.3	58.5	61.8	0.4	0.0	1.5

**TABLE 8: USA GRADING OF WHITE MAIZE (2020/21) (continue)**

Number of samples	Region	Damaged kernels						% Broken corn and foreign material			Bushel weight (lbs)			% Other colour		
		% Heat damaged			% Total damaged			ave.	min.	max.	ave.	min.	max.	ave.	min.	max.
		ave.	min.	max.	ave.	min.	max.									
<b>GRADE: US No.2</b>																
8	Region 34	0.0	0.0	0.0	3.6	2.2	4.6	0.7	0.2	1.1	59.6	54.1	61.3	0.2	0.0	0.4
1	Region 35	0.0	-	-	3.8	-	-	0.0	-	-	61.0	-	-	0.0	-	-
<b>135</b>	<b>Ave. US No.2</b>	<b>0.0</b>			<b>3.6</b>			<b>0.8</b>			<b>59.2</b>			<b>0.3</b>		
	<b>Min. US No.2</b>	<b>0.0</b>			<b>0.7</b>			<b>0.0</b>			<b>54.1</b>			<b>0.0</b>		
	<b>Max. US No.2</b>	<b>0.2</b>			<b>5.0</b>			<b>2.9</b>			<b>61.8</b>			<b>1.9</b>		
<b>GRADE: US No.3</b>																
1	Region 11	0.0	-	-	6.1	-	-	1.2	-	-	59.4	-	-	0.0	-	-
2	Region 12	0.0	0.0	0.0	5.2	4.9	5.5	2.9	2.1	3.7	58.0	57.7	58.2	0.0	0.0	0.0
3	Region 14	0.0	0.0	0.0	5.5	5.3	5.6	0.7	0.4	0.8	58.7	58.6	58.9	0.0	0.0	0.0
3	Region 17	0.0	0.0	0.0	5.4	5.1	5.7	1.0	0.9	1.1	58.4	58.2	58.9	0.0	0.0	0.0
1	Region 18	0.0	-	-	1.2	-	-	3.4	-	-	58.5	-	-	0.0	-	-
4	Region 19	0.1	0.0	0.3	3.9	2.5	4.8	2.6	0.7	4.0	56.2	53.2	57.5	0.0	0.0	0.0
2	Region 21	0.0	0.0	0.0	6.0	5.1	7.0	0.5	0.5	0.5	58.4	57.9	58.9	0.0	0.0	0.0
4	Region 23	0.0	0.0	0.0	2.9	1.5	5.2	2.9	1.4	3.6	59.9	59.0	60.5	0.0	0.0	0.0
2	Region 24	0.0	0.0	0.0	3.6	1.0	6.2	2.0	0.6	3.4	57.3	54.9	59.6	0.0	0.0	0.0
1	Region 27	0.0	-	-	0.4	-	-	3.5	-	-	59.1	-	-	0.3	-	-
5	Region 30	0.0	0.0	0.2	4.7	0.0	6.6	2.0	0.4	3.8	59.8	58.6	60.9	0.5	0.0	1.8
5	Region 31	0.0	0.0	0.0	6.2	5.6	6.9	0.8	0.5	1.1	60.4	59.6	61.4	0.7	0.0	1.5
5	Region 32	0.0	0.0	0.0	6.1	5.1	7.0	0.6	0.4	0.8	60.3	58.2	62.1	0.3	0.0	0.9
8	Region 33	0.0	0.0	0.2	5.7	5.1	6.5	0.7	0.1	1.7	59.4	56.9	61.6	0.4	0.0	1.0
3	Region 34	0.0	0.0	0.0	4.4	1.5	6.2	0.5	0.0	1.1	57.3	53.4	60.4	0.1	0.0	0.2
8	Region 36	0.1	0.0	0.4	5.7	4.0	6.8	0.4	0.1	0.6	60.5	59.8	61.1	0.2	0.0	0.5
<b>57</b>	<b>Ave. US No.3</b>	<b>0.0</b>			<b>5.0</b>			<b>1.3</b>			<b>59.2</b>			<b>0.2</b>		
	<b>Min. US No.3</b>	<b>0.0</b>			<b>0.0</b>			<b>0.0</b>			<b>53.2</b>			<b>0.0</b>		
	<b>Max. US No.3</b>	<b>0.4</b>			<b>7.0</b>			<b>4.0</b>			<b>62.1</b>			<b>1.8</b>		
<b>GRADE: US No.4</b>																
1	Region 13	0.0	-	-	8.6	-	-	1.7	-	-	56.3	-	-	0.0	-	-
4	Region 14	0.3	0.0	1.0	7.2	4.7	8.4	0.8	0.6	1.1	59.3	57.7	60.1	0.0	0.0	0.0
2	Region 19	0.0	0.0	0.0	2.7	2.5	2.9	4.6	4.4	4.8	56.6	55.5	57.7	0.4	0.0	0.8
1	Region 20	0.0	-	-	9.9	-	-	0.4	-	-	58.4	-	-	1.5	-	-
2	Region 21	0.0	0.0	0.0	0.6	0.0	1.1	4.4	4.4	4.5	57.0	56.5	57.5	0.3	0.0	0.6
1	Region 23	0.0	-	-	8.9	-	-	0.7	-	-	60.2	-	-	0.0	-	-
1	Region 24	0.8	-	-	3.9	-	-	0.8	-	-	59.6	-	-	0.0	-	-
1	Region 25	0.0	-	-	2.6	-	-	4.9	-	-	59.1	-	-	1.0	-	-
1	Region 30	0.0	-	-	7.6	-	-	2.0	-	-	59.8	-	-	0.0	-	-
2	Region 32	0.0	0.0	0.0	8.1	8.0	8.2	0.7	0.4	1.0	60.7	60.3	61.1	0.9	0.4	1.4
2	Region 33	0.0	0.0	0.0	8.5	7.5	9.5	0.6	0.4	0.7	58.4	57.7	59.0	0.3	0.0	0.7
1	Region 35	0.0	-	-	8.2	-	-	0.9	-	-	59.6	-	-	0.0	-	-
<b>19</b>	<b>Ave. US No.4</b>	<b>0.1</b>			<b>6.2</b>			<b>1.9</b>			<b>58.7</b>			<b>0.3</b>		
	<b>Min. US No.4</b>	<b>0.0</b>			<b>0.0</b>			<b>0.4</b>			<b>55.5</b>			<b>0.0</b>		
	<b>Max. US No.4</b>	<b>1.0</b>			<b>9.9</b>			<b>4.9</b>			<b>61.1</b>			<b>1.5</b>		
<b>GRADE: US No.5</b>																
1	Region 12	0.0	-	-	4.2	-	-	5.2	-	-	57.7	-	-	0.0	-	-
1	Region 16	0.0	-	-	12.1	-	-	1.2	-	-	59.7	-	-	0.2	-	-
3	Region 17	0.0	0.0	0.0	12.4	11.0	14.4	0.5	0.1	0.7	58.0	56.3	60.3	0.0	0.0	0.0
3	Region 18	0.6	0.0	1.7	6.1	1.9	12.6	4.0	0.9	6.4	58.4	57.4	59.4	0.3	0.0	0.5
1	Region 19	0.0	-	-	13.1	-	-	1.1	-	-	56.6	-	-	0.9	-	-
1	Region 21	0.0	-	-	1.2	-	-	5.5	-	-	57.4	-	-	0.0	-	-
2	Region 22	1.4	1.3	1.5	7.7	7.1	8.3	4.2	3.2	5.2	57.9	56.4	59.5	0.7	0.0	1.4

**TABLE 8: USA GRADING OF WHITE MAIZE (2020/21) (continue)**

Number of samples	Region	Damaged kernels						% Broken corn and foreign material			Bushel weight (lbs)			% Other colour		
		% Heat damaged			% Total damaged			ave.	min.	max.	ave.	min.	max.	ave.	min.	max.
		ave.	min.	max.	ave.	min.	max.									
<b>GRADE: US No.5</b>																
1	23	0.0	-	-	10.9	-	-	1.9	-	-	58.3	-	-	0.0	-	-
1	29	0.0	-	-	12.6	-	-	0.0	-	-	59.5	-	-	0.0	-	-
1	32	0.0	-	-	10.2	-	-	0.5	-	-	59.1	-	-	1.4	-	-
1	35	1.1	-	-	3.8	-	-	0.1	-	-	59.9	-	-	0.2	-	-
1	36	1.4	-	-	5.6	-	-	0.7	-	-	60.2	-	-	0.4	-	-
<b>17</b>	<b>Ave. US No.5</b>	<b>0.4</b>			<b>8.5</b>			<b>2.2</b>			<b>58.4</b>			<b>0.3</b>		
	<b>Min. US No.5</b>	<b>0.0</b>			<b>1.2</b>			<b>0.0</b>			<b>56.3</b>			<b>0.0</b>		
	<b>Max. US no.5</b>	<b>1.7</b>			<b>14.4</b>			<b>6.4</b>			<b>60.3</b>			<b>1.4</b>		
<b>GRADE: Sample Grade</b>																
3	17	0.6	0.0	1.8	17.2	15.2	20.8	0.7	0.1	1.8	56.5	53.8	58.0	0.0	0.0	0.1
1	19	0.0	-	-	6.4	-	-	7.4	-	-	57.9	-	-	0.0	-	-
4	21	0.0	0.0	0.0	1.8	0.7	2.3	7.8	1.3	11.8	57.7	56.9	59.8	0.0	0.0	0.0
1	22	0.0	-	-	11.2	-	-	14.0	-	-	55.6	-	-	0.6	-	-
3	23	1.5	0.0	4.4	13.1	4.8	19.4	3.1	0.4	7.7	59.2	59.0	59.4	0.1	0.0	0.2
2	26	0.0	0.0	0.0	4.2	0.4	8.0	14.1	9.6	18.6	57.9	57.4	58.4	0.8	0.4	1.3
1	31	0.0	-	-	55.4	-	-	1.8	-	-	56.0	-	-	0.6	-	-
1	32	0.0	-	-	1.5	-	-	2.6	-	-	59.7	-	-	0.0	-	-
3	36	0.7	0.0	1.1	4.9	3.0	6.3	0.9	0.1	2.3	58.8	58.2	59.7	0.7	0.1	1.8
<b>19</b>	<b>Ave. Sample Grade</b>	<b>0.4</b>			<b>10.3</b>			<b>5.2</b>			<b>57.9</b>			<b>0.3</b>		
	<b>Min. Sample Grade</b>	<b>0.0</b>			<b>0.4</b>			<b>0.1</b>			<b>53.8</b>			<b>0.0</b>		
	<b>Max. Sample Grade</b>	<b>4.4</b>			<b>55.4</b>			<b>18.6</b>			<b>59.8</b>			<b>1.8</b>		
<b>GRADE: Mixed Corn</b>																
1	16	0.0	-	-	1.0	-	-	0.8	-	-	59.5	-	-	4.0	-	-
2	18	0.0	0.0	0.0	1.0	0.6	1.4	0.6	0.4	0.7	56.6	56.2	57.0	2.3	2.2	2.4
1	26	0.0	-	-	1.1	-	-	0.2	-	-	59.0	-	-	5.7	-	-
6	30	0.0	0.0	0.0	4.0	2.1	5.3	0.7	0.3	1.6	59.6	57.0	62.2	4.3	2.5	6.7
3	31	0.0	0.0	0.0	4.6	2.0	9.0	0.5	0.3	0.6	58.1	57.3	58.8	5.1	2.5	7.9
5	33	0.0	0.0	0.0	3.0	1.8	4.5	1.4	0.3	2.3	58.0	55.8	58.9	5.6	2.3	12.8
<b>18</b>	<b>Ave. Mixed Corn</b>	<b>0.0</b>			<b>3.2</b>			<b>0.8</b>			<b>58.5</b>			<b>4.6</b>		
	<b>Min. Mixed Corn</b>	<b>0.0</b>			<b>0.6</b>			<b>0.2</b>			<b>55.8</b>			<b>2.2</b>		
	<b>Max. Mixed Corn</b>	<b>0.0</b>			<b>9.0</b>			<b>2.3</b>			<b>62.2</b>			<b>12.8</b>		
<b>560</b>	<b>Ave. WM</b>	<b>0.0</b>			<b>3.3</b>			<b>1.0</b>			<b>59.0</b>			<b>0.4</b>		
	<b>Min. WM</b>	<b>0.0</b>			<b>0.0</b>			<b>0.0</b>			<b>53.2</b>			<b>0.0</b>		
	<b>Max. WM</b>	<b>4.4</b>			<b>55.4</b>			<b>18.6</b>			<b>63.0</b>			<b>12.8</b>		
<b>1000</b>	<b>Ave. Maize</b>	<b>0.0</b>			<b>2.8</b>			<b>0.9</b>			<b>59.2</b>			<b>0.3</b>		
	<b>Min. Maize</b>	<b>0.0</b>			<b>0.0</b>			<b>0.0</b>			<b>53.2</b>			<b>0.0</b>		
	<b>Max. Maize</b>	<b>4.4</b>			<b>55.4</b>			<b>20.2</b>			<b>64.1</b>			<b>12.8</b>		

\*The following white maize samples were downgraded to Sample Grade due to the number of poisonous seeds exceeding the maximum allowance

Region	Cockleburs ( <i>Xanthium</i> spp.) exceeding 8 seeds
21	24 <i>Xanthium Strumarium</i>
32	10 <i>Xanthium Strumarium</i>
36	24 <i>Xanthium Strumarium</i>
36	12 <i>Xanthium Strumarium</i>
36	12 <i>Xanthium Strumarium</i>

**TABLE 9: USA GRADING OF YELLOW MAIZE (2020/21)**

Number of samples	Region	Damaged kernels						% Broken corn and foreign material			Bushel weight (lbs)			% Other colour		
		% Heat damaged			% Total damaged			ave.	min.	max.	ave.	min.	max.	ave.	min.	max.
		ave.	min.	max.	ave.	min.	max.									
<b>GRADE: US No.1</b>																
23	Region 10	0.0	0.0	0.0	1.0	0.3	2.6	0.4	0.0	1.3	61.3	58.6	64.1	0.0	0.0	0.2
13	Region 11	0.0	0.0	0.0	1.4	0.2	2.7	0.3	0.0	0.6	61.4	59.4	62.2	0.0	0.0	0.2
1	Region 12	0.0	-	-	0.4	-	-	0.0	-	-	59.5	-	-	0.2	-	-
5	Region 13	0.0	0.0	0.0	1.7	0.4	2.8	0.5	0.2	1.0	59.7	57.9	60.9	0.2	0.0	0.8
6	Region 14	0.0	0.0	0.0	0.9	0.0	2.2	0.6	0.1	1.6	59.8	58.2	61.4	0.1	0.0	0.4
1	Region 16	0.0	-	-	1.1	-	-	0.7	-	-	59.6	-	-	0.0	-	-
4	Region 17	0.0	0.0	0.0	1.6	0.9	2.4	0.7	0.1	1.6	60.0	58.8	62.0	0.1	0.0	0.1
7	Region 18	0.0	0.0	0.0	1.8	0.3	2.6	0.5	0.2	0.9	58.9	56.0	60.3	0.3	0.0	1.5
7	Region 19	0.0	0.0	0.0	1.6	0.3	2.7	0.7	0.4	1.2	58.5	57.0	61.3	0.1	0.0	0.2
3	Region 20	0.0	0.0	0.0	1.5	0.9	2.4	0.4	0.2	0.7	60.1	59.3	61.0	0.0	0.0	0.0
5	Region 21	0.0	0.0	0.0	1.4	0.4	2.2	1.2	0.6	1.9	58.9	57.4	60.6	0.1	0.0	0.6
1	Region 22	0.0	-	-	2.1	-	-	0.9	-	-	60.6	-	-	0.0	-	-
4	Region 23	0.0	0.0	0.0	0.9	0.6	1.3	0.5	0.4	0.7	59.5	58.6	60.4	0.2	0.0	0.5
4	Region 24	0.0	0.0	0.0	1.5	0.7	2.3	0.8	0.3	1.9	60.0	58.5	60.8	0.0	0.0	0.0
18	Region 25	0.0	0.0	0.0	1.6	0.7	2.8	0.5	0.0	1.5	58.5	57.8	59.5	0.0	0.0	0.3
12	Region 26	0.0	0.0	0.0	1.5	0.1	2.9	0.7	0.1	1.7	58.5	56.1	60.0	0.0	0.0	0.1
5	Region 27	0.0	0.0	0.0	0.9	0.5	1.4	1.0	0.5	1.5	58.5	57.9	59.6	0.0	0.0	0.0
32	Region 28	0.0	0.0	0.0	1.4	0.3	3.0	0.6	0.0	2.0	58.8	56.0	61.2	0.0	0.0	0.3
31	Region 29	0.0	0.0	0.0	1.6	0.4	3.0	0.6	0.0	1.9	59.7	57.7	61.7	0.1	0.0	0.7
27	Region 30	0.0	0.0	0.0	2.0	0.7	3.0	0.5	0.0	1.4	59.4	57.7	61.0	0.0	0.0	0.6
30	Region 31	0.0	0.0	0.0	1.7	0.4	2.8	0.5	0.1	1.2	60.2	57.7	62.2	0.0	0.0	0.5
30	Region 32	0.0	0.0	0.0	2.1	0.8	2.9	0.5	0.1	1.8	60.1	57.1	62.4	0.1	0.0	4.1
16	Region 33	0.0	0.0	0.0	1.6	0.6	2.8	0.4	0.0	1.5	59.3	57.6	61.9	0.1	0.0	0.3
17	Region 34	0.0	0.0	0.0	1.6	0.1	3.0	0.6	0.1	1.8	59.7	57.6	61.3	0.0	0.0	0.2
6	Region 35	0.0	0.0	0.0	1.6	1.1	2.2	0.2	0.0	0.5	60.0	57.2	62.1	0.0	0.0	0.0
13	Region 36	0.0	0.0	0.0	1.9	0.5	2.7	0.3	0.0	0.6	59.5	58.3	61.8	0.2	0.0	0.9
<b>321</b>	<b>Ave. US No.1</b>	<b>0.0</b>			<b>1.6</b>			<b>0.5</b>			<b>59.7</b>			<b>0.1</b>		
	<b>Min. US No.1</b>	<b>0.0</b>			<b>0.0</b>			<b>0.0</b>			<b>56.0</b>			<b>0.0</b>		
	<b>Max. US No.1</b>	<b>0.0</b>			<b>3.0</b>			<b>2.0</b>			<b>64.1</b>			<b>4.1</b>		
<b>GRADE: US No.2</b>																
4	Region 10	0.0	0.0	0.0	3.0	2.1	3.5	1.0	0.3	2.2	59.9	57.3	61.5	0.0	0.0	0.0
1	Region 11	0.0	-	-	1.0	-	-	0.9	-	-	55.6	-	-	0.0	-	-
1	Region 13	0.0	-	-	3.6	-	-	0.2	-	-	60.6	-	-	0.0	-	-
1	Region 14	0.0	-	-	3.5	-	-	0.4	-	-	60.3	-	-	0.2	-	-
1	Region 18	0.0	-	-	1.7	-	-	2.6	-	-	58.4	-	-	0.0	-	-
1	Region 19	0.0	-	-	3.0	-	-	2.3	-	-	55.3	-	-	0.3	-	-
3	Region 20	0.0	0.0	0.0	3.5	3.2	3.7	0.3	0.3	0.4	60.1	59.1	61.5	0.0	0.0	0.0
1	Region 21	0.0	-	-	4.6	-	-	0.1	-	-	59.1	-	-	0.7	-	-
4	Region 22	0.0	0.0	0.0	3.6	1.4	5.0	0.8	0.1	2.5	60.1	58.9	60.8	0.0	0.0	0.0
1	Region 24	0.0	-	-	1.8	-	-	1.3	-	-	55.8	-	-	0.0	-	-
3	Region 25	0.0	0.0	0.0	3.4	3.3	3.6	1.7	1.1	2.1	57.4	57.2	57.7	0.2	0.0	0.7
4	Region 26	0.0	0.0	0.0	3.9	3.4	4.3	0.1	0.0	0.1	58.1	56.5	60.0	1.4	0.3	2.8
3	Region 27	0.0	0.0	0.0	0.5	0.4	0.7	2.0	0.5	2.9	57.1	55.5	58.6	0.0	0.0	0.0
1	Region 28	0.0	-	-	3.2	-	-	0.3	-	-	58.4	-	-	0.0	-	-
13	Region 30	0.0	0.0	0.0	3.4	1.1	4.9	0.5	0.2	1.1	58.8	55.6	60.5	0.0	0.0	0.5
9	Region 31	0.0	0.0	0.0	4.0	3.1	4.6	0.4	0.1	0.9	60.7	59.2	61.6	0.0	0.0	0.2
8	Region 32	0.0	0.0	0.0	3.5	2.5	4.7	1.1	0.3	2.7	59.3	57.6	60.6	0.0	0.0	0.0
7	Region 33	0.0	0.0	0.0	3.7	3.1	4.6	0.4	0.1	1.1	59.2	58.6	60.0	0.0	0.0	0.2



**TABLE 9: USA GRADING OF YELLOW MAIZE (2020/21) (continue)**

Number of samples	Region	Damaged kernels						% Broken corn and foreign material			Bushel weight (lbs)			% Other colour		
		% Heat damaged			% Total damaged			ave.	min.	max.	ave.	min.	max.	ave.	min.	max.
		ave.	min.	max.	ave.	min.	max.									
<b>GRADE: US No.2</b>																
5	Region 34	0.0	0.0	0.0	3.2	1.2	4.6	1.2	0.4	2.3	58.3	55.1	61.8	0.2	0.0	0.6
5	Region 36	0.0	0.0	0.0	3.6	3.1	4.4	0.4	0.0	1.0	59.0	58.4	59.3	0.1	0.0	0.4
<b>76</b>	<b>Ave. US No.2</b>	<b>0.0</b>			<b>3.4</b>			<b>0.8</b>			<b>59.0</b>			<b>0.1</b>		
	<b>Min. US No.2</b>	<b>0.0</b>			<b>0.4</b>			<b>0.0</b>			<b>55.1</b>			<b>0.0</b>		
	<b>Max. US No.2</b>	<b>0.0</b>			<b>5.0</b>			<b>2.9</b>			<b>61.8</b>			<b>2.8</b>		
<b>GRADE: US No.3</b>																
1	Region 11	0.0	-	-	6.2	-	-	0.9	-	-	60.4	-	-	0.0	-	-
1	Region 19	0.0	-	-	1.0	-	-	3.9	-	-	58.2	-	-	1.6	-	-
2	Region 22	0.0	0.0	0.0	3.7	1.4	6.0	1.7	0.1	3.2	59.7	58.9	60.5	0.1	0.0	0.1
2	Region 25	0.1	0.0	0.2	5.3	5.2	5.3	2.3	2.2	2.4	57.3	57.2	57.4	0.1	0.0	0.3
1	Region 27	0.0	-	-	0.4	-	-	3.2	-	-	58.4	-	-	0.0	-	-
3	Region 30	0.0	0.0	0.0	4.6	2.8	5.7	1.9	0.3	3.6	60.1	59.3	60.7	0.1	0.0	0.4
1	Region 31	0.0	-	-	6.3	-	-	1.1	-	-	62.3	-	-	0.0	-	-
1	Region 32	0.0	-	-	6.6	-	-	0.3	-	-	60.1	-	-	0.0	-	-
2	Region 33	0.0	0.0	0.0	5.5	5.4	5.6	0.2	0.1	0.3	57.7	56.5	58.9	0.0	0.0	0.0
<b>14</b>	<b>Ave. US No.3</b>	<b>0.0</b>			<b>4.5</b>			<b>1.7</b>			<b>59.2</b>			<b>0.2</b>		
	<b>Min. US No.3</b>	<b>0.0</b>			<b>0.4</b>			<b>0.1</b>			<b>56.5</b>			<b>0.0</b>		
	<b>Max. US No.3</b>	<b>0.2</b>			<b>6.6</b>			<b>3.9</b>			<b>62.3</b>			<b>1.6</b>		
<b>GRADE: US No.4</b>																
1	Region 12	0.0	-	-	8.7	-	-	0.2	-	-	58.4	-	-	0.0	-	-
1	Region 20	0.6	-	-	1.5	-	-	0.8	-	-	59.6	-	-	0.0	-	-
1	Region 24	0.0	-	-	2.5	-	-	4.3	-	-	57.5	-	-	0.0	-	-
1	Region 26	0.0	-	-	5.4	-	-	4.8	-	-	59.1	-	-	0.1	-	-
1	Region 27	0.0	-	-	6.9	-	-	4.3	-	-	56.0	-	-	0.2	-	-
1	Region 28	0.0	-	-	2.9	-	-	4.6	-	-	57.0	-	-	0.0	-	-
2	Region 36	0.3	0.0	0.5	5.5	1.5	9.5	2.7	0.4	4.9	58.4	58.2	58.6	0.3	0.0	0.7
<b>8</b>	<b>Ave. US No.4</b>	<b>0.1</b>			<b>4.9</b>			<b>3.0</b>			<b>58.1</b>			<b>0.1</b>		
	<b>Min. US No.4</b>	<b>0.0</b>			<b>1.5</b>			<b>0.2</b>			<b>56.0</b>			<b>0.0</b>		
	<b>Max. US No.4</b>	<b>0.6</b>			<b>9.5</b>			<b>4.9</b>			<b>59.6</b>			<b>0.7</b>		
<b>GRADE: US No.5</b>																
1	10	1.7	-	-	1.8	-	-	0.8	-	-	61.1	-	-	0.0	-	-
1	18	0.0	-	-	4.0	-	-	5.4	-	-	58.2	-	-	0.2	-	-
1	31	0.0	-	-	12.8	-	-	0.4	-	-	58.5	-	-	0.0	-	-
1	33	0.0	-	-	1.7	-	-	5.2	-	-	61.7	-	-	0.0	-	-
<b>4</b>	<b>Ave. US No.5</b>	<b>0.4</b>			<b>5.1</b>			<b>2.9</b>			<b>59.9</b>			<b>0.0</b>		
	<b>Min. US No.5</b>	<b>0.0</b>			<b>1.7</b>			<b>0.4</b>			<b>58.2</b>			<b>0.0</b>		
	<b>Max. US No.5</b>	<b>1.7</b>			<b>12.8</b>			<b>5.4</b>			<b>61.7</b>			<b>0.2</b>		
<b>GRADE: Sample Grade</b>																
1	Region 25	0.2	-	-	12.1	-	-	20.2	-	-	54.9	-	-	0.0	-	-
2	Region 26	0.0	0.0	0.0	3.5	2.1	4.9	1.0	0.3	1.6	58.1	57.0	59.1	0.0	0.0	0.0
2	Region 30	0.0	0.0	0.0	2.2	1.0	3.3	0.5	0.2	0.7	59.7	59.6	59.8	0.0	0.0	0.0
2	Region 31	0.0	0.0	0.0	4.8	4.0	5.6	1.1	0.5	1.6	59.3	59.2	59.4	0.0	0.0	0.0
1	Region 32	0.0	-	-	4.1	-	-	0.3	-	-	59.6	-	-	0.0	-	-
3	Region 33	0.8	0.0	2.3	15.3	4.7	22.7	0.7	0.4	1.0	57.9	55.4	60.5	0.3	0.0	0.8
2	Region 34	0.0	0.0	0.0	1.5	1.3	1.7	1.0	0.8	1.1	58.1	56.8	59.3	0.0	0.0	0.0
<b>13</b>	<b>Ave. Sample Grade</b>	<b>0.2</b>			<b>6.6</b>			<b>2.3</b>			<b>58.3</b>			<b>0.1</b>		
	<b>Min. Sample Grade</b>	<b>0.0</b>			<b>1.0</b>			<b>0.2</b>			<b>54.9</b>			<b>0.0</b>		
	<b>Max. Sample Grade</b>	<b>2.3</b>			<b>22.7</b>			<b>20.2</b>			<b>60.5</b>			<b>0.8</b>		

**TABLE 9: USA GRADING OF YELLOW MAIZE (2020/21) (continue)**

Number of samples	Region	Damaged kernels						% Broken corn and foreign material			Bushel weight (lbs)			% Other colour		
		% Heat damaged			% Total damaged			ave.	min.	max.	ave.	min.	max.	ave.	min.	max.
		ave.	min.	max.	ave.	min.	max.									
<b>CLASS: Mixed Corn</b>																
1	Region 14	0.0	-	-	3.7	-	-	0.7	-	-	59.0	-	-	8.5	-	-
1	Region 21	0.0	-	-	1.0	-	-	0.4	-	-	59.1	-	-	8.2	-	-
1	Region 26	0.0	-	-	6.8	-	-	0.0	-	-	57.7	-	-	5.4	-	-
1	Region 31	0.0	-	-	1.9	-	-	0.5	-	-	58.2	-	-	9.6	-	-
<b>4</b>	<b>Ave. Mixed Corn</b>	<b>0.0</b>			<b>3.3</b>			<b>0.4</b>			<b>58.5</b>			<b>7.9</b>		
	<b>Min. Mixed Corn</b>	<b>0.0</b>			<b>1.0</b>			<b>0.0</b>			<b>57.7</b>			<b>5.4</b>		
	<b>Max. Mixed Corn</b>	<b>0.0</b>			<b>6.8</b>			<b>0.7</b>			<b>59.1</b>			<b>9.6</b>		
<b>440</b>	<b>Ave. YM</b>	<b>0.0</b>			<b>2.2</b>			<b>0.7</b>			<b>59.5</b>			<b>0.2</b>		
	<b>Min. YM</b>	<b>0.0</b>			<b>0.0</b>			<b>0.0</b>			<b>54.9</b>			<b>0.0</b>		
	<b>Max. YM</b>	<b>2.3</b>			<b>22.7</b>			<b>20.2</b>			<b>64.1</b>			<b>9.6</b>		
<b>1000</b>	<b>Ave. Maize</b>	<b>0.0</b>			<b>2.8</b>			<b>0.9</b>			<b>59.2</b>			<b>0.3</b>		
	<b>Min. Maize</b>	<b>0.0</b>			<b>0.0</b>			<b>0.0</b>			<b>53.2</b>			<b>0.0</b>		
	<b>Max. Maize</b>	<b>4.4</b>			<b>55.4</b>			<b>20.2</b>			<b>64.1</b>			<b>12.8</b>		

**\*The following yellow maize samples were downgraded to Sample Grade due to the number of poisonous seeds exceeding the maximum allowance**

Region	Cockleburs ( <i>Xanthium</i> spp.) exceeding 8 seeds
26	12 <i>Xanthium Strumarium</i>
26	12 <i>Xanthium Strumarium</i>
30	12 <i>Xanthium Strumarium</i>
30	12 <i>Xanthium Strumarium</i>
31	96 <i>Xanthium Strumarium</i>
31	12 <i>Xanthium Strumarium</i>
32	12 <i>Xanthium Strumarium</i>
33	24 <i>Xanthium Strumarium</i>
34	12 <i>Xanthium Strumarium</i>
34	24 <i>Xanthium Strumarium</i>

TABLE 10: STANDARDS FOR GRADES OF CLASS WHITE MAIZE AND CLASS YELLOW MAIZE							
Deviation		Maximum permissible deviation allowed (m/m)					
		White maize			Yellow maize		
		WM1	WM2	WM3	YM1	YM2	YM3
1	Foreign matter	0.3%	0.5%	0.75%	0.3%	0.5%	0.75%
2	Defective maize kernels, above and below the 6.35 mm round-hole sieve	7%	13%	30%	*	*	*
3	Defective maize kernels that can pass through the 6.35 mm round-hole sieve	*	*	*	4%	10%	30%
4	Defective maize kernels that can not pass through the 6.35 mm round-hole sieve	*	*	*	9%	20%	30%
5	Other colour maize kernels	3%	6%	10%	2%	5%	5%
6	Deviations referred to in items 1, 2, 3, 4 and 5 collectively: Provided that the deviations are individually within the specified limits	8%	16%	30%	9%	20%	30%
7	Pinked maize kernels	12%	12%	12%	*	*	*

A consignment of maize shall be classified as Class Other Maize if the consignment does not comply with the standards for Class White Maize or Class Yellow Maize.

\* Not specified

Regulations relating to the Grading, Packing and Marking of Maize intended for sale in the Republic of South Africa as published in the Government Gazette No. 32190, Government Notice No. R. 473 of 8 May 2009, as well as Industry-wide Dispensation REF NO: 20/4/14/1 of 15 April 2010.

TABLE 11: GRADES AND GRADE REQUIREMENTS FOR CLASS WHITE AND YELLOW MAIZE ACCORDING TO USA GRADING REGULATIONS					
		Maximum limits of -			
		Damaged kernels		Broken corn and foreign material (percent)	
Grades	Minimum test weight per bushel (pounds)		Heat damaged kernels (percent)		Total (percent)
U.S. No. 1	56.0	72.1 kg/hl	0.1	3.0	2.0
U.S. No. 2	54.0	69.5 kg/hl	0.2	5.0	3.0
U.S. No. 3	52.0	66.9 kg/hl	0.5	7.0	4.0
U.S. No. 4	49.0	63.1 kg/hl	1.0	10.0	5.0
U.S. No. 5	46.0	59.2 kg/hl	3.0	15.0	7.0
U.S. Sample Grade	< 46.0	< 59.2 kg/hl	>3.0	>15.0	>7.0
U.S. Sample grade is corn that:					
a) Does not meet the requirements for the grades U.S. Nos. 1, 2, 3, 4 or 5; or					
b) Contains stones which have an aggregate weight in excess of 0.1 percent of the sample weight, 2 or more pieces of glass, 3 or more crotalaria seeds ( <i>Crotalaria spp.</i> ), 2 or more castor beans ( <i>Ricinus communis L.</i> ), 4 or more particles of an unknown foreign substance(s) or a commonly recognized harmful or toxic substance(s), 8 or more cockleburrs ( <i>Xanthium spp.</i> ) or similar seeds singly or in combination, or animal filth in excess of 0.20 percent in 1,000 grams; or					
c) Has a musty, sour, or commercially objectionable foreign odor; or					
d) Is heating or otherwise of distinctly low quality.					
Mixed corn class	When % other colour in yellow maize samples > 5% and white maize samples > 2%				

Source: Official United States Standard of Grain (excluding metric conversions).

TABLE 12: PHYSICAL QUALITY CHARACTERISTICS OF WHITE MAIZE ACCORDING TO GRADE (2020/21)																															
Number of samples	Region	Test weight (kg/hi)			100 kernel mass (g)			Kernel size (%)			Breakage susceptibility (%)			Stress cracks (%)			SAGL Milling index 2021			GYA											
		ave.	min.	max.	ave.	min.	max.	Above 10 mm sieve	Above 8 mm sieve	Below 8 mm sieve	< 6.35 mm sieve	< 4.75 mm sieve	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.										
								ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.									
<b>GRADE: WM1</b>																															
8	Region 12	74.2	71.9	75.3	34.2	30.7	36.3	32.5	13.5	41.8	60.0	52.4	76.4	7.5	5.8	10.1	1.4	0.4	3.2	1.3	0.4	3.0	18	7	32	69	63	78	62	61	64
23	Region 13	74.2	72.2	76.8	31.9	26.3	38.1	20.2	0.2	50.7	64.6	46.8	77.7	15.2	2.5	33.0	0.8	0.0	2.5	0.6	0.0	2.2	13	5	31	71	61	79	63	60	65
25	Region 14	75.9	73.9	78.1	32.7	28.7	36.3	26.6	6.3	36.5	63.8	57.5	73.9	9.7	4.0	22.9	0.7	0.0	2.5	0.6	0.0	1.7	11	5	19	69	57	83	62	59	65
6	Region 16	75.5	75.2	75.9	30.9	30.2	32.0	15.8	11.8	24.7	66.8	62.6	71.4	17.4	12.7	25.1	1.0	0.2	2.7	0.7	0.0	2.0	11	3	22	66	62	71	61	60	63
15	Region 17	75.3	73.8	76.8	32.7	30.1	34.6	25.9	2.8	42.4	62.6	52.8	72.9	11.6	4.8	24.3	0.5	0.1	1.5	0.4	0.0	1.4	10	5	18	68	51	78	62	58	64
24	Region 18	74.9	72.4	77.3	31.9	28.7	35.8	22.6	3.2	47.1	66.0	48.9	77.7	11.4	4.0	29.8	0.9	0.0	2.9	0.6	0.0	1.8	14	3	33	72	64	86	63	61	66
17	Region 19	74.2	68.5	77.9	31.0	25.8	36.2	13.6	4.2	27.9	70.0	63.4	83.1	16.4	5.3	26.3	0.9	0.0	2.9	0.7	0.0	2.3	13	5	28	71	57	87	63	59	67
15	Region 20	75.5	74.2	77.1	31.2	28.5	35.6	15.1	4.7	32.2	67.4	57.5	76.8	17.5	8.5	28.9	0.6	0.0	1.1	0.4	0.0	0.9	9	2	17	67	48	75	62	57	64
27	Region 21	76.1	74.0	78.1	31.7	26.8	38.7	21.1	3.3	47.8	64.8	47.3	73.0	14.1	4.9	29.8	0.6	0.0	3.0	0.4	0.0	1.6	10	3	28	72	63	82	63	61	65
9	Region 22	76.2	73.8	78.3	34.5	29.8	37.3	27.1	11.7	36.0	63.8	57.6	71.9	9.1	2.4	20.5	1.5	0.4	2.7	1.1	0.3	1.6	18	8	29	69	62	74	62	60	63
31	Region 23	77.0	75.1	78.4	34.7	31.1	40.6	28.9	17.2	48.0	61.2	48.6	69.2	9.9	3.1	17.9	0.5	0.0	1.7	0.4	0.0	1.3	8	2	20	69	57	79	62	59	64
10	Region 24	76.4	73.5	78.1	33.6	29.4	38.2	26.2	0.9	40.0	61.5	53.0	68.4	12.3	6.3	41.8	0.4	0.0	1.1	0.3	0.0	0.8	10	1	18	69	37	80	62	54	65
2	Region 25	74.7	72.7	76.7	32.2	31.0	33.4	15.4	4.0	26.7	67.3	65.6	69.0	17.4	7.7	27.0	0.7	0.4	0.9	0.5	0.3	0.7	17	14	19	62	58	66	60	60	61
5	Region 26	76.0	73.6	77.4	32.3	30.9	35.2	16.1	11.0	22.0	71.6	68.3	74.1	12.4	5.8	20.7	1.3	0.2	3.0	0.9	0.0	2.4	17	4	41	68	58	78	62	59	64
3	Region 27	75.8	74.8	77.0	30.2	29.6	31.0	11.7	0.2	21.6	67.5	61.9	70.6	20.8	7.8	37.9	1.3	1.1	1.5	1.0	0.8	1.3	18	14	23	63	48	78	61	57	64
12	Region 28	77.8	75.9	81.1	34.7	29.1	40.6	21.8	6.1	41.6	66.8	51.5	76.3	11.5	2.5	19.4	0.4	0.0	1.3	0.3	0.0	1.1	12	3	26	81	65	111	65	61	72
26	Region 29	76.2	73.9	79.3	34.1	30.2	39.8	23.3	5.1	48.3	65.7	46.8	78.3	11.0	3.1	23.9	0.6	0.0	1.5	0.5	0.0	1.2	10	3	21	76	64	90	64	61	67
23	Region 30	77.0	74.0	78.9	35.4	31.0	39.8	21.7	7.2	38.4	67.1	55.5	76.5	11.2	5.5	20.5	0.6	0.0	1.5	0.4	0.0	1.1	11	2	26	78	70	88	64	62	67
10	Region 31	76.4	73.5	78.6	31.7	24.9	35.6	10.5	1.9	17.6	68.1	63.7	73.4	21.4	9.9	29.8	0.6	0.2	1.3	0.4	0.0	0.9	9	1	19	77	67	93	64	62	68
13	Region 32	77.0	73.6	79.0	34.2	27.3	39.6	16.1	3.4	28.9	67.8	59.5	73.6	16.0	4.5	31.6	0.6	0.0	1.7	0.5	0.0	1.2	9	4	16	77	58	90	64	59	67
40	Region 33	77.2	73.8	79.4	34.7	26.9	40.4	17.9	3.0	39.0	68.5	52.2	77.8	13.6	1.7	33.6	0.4	0.0	2.0	0.3	0.0	1.8	9	3	23	76	64	90	64	61	67
21	Region 34	76.2	68.8	78.9	32.5	25.6	39.5	16.3	2.5	46.2	62.5	41.8	75.6	21.2	1.6	55.7	0.9	0.0	3.9	0.6	0.0	2.6	13	3	36	76	57	86	64	59	66
6	Region 35	78.1	77.1	78.9	34.3	30.0	39.2	18.4	6.0	23.8	69.6	61.7	82.1	12.0	3.8	14.5	0.6	0.3	1.1	0.5	0.2	0.9	13	5	30	83	75	93	66	64	68
17	Region 36	77.3	75.0	78.9	33.5	27.1	38.4	17.5	5.4	30.9	67.3	55.9	72.6	15.2	4.7	37.3	0.6	0.0	1.5	0.5	0.0	1.4	10	3	19	82	72	89	65	63	67
<b>388</b>	<b>Ave. WM1</b>	<b>76.1</b>			<b>33.2</b>			<b>21.0</b>			<b>65.6</b>			<b>13.4</b>			<b>0.7</b>			<b>0.5</b>			<b>11</b>			<b>73</b>			<b>63</b>		
	<b>Min. WM1</b>	<b>68.5</b>			<b>24.9</b>			<b>0.2</b>			<b>41.8</b>			<b>1.6</b>			<b>0.0</b>			<b>0.0</b>			<b>1</b>			<b>37</b>			<b>54</b>		
	<b>Max. WM1</b>	<b>81.1</b>			<b>40.6</b>			<b>50.7</b>			<b>83.1</b>			<b>55.7</b>			<b>3.9</b>			<b>3.0</b>			<b>41</b>			<b>111</b>			<b>72</b>		

**TABLE 12: PHYSICAL QUALITY CHARACTERISTICS OF WHITE MAIZE ACCORDING TO GRADE (2020/21)**  
(continue)

Number of samples	Region	Test weight (kg/ha)			100 kernel mass (g)			Kernel size (%)						Breakage susceptibility (%)						Stress cracks (%)			SAGL Milling index 2021			GYA			
		ave.	min.	max.	ave.	min.	max.	Above 10 mm sieve		Above 8 mm sieve		Below 8 mm sieve		< 6.35 mm sieve		< 4.75 mm sieve		ave.	min.	max.	ave.	min.	max.	ave.	min.	max.			
								ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.										min.	max.	
1	Region 11	76.5	76.5	76.5	30.7	30.7	30.7	4.5	4.5	4.5	71.2	71.2	71.2	24.3	24.3	24.3	0.4	0.4	0.4	8	8	8	91	91	91	68	68	68	
3	Region 12	74.6	73.9	75.6	35.4	34.3	36.9	28.1	24.2	33.6	62.9	57.9	65.7	9.0	7.8	10.7	1.6	0.7	2.8	25	18	33	70	64	76	62	61	64	
4	Region 13	74.4	72.5	76.1	31.6	28.5	36.4	20.1	7.6	37.5	68.0	59.6	79.2	11.9	2.9	21.8	0.7	0.0	1.6	13	4	22	70	59	75	62	60	63	
9	Region 14	75.6	73.0	77.4	32.5	30.5	34.6	30.3	2.1	44.6	60.9	48.5	71.2	8.7	4.5	26.7	0.8	0.0	2.9	10	5	23	67	61	74	62	60	63	
1	Region 16	76.6	76.6	76.6	32.0	32.0	32.0	24.3	24.3	24.3	66.0	66.0	66.0	9.7	9.7	9.7	1.1	1.1	1.1	11	11	11	79	79	79	65	65	65	
2	Region 17	74.5	74.1	74.9	34.5	33.6	35.3	34.4	20.5	48.2	57.0	48.4	65.5	8.7	3.4	14.0	1.1	0.5	1.6	16	10	22	69	68	71	62	62	63	
3	Region 18	73.5	72.2	74.4	31.5	28.9	34.2	21.8	16.3	31.1	67.9	61.0	74.1	10.3	7.9	13.4	1.0	0.5	1.6	18	9	26	69	64	75	62	61	64	
5	Region 19	73.1	71.0	74.2	30.3	24.7	34.1	17.8	9.4	37.8	68.1	58.5	74.8	14.1	3.7	28.1	1.8	0.7	3.0	21	15	28	65	52	73	61	58	63	
1	Region 20	75.1	75.1	75.1	35.1	35.1	35.1	43.3	43.3	43.3	46.6	46.6	46.6	10.1	10.1	10.1	0.4	0.4	0.4	5	5	5	57	57	57	59	59	59	
3	Region 21	75.1	73.4	76.0	33.3	31.0	36.7	20.5	16.3	28.5	66.6	63.0	71.4	12.9	8.5	18.3	0.7	0.6	0.9	12	3	18	69	58	87	62	59	66	
2	Region 22	76.1	75.6	76.6	34.8	33.8	35.8	32.5	24.5	40.4	60.7	53.3	68.0	6.9	6.3	7.5	2.0	1.7	2.2	26	25	26	63	63	63	61	61	61	
6	Region 23	76.3	73.6	77.6	36.3	33.0	39.0	33.4	22.8	48.7	60.7	50.0	69.0	5.9	1.3	9.7	0.7	0.0	2.7	13	1	24	69	49	82	62	57	65	
2	Region 24	73.7	70.7	76.7	32.0	31.3	32.7	23.8	13.5	34.0	65.4	55.8	75.0	10.9	10.2	11.5	2.4	0.4	4.3	23	8	38	66	62	70	61	60	62	
2	Region 26	75.6	75.2	75.9	30.2	28.0	32.3	14.0	12.1	15.9	73.1	73.0	73.2	12.9	11.1	14.7	0.5	0.3	0.7	9	3	15	71	69	73	63	62	63	
2	Region 27	75.9	75.7	76.0	31.5	29.6	33.3	21.2	19.7	22.7	69.9	69.2	70.5	9.0	8.1	9.8	1.6	1.4	1.7	20	17	22	69	68	70	62	62	62	
1	Region 29	76.6	76.6	76.6	37.4	37.4	37.4	36.8	36.8	36.8	58.1	58.1	58.1	5.1	5.1	5.1	0.9	0.9	0.9	14	14	14	82	82	82	65	65	65	
7	Region 30	77.1	73.4	80.0	35.9	30.9	40.1	19.0	9.2	28.0	67.4	63.5	72.0	13.6	7.9	20.5	0.7	0.0	1.3	13	10	16	83	76	94	65	64	68	
7	Region 31	77.5	74.9	79.0	35.6	32.9	39.1	15.7	2.1	27.1	68.1	55.9	83.6	16.2	9.9	22.8	0.7	0.1	1.8	13	5	26	78	62	84	64	61	66	
8	Region 32	78.0	76.1	79.9	36.4	32.4	39.2	23.3	8.5	38.3	63.0	57.5	76.3	13.6	4.0	28.5	0.4	0.1	1.1	9	3	15	81	72	88	65	63	67	
10	Region 33	76.0	73.3	79.6	37.7	30.5	42.6	28.2	7.9	42.3	63.0	50.8	75.9	8.8	3.4	16.9	1.2	0.0	2.7	21	12	39	78	64	87	64	61	66	
4	Region 34	76.7	74.9	78.6	31.4	29.3	34.7	12.3	10.2	14.3	59.0	45.8	78.3	28.8	9.2	42.2	1.1	0.0	3.5	12	2	32	79	70	86	65	62	66	
1	Region 35	76.8	76.8	76.8	32.0	32.0	32.0	13.6	13.6	13.6	66.0	66.0	66.0	20.4	20.4	20.4	1.2	1.2	1.2	18	18	18	80	80	80	65	65	65	
6	Region 36	77.6	76.9	78.4	33.2	30.2	34.7	14.6	6.3	24.5	65.5	57.9	71.0	19.9	13.4	31.8	0.9	0.2	1.7	10	5	16	76	69	81	64	62	65	
<b>90</b>	<b>Ave. WM2</b>	<b>76.0</b>	<b>70.7</b>	<b>80.0</b>	<b>34.2</b>	<b>24.7</b>	<b>42.6</b>	<b>23.1</b>	<b>2.1</b>	<b>48.7</b>	<b>64.3</b>	<b>45.8</b>	<b>83.6</b>	<b>12.6</b>	<b>1.3</b>	<b>42.2</b>	<b>1.0</b>	<b>0.0</b>	<b>4.3</b>	<b>14</b>	<b>1</b>	<b>39</b>	<b>74</b>	<b>49</b>	<b>94</b>	<b>63</b>	<b>57</b>	<b>68</b>	
	<b>Min. WM2</b>																												
	<b>Max. WM2</b>																												

**TABLE 12: PHYSICAL QUALITY CHARACTERISTICS OF WHITE MAIZE ACCORDING TO GRADE (2020/21)**  
(continue)

Number of samples	Region	Test weight (kg/ha)		100 kernel mass (g)		Kernel size (%)			Breakage susceptibility (%)			Stress cracks (%)			SAGL Milling index 2021			GYA														
		ave.	min.	max.	Above 10 mm sieve		Above 8 mm sieve		Below 8 mm sieve		< 6.35 mm sieve		< 4.75 mm sieve		ave.	min.	max.	ave.	min.	max.	ave.	min.	max.									
					ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.										min.	max.							
<b>GRADE: WM3</b>																																
1	Region 13	71.5	71.5	71.5	28.4	28.4	28.4	28.4	4.5	4.5	4.5	71.3	71.3	71.3	24.2	24.2	24.2	1.5	1.5	1.5	1.0	1.0	1.0	21	21	21	62	62	62	60	60	60
3	Region 14	75.6	75.1	75.9	34.0	32.2	35.7	34.8	20.3	49.1	58.3	46.6	67.4	6.9	4.1	12.3	0.3	0.0	0.6	0.3	0.0	0.5	10	6	12	67	65	71	62	61	63	
1	Region 16	76.9	76.9	76.9	37.5	37.5	37.5	25.4	25.4	25.4	66.3	66.3	66.3	8.3	8.3	8.3	1.1	1.1	1.1	1.0	1.0	1.0	10	10	10	78	78	78	64	64	64	
8	Region 17	73.9	69.2	77.6	32.7	28.2	36.9	26.5	20.5	43.4	65.3	51.3	72.8	8.3	5.3	12.6	1.9	0.2	3.9	1.5	0.2	3.3	19	11	28	71	62	86	63	60	66	
1	Region 18	76.4	76.4	76.4	35.0	35.0	35.0	39.5	39.5	39.5	57.9	57.9	57.9	2.6	2.6	2.6	1.0	1.0	1.0	0.4	0.4	0.4	22	22	22	84	84	84	66	66	66	
5	Region 19	73.5	71.4	75.4	29.7	27.0	33.5	18.7	4.3	28.9	66.2	55.8	73.4	15.0	6.3	23.5	1.3	0.4	2.0	0.9	0.3	1.4	23	13	29	67	49	75	62	57	63	
1	Region 20	75.6	75.6	75.6	30.5	30.5	30.5	22.3	22.3	22.3	59.1	59.1	59.1	18.6	18.6	18.6	0.1	0.1	0.1	0.0	0.0	0.0	8	8	8	70	70	70	62	62	62	
3	Region 21	73.8	72.7	75.5	29.5	28.6	30.1	18.5	16.5	19.8	68.2	66.4	70.5	13.3	10.3	17.1	0.5	0.0	0.9	0.5	0.0	0.8	5	3	6	69	66	74	62	61	63	
5	Region 23	76.5	75.9	77.5	37.6	35.6	40.2	36.5	27.1	49.5	57.0	47.8	62.3	6.6	2.5	11.2	0.9	0.4	2.0	0.6	0.3	1.4	10	4	22	68	64	72	62	61	63	
1	Region 26	73.9	73.9	73.9	30.5	30.5	30.5	11.4	11.4	11.4	61.5	61.5	61.5	27.1	27.1	27.1	1.7	1.7	1.7	1.0	1.0	1.0	20	20	20	69	69	69	62	62	62	
2	Region 30	76.8	75.4	78.1	30.5	25.5	35.4	7.4	3.0	11.8	66.3	59.0	73.5	26.4	14.7	38.0	0.9	0.3	1.5	0.8	0.3	1.3	16	10	21	75	73	78	64	63	64	
2	Region 31	75.2	74.7	75.7	33.0	31.3	34.7	13.9	11.5	16.3	75.7	71.9	79.4	10.5	9.1	11.8	0.4	0.1	0.7	0.3	0.0	0.7	9	8	10	74	68	81	63	62	65	
1	Region 32	74.9	74.9	74.9	31.8	31.8	31.8	20.5	20.5	20.5	63.3	63.3	63.3	16.2	16.2	16.2	0.9	0.9	0.9	0.7	0.7	0.7	13	13	13	72	72	72	63	63	63	
1	Region 33	75.1	75.1	75.1	31.3	31.3	31.3	13.0	13.0	13.0	69.9	69.9	69.9	17.1	17.1	17.1	0.9	0.9	0.9	0.8	0.8	0.8	14	14	14	76	76	76	64	64	64	
1	Region 34	75.5	75.5	75.5	34.0	34.0	34.0	14.7	14.7	14.7	65.1	65.1	65.1	20.2	20.2	20.2	0.4	0.4	0.4	0.3	0.3	0.3	12	12	12	75	75	75	64	64	64	
<b>36</b>	<b>Ave. WM3</b>	<b>74.8</b>			<b>32.6</b>			<b>23.4</b>	<b>3.0</b>		<b>64.3</b>			<b>12.3</b>			<b>1.1</b>			<b>0.8</b>			<b>15</b>			<b>71</b>			<b>62</b>			
	<b>Min. WM3</b>	<b>69.2</b>			<b>25.5</b>			<b>3.0</b>			<b>46.6</b>			<b>2.5</b>			<b>0.0</b>			<b>0.0</b>			<b>3</b>			<b>49</b>			<b>57</b>			
	<b>Max. WM3</b>	<b>78.1</b>			<b>40.2</b>			<b>49.5</b>			<b>79.4</b>			<b>38.0</b>			<b>3.9</b>			<b>3.3</b>			<b>29</b>			<b>86</b>			<b>66</b>			

**TABLE 12: PHYSICAL QUALITY CHARACTERISTICS OF WHITE MAIZE ACCORDING TO GRADE (2020/21)**  
(continue)

Number of samples	Region	Test weight (kg/hl)		100 kernel mass (g)		Kernel size (%)						Breakage susceptibility (%)						Stress cracks (%)			SAGL Milling index 2021			GYA						
		ave.	min.	max.	ave.	min.	max.	Above 10 mm sieve		Above 8 mm sieve		Below 8 mm sieve		< 6.35 mm sieve		< 4.75 mm sieve		ave.	min.	max.	ave.	min.	max.	ave.	min.	max.				
								ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.										min.	max.		
<b>CLASS: COM</b>																														
1	Region 11	69.9	69.9	69.9	18.9	18.9	18.9	0.6	0.6	0.6	45.7	45.7	53.7	53.7	53.7	2.0	2.0	2.0	1.8	1.8	1.8	11	11	11	74	74	74	63	63	63
3	Region 12	74.6	74.3	74.9	34.6	33.3	35.9	35.6	21.0	44.6	57.5	49.5	67.0	7.0	3.0	12.0	1.7	1.5	1.8	1.4	1.2	22	18	26	74	67	77	63	62	64
1	Region 13	74.1	74.1	74.1	34.2	34.2	34.2	38.5	38.5	38.5	56.8	56.8	4.7	4.7	4.7	0.1	0.1	0.1	0.0	0.0	0.0	3	3	3	80	80	80	65	65	65
3	Region 14	75.3	74.9	75.6	34.1	31.9	37.1	35.5	25.4	55.1	55.4	43.4	64.2	9.1	1.5	16.0	1.2	0.5	2.2	0.7	0.3	11	7	19	70	67	72	62	62	63
4	Region 17	74.7	72.2	75.9	32.1	28.9	35.1	21.7	3.1	39.1	65.2	58.2	71.7	13.1	2.7	25.2	0.8	0.2	1.3	0.6	0.0	13	8	18	64	46	71	61	57	63
1	Region 18	75.0	75.0	75.0	35.3	35.3	35.3	5.0	5.0	5.0	74.5	74.5	20.5	20.5	20.5	0.8	0.8	0.8	0.6	0.6	0.6	14	14	14	83	83	83	66	66	66
2	Region 19	75.6	74.0	77.1	31.0	27.1	34.8	19.4	0.0	38.8	53.2	51.5	54.8	27.5	6.4	48.5	1.1	0.7	1.5	0.8	0.6	17	15	18	73	62	85	63	60	66
3	Region 21	74.8	73.7	76.9	29.4	26.0	33.0	21.8	11.4	31.9	64.3	60.8	69.7	13.9	7.3	26.2	0.4	0.0	1.0	0.2	0.0	5	3	7	75	68	87	63	62	67
2	Region 22	72.1	71.6	72.6	34.1	33.3	34.9	27.0	9.6	44.3	66.4	52.2	80.6	6.7	3.5	9.8	2.4	1.4	3.4	1.8	1.3	20	18	22	65	65	65	61	61	61
5	Region 23	76.3	74.6	77.8	36.1	32.4	37.8	34.8	24.0	43.3	59.1	53.2	67.2	6.0	3.5	8.8	0.9	0.0	1.4	0.7	0.0	12	4	20	70	67	72	62	62	63
1	Region 24	77.2	77.2	77.2	30.9	30.9	30.9	14.4	14.4	14.4	65.4	65.4	65.4	20.2	20.2	20.2	1.8	1.8	1.8	1.2	1.2	15	15	15	68	68	68	62	62	62
1	Region 25	76.0	76.0	76.0	26.7	26.7	26.7	4.3	4.3	4.3	63.4	63.4	63.4	32.3	32.3	32.3	1.3	1.3	1.3	1.1	1.1	23	23	23	48	48	48	57	57	57
1	Region 28	77.5	77.5	77.5	33.1	33.1	33.1	10.3	10.3	10.3	67.0	67.0	67.0	22.7	22.7	22.7	0.8	0.8	0.8	0.5	0.5	6	6	6	67	67	67	62	62	62
2	Region 30	76.7	76.3	77.0	37.1	35.5	38.6	34.1	33.8	34.4	60.1	57.9	62.3	5.8	3.3	8.3	0.5	0.0	0.9	0.4	0.0	14	12	15	73	73	73	63	63	63
1	Region 31	72.0	72.0	72.0	29.7	29.7	29.7	28.3	28.3	28.3	65.5	65.5	65.5	6.2	6.2	6.2	2.8	2.8	2.8	2.2	2.2	40	40	40	60	60	60	60	60	60
4	Region 32	75.9	73.7	78.2	34.3	29.6	37.2	25.0	9.0	33.6	65.5	60.6	74.5	9.5	5.8	16.5	0.6	0.1	0.9	0.5	0.1	8	4	12	76	71	83	64	63	66
3	Region 33	73.3	71.9	74.9	39.3	37.6	40.4	18.0	8.0	27.2	71.5	61.2	78.8	10.4	6.5	13.2	0.3	0.1	0.7	0.2	0.0	21	18	24	88	85	90	67	66	67
4	Region 34	75.4	69.6	78.3	31.2	29.1	33.3	11.8	9.3	14.6	68.9	64.8	74.8	19.4	15.9	21.9	1.1	0.3	2.1	0.7	0.2	11	4	22	77	68	82	64	62	65
4	Region 36	75.6	74.9	76.9	32.0	27.6	35.7	9.4	6.4	16.8	71.1	66.1	76.9	19.6	13.2	27.5	0.8	0.2	1.4	0.5	0.1	17	8	25	74	65	81	63	61	65
<b>46</b>	<b>Ave. COM</b>	<b>75.0</b>	<b>69.6</b>	<b>78.3</b>	<b>33.1</b>	<b>18.9</b>	<b>40.4</b>	<b>22.6</b>	<b>0.0</b>	<b>55.1</b>	<b>63.5</b>	<b>43.4</b>	<b>80.6</b>	<b>13.9</b>	<b>1.5</b>	<b>53.7</b>	<b>1.0</b>	<b>0.0</b>	<b>3.4</b>	<b>0.7</b>	<b>0.0</b>	<b>14</b>	<b>3</b>	<b>40</b>	<b>72</b>	<b>46</b>	<b>90</b>	<b>63</b>	<b>57</b>	<b>67</b>
<b>560</b>	<b>Ave. WM</b>	<b>75.9</b>	<b>68.5</b>	<b>81.1</b>	<b>33.3</b>	<b>18.9</b>	<b>42.6</b>	<b>21.6</b>	<b>0.0</b>	<b>55.1</b>	<b>65.1</b>	<b>41.8</b>	<b>83.6</b>	<b>13.3</b>	<b>1.3</b>	<b>55.7</b>	<b>0.8</b>	<b>0.0</b>	<b>4.3</b>	<b>0.6</b>	<b>0.0</b>	<b>12</b>	<b>1</b>	<b>41</b>	<b>73</b>	<b>37</b>	<b>111</b>	<b>63</b>	<b>54</b>	<b>72</b>
<b>1000</b>	<b>Ave. Maize</b>	<b>76.2</b>	<b>68.5</b>	<b>82.6</b>	<b>32.4</b>	<b>18.9</b>	<b>43.8</b>	<b>15.4</b>	<b>0.0</b>	<b>55.1</b>	<b>65.5</b>	<b>5.6</b>	<b>83.6</b>	<b>19.2</b>	<b>1.3</b>	<b>88.1</b>	<b>0.8</b>	<b>0.0</b>	<b>4.3</b>	<b>0.6</b>	<b>0.0</b>	<b>12</b>	<b>1</b>	<b>49</b>	<b>74</b>	<b>37</b>	<b>111</b>	<b>63</b>	<b>54</b>	<b>72</b>
	<b>Min. Maize</b>																													
	<b>Max. Maize</b>																													

TABLE 13: PHYSICAL QUALITY CHARACTERISTICS OF WHITE MAIZE (2020/21)																															
Number of samples	Region	Test weight (kg/hl)			100 kernel mass (g)			Kernel size (%)						Breakage susceptibility (%)						Stress cracks (%)			SAGL Milling index 2021			GYA					
		ave.	min.	max.	ave.	min.	max.	Above 10 mm sieve		Above 8 mm sieve		Below 8 mm sieve		< 6.35 mm sieve		< 4.75 mm sieve		ave.	min.	max.	ave.	min.	max.	ave.	min.	max.					
								ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.										min.	max.			
<b>WHITE</b>																															
2	Region 11	73.2	69.9	76.5	24.8	18.9	30.7	2.6	0.6	4.5	58.5	45.7	71.2	39.0	24.3	53.7	1.2	0.4	2.0	1.1	0.4	1.8	10	8	11	83	74	91	65	63	68
14	Region 12	74.4	71.9	75.6	34.5	30.7	36.9	32.2	13.5	44.6	60.1	49.5	76.4	7.7	3.0	12.0	1.5	0.4	3.2	1.3	0.4	3.0	20	7	33	70	63	78	62	61	64
29	Region 13	74.1	71.5	76.8	31.8	26.3	38.1	20.3	0.2	50.7	65.0	46.8	79.2	14.7	2.5	33.0	0.8	0.0	2.5	0.6	0.0	2.2	13	3	31	71	59	80	63	60	65
40	Region 14	75.7	73.0	78.1	32.9	28.7	37.1	28.7	2.1	55.1	62.1	43.4	73.9	9.2	1.5	26.7	0.8	0.0	2.9	0.6	0.0	1.8	10	5	23	68	57	83	62	59	65
8	Region 16	75.8	75.2	76.9	31.9	30.2	37.5	18.1	11.8	25.4	66.7	62.6	71.4	15.3	8.3	25.1	1.0	0.2	2.7	0.8	0.0	2.0	11	3	22	69	62	79	62	60	65
29	Region 17	74.7	69.2	77.6	32.7	28.2	36.9	26.0	2.8	48.2	63.3	48.4	72.9	10.7	2.7	25.2	1.0	0.1	3.9	0.8	0.0	3.3	13	5	28	68	46	86	62	57	66
29	Region 18	74.8	72.2	77.3	32.1	28.7	35.8	22.5	3.2	47.1	66.2	48.9	77.7	11.3	2.6	29.8	0.9	0.0	2.9	0.6	0.0	1.8	14	3	33	72	64	86	63	61	66
29	Region 19	74.0	68.5	77.9	30.7	24.7	36.2	15.6	0.0	38.8	67.9	51.5	83.1	16.5	3.7	48.5	1.1	0.0	3.0	0.9	0.0	2.5	16	5	29	69	49	87	62	57	67
17	Region 20	75.5	74.2	77.1	31.4	28.5	35.6	17.2	4.7	43.3	65.7	46.6	76.8	17.1	8.5	28.9	0.5	0.0	1.1	0.4	0.0	0.9	8	2	17	67	48	75	62	57	64
36	Region 21	75.7	72.7	78.1	31.5	26.0	38.7	20.9	3.3	47.8	65.2	47.3	73.0	13.9	4.9	29.8	0.6	0.0	3.0	0.4	0.0	1.6	10	3	28	72	58	87	63	59	67
13	Region 22	75.5	71.6	78.3	34.5	29.8	37.3	27.9	9.6	44.3	63.7	52.2	80.6	8.4	2.4	20.5	1.7	0.4	3.4	1.2	0.3	2.3	19	8	29	67	62	74	62	60	63
47	Region 23	76.8	73.6	78.4	35.3	31.1	40.6	30.9	17.2	49.5	60.5	47.8	69.2	8.6	1.3	17.9	0.6	0.0	2.7	0.5	0.0	1.7	9	1	24	69	49	82	62	57	65
13	Region 24	76.0	70.7	78.1	33.1	29.4	38.2	24.9	0.9	40.0	62.4	53.0	75.0	12.7	6.3	41.8	0.8	0.0	4.3	0.5	0.0	2.1	12	1	38	68	37	80	62	54	65
3	Region 25	75.1	72.7	76.7	30.4	26.7	33.4	11.7	4.0	26.7	66.0	63.4	69.0	22.3	7.7	32.3	0.9	0.4	1.3	0.7	0.3	1.1	19	14	23	57	48	66	59	57	61
8	Region 26	75.7	73.6	77.4	31.5	28.0	35.2	15.0	11.0	22.0	70.7	61.5	74.1	14.4	5.8	27.1	1.2	0.2	3.0	0.8	0.0	2.4	16	3	41	69	58	78	62	59	64
5	Region 27	75.8	74.8	77.0	30.7	29.6	33.3	15.5	0.2	22.7	68.4	61.9	70.6	16.1	7.8	37.9	1.4	1.1	1.7	1.0	0.8	1.3	18	14	23	66	48	78	61	57	64
13	Region 28	77.8	75.9	81.1	34.6	29.1	40.6	20.9	6.1	41.6	66.8	51.5	76.3	12.4	2.5	22.7	0.5	0.0	1.3	0.3	0.0	1.1	11	3	26	80	65	111	65	61	72
27	Region 29	76.2	73.9	79.3	34.3	30.2	39.8	23.8	5.1	48.3	65.4	46.8	78.3	10.8	3.1	23.9	0.6	0.0	1.5	0.5	0.0	1.2	10	3	21	77	64	90	64	61	67
34	Region 30	77.0	73.4	80.0	35.3	25.5	40.1	21.0	3.0	38.4	66.7	55.5	76.5	12.3	3.3	38.0	0.6	0.0	1.5	0.5	0.0	1.3	12	2	26	79	70	94	64	62	68
20	Region 31	76.4	72.0	79.0	33.1	24.9	39.1	13.6	1.9	28.3	68.7	55.9	83.6	17.7	6.2	29.8	0.7	0.1	2.8	0.5	0.0	2.2	12	1	40	76	60	93	64	60	68
26	Region 32	77.0	73.6	79.9	34.8	27.3	39.6	19.9	3.4	38.3	65.8	57.5	76.3	14.3	4.0	31.6	0.6	0.0	1.7	0.4	0.0	1.2	9	3	16	78	58	90	64	59	67
54	Region 33	76.7	71.9	79.6	35.5	26.9	42.6	19.7	3.0	42.3	67.7	50.8	78.8	12.6	1.7	33.6	0.6	0.0	2.7	0.5	0.0	2.6	12	3	39	77	64	90	64	61	67
30	Region 34	76.2	68.8	78.9	32.2	25.6	39.5	15.1	2.5	46.2	63.0	41.8	78.3	21.9	1.6	55.7	0.9	0.0	3.9	0.7	0.0	2.6	13	2	36	76	57	86	64	59	66
7	Region 35	77.9	76.8	78.9	34.0	30.0	39.2	17.7	6.0	23.8	69.1	61.7	82.1	13.2	3.8	20.4	0.7	0.3	1.2	0.6	0.2	0.9	13	5	30	83	75	93	65	64	68
27	Region 36	77.1	74.9	78.9	33.2	27.1	38.4	15.7	5.4	30.9	67.5	55.9	76.9	16.9	4.7	37.3	0.7	0.0	1.7	0.6	0.0	1.6	11	3	25	79	65	89	65	61	67
560	Ave. White	75.9			33.3			21.6			65.1			13.3			0.8			0.6			12			73			63		
	Min. White	68.5			18.9			0.0			41.8			1.3			0.0			0.0			1			37			54		
	Max. White	81.1			42.6			55.1			83.6			55.7			4.3			3.3			41			111			72		



TABLE 14: PHYSICAL QUALITY CHARACTERISTICS OF YELLOW MAIZE ACCORDING TO GRADE (2020/21)																																
Number of samples	Region	Test weight (kg/hl)			100 kernel mass (g)			Kernel size (%)						Breakage susceptibility (%)						Stress cracks (%)			SAGL Milling index 2021			GYA						
		min.		max.	min.		max.	Above 10 mm sieve		Above 8 mm sieve		Below 8 mm sieve		< 6.35 mm sieve		< 4.75 mm sieve		ave.		min.		max.		ave.		min.		max.				
		ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.				
<b>GRADE: YM1</b>																																
24	Region 10	78.9	75.5	82.6	36.0	32.0	43.8	3.4	0.0	10.1	68.1	48.7	81.0	28.6	9.1	48.8	0.4	0.0	1.8	0.3	0.0	1.2	7.1	1	18	74	41	84	63	55	66	
13	Region 11	79.0	76.4	80.1	34.7	29.5	38.0	4.9	1.7	20.7	69.2	63.3	78.4	25.9	16.0	34.1	0.7	0.0	1.5	0.5	0.0	1.1	10.3	3	19	75	65	87	64	61	67	
5	Region 13	77.6	76.8	78.4	28.6	25.8	30.2	7.4	4.1	15.6	71.6	68.3	75.4	21.0	12.6	27.6	0.7	0.1	1.3	0.5	0.0	1.1	9.4	4	14	86	83	88	66	66	67	
6	Region 14	77.4	75.6	79.0	31.0	28.1	34.3	6.9	3.0	13.2	66.9	55.0	78.8	26.1	10.9	41.8	1.2	0.4	2.0	0.9	0.3	1.4	15.6	6	25	84	71	91	66	63	67	
3	Region 17	77.5	75.7	79.8	29.1	27.4	30.7	3.6	2.1	6.1	63.9	59.2	69.9	32.5	24.0	38.3	0.4	0.0	1.2	0.3	0.0	1.0	10.5	5	15	81	73	86	65	63	66	
6	Region 18	76.0	72.1	77.6	29.5	25.5	34.1	7.4	4.7	9.7	70.6	65.0	77.6	22.1	16.3	26.0	1.3	0.4	2.2	1.0	0.3	1.9	16.7	7	32	83	80	87	65	65	66	
7	Region 19	74.3	71.2	77.7	27.0	24.0	29.0	9.8	3.1	15.8	68.1	60.1	74.1	22.1	15.2	34.7	1.0	0.4	2.9	0.8	0.3	2.1	10.5	5	20	84	79	87	66	65	66	
7	Region 20	77.2	76.1	79.1	30.1	27.1	37.1	3.7	1.5	6.8	67.8	60.2	75.5	28.5	22.0	37.3	0.7	0.1	1.5	0.5	0.1	1.3	13.4	4	33	77	69	82	64	62	65	
3	Region 21	76.4	75.1	78.0	30.3	26.8	32.7	7.7	1.9	13.7	67.6	65.0	70.1	24.7	21.3	30.4	1.0	0.2	2.4	0.7	0.1	1.7	13.2	2	20	85	85	86	66	66	66	
5	Region 22	77.8	77.1	78.3	28.5	27.6	29.6	6.0	4.8	7.4	67.7	63.8	71.4	26.3	22.7	31.4	0.7	0.4	1.2	0.5	0.3	1.2	7.3	3	11	84	79	94	66	65	68	
4	Region 23	76.6	75.4	77.8	28.4	27.1	30.0	6.6	2.5	11.6	60.7	46.2	68.8	32.7	19.6	51.3	0.9	0.4	1.4	0.7	0.4	1.0	9.4	4	15	78	62	90	64	60	67	
4	Region 24	77.3	75.3	78.3	30.5	26.5	33.7	2.7	1.6	3.7	62.7	60.2	66.8	34.6	30.0	37.0	0.7	0.0	1.1	0.6	0.0	0.9	12.5	5	23	75	60	82	64	60	65	
18	Region 25	75.2	73.7	76.6	29.2	26.3	32.1	6.5	0.8	26.1	64.6	52.4	69.7	28.9	17.6	39.1	0.8	0.0	1.6	0.7	0.0	1.5	15.6	6	33	69	56	88	62	59	67	
11	Region 26	75.9	72.9	77.2	28.6	25.5	34.2	6.6	1.7	26.8	65.1	60.0	70.3	28.3	6.2	35.1	0.7	0.2	2.0	0.6	0.1	1.7	13.4	4	25	79	62	100	65	60	70	
4	Region 27	74.5	71.4	76.8	25.6	22.8	27.9	3.2	0.8	5.0	61.3	57.1	63.8	35.5	32.3	42.1	0.9	0.7	1.2	0.8	0.6	1.0	16.8	8	25	77	75	79	64	64	65	
28	Region 28	75.9	74.3	78.8	30.2	23.9	38.5	6.4	0.1	32.0	65.4	55.5	76.5	28.2	8.2	44.4	0.8	0.0	2.3	0.6	0.0	1.5	13.3	3	28	73	55	86	63	59	66	
26	Region 29	77.0	74.2	79.4	31.4	27.9	35.2	7.7	2.2	16.8	69.4	58.8	78.7	22.9	10.5	38.2	0.7	0.0	1.6	0.5	0.0	1.3	14.4	4	49	80	59	89	65	60	67	
39	Region 30	76.3	71.6	78.5	33.5	29.2	39.5	11.6	2.5	32.5	68.4	53.3	80.1	20.0	6.1	39.3	0.5	0.0	1.5	0.3	0.0	1.4	10.3	3	32	77	60	95	64	60	68	
39	Region 31	77.7	74.3	80.2	32.3	28.0	37.1	9.0	0.4	23.0	67.2	58.9	81.8	23.8	9.8	37.0	0.5	0.0	1.6	0.4	0.0	1.4	10.2	2	23	78	66	93	64	61	68	
32	Region 32	77.3	73.5	80.3	33.1	24.1	37.2	10.7	0.6	20.9	67.1	49.8	76.1	22.2	9.7	49.4	0.7	0.0	1.8	0.5	0.0	1.4	11.4	4	21	80	72	89	65	63	67	
20	Region 33	76.5	75.0	79.7	32.1	26.1	37.1	7.5	1.4	14.0	65.9	47.6	73.9	26.7	13.6	50.5	0.8	0.0	2.0	0.6	0.0	1.7	12.4	4	23	69	59	81	62	60	65	
16	Region 34	77.0	74.5	79.5	30.3	22.7	37.4	9.3	1.0	31.8	62.2	42.3	74.3	28.5	13.0	53.8	0.6	0.0	1.8	0.4	0.0	1.2	11.5	5	21	82	75	93	65	64	68	
6	Region 35	77.3	73.7	79.9	29.4	22.4	34.1	5.2	1.2	13.2	62.9	51.5	72.1	31.9	20.7	46.8	0.9	0.2	1.8	0.8	0.2	1.5	12.8	8	15	79	50	89	65	58	67	
17	Region 36	76.3	75.0	79.5	31.5	28.5	34.9	8.7	0.0	28.6	62.4	44.2	71.9	28.9	13.5	55.2	0.7	0.1	2.0	0.5	0.0	1.4	11.4	4	29	73	49	87	63	57	66	
343	Ave. YM1	76.9	71.2	82.6	31.6	22.4	43.8	7.8	0.0	32.5	66.6	42.3	81.8	25.6	6.1	55.2	0.7	0.0	2.9	0.5	0.0	2.1	11.1	1	49	77	41	100	64	55	70	
	Min. YM1																															
	Max. YM1																															

**TABLE 14: PHYSICAL QUALITY CHARACTERISTICS OF YELLOW MAIZE ACCORDING TO GRADE (2020/21)**  
(continue)

Number of samples	Region	Test weight (kg/hl)			100 kernel mass (g)			Kernel size (%)						Breakage susceptibility (%)						Stress cracks (%)			SAGL Milling index 2021			GYA					
		ave.	min.	max.	ave.	min.	max.	Above 10 mm sieve		Above 8 mm sieve		Below 8 mm sieve		< 6.35 mm sieve		< 4.75 mm sieve		ave.	min.	max.	ave.	min.	max.	ave.	min.	max.					
								ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.										min.	max.			
<b>GRADE: YM2</b>																															
3	Region 10	76.6	73.8	78.8	35.5	33.7	37.1	2.4	1.3	4.6	70.3	59.6	77.6	27.3	21.1	39.1	0.8	0.1	1.5	0.7	0.1	1.4	10	4	20	64	49	75	61	57	64
2	Region 11	74.7	71.6	77.8	29.5	23.6	35.3	5.0	0.6	9.4	54.1	39.3	68.8	41.0	21.8	60.1	1.4	0.6	2.2	1.2	0.5	1.8	17	15	18	63	51	76	61	58	64
1	Region 12	75.1	75.1	75.1	36.4	36.4	36.4	5.6	5.6	5.6	66.6	66.6	66.6	27.8	27.8	27.8	0.9	0.9	0.9	0.8	0.8	0.8	10	10	10	77	77	77	64	64	64
1	Region 13	74.5	74.5	74.5	26.9	26.9	26.9	4.3	4.3	4.3	65.7	65.7	65.7	30.0	30.0	30.0	0.8	0.8	0.8	0.5	0.5	0.5	8	8	8	82	82	82	65	65	65
1	Region 14	74.9	74.9	74.9	27.1	27.1	27.1	3.0	3.0	3.0	60.5	60.5	60.5	36.5	36.5	36.5	1.1	1.1	1.1	0.9	0.9	0.9	10	10	10	80	80	80	65	65	65
1	Region 16	76.7	76.7	76.7	28.7	28.7	28.7	3.3	3.3	3.3	64.8	64.8	64.8	31.9	31.9	31.9	0.7	0.7	0.7	0.6	0.6	0.6	6	6	6	71	71	71	63	63	63
1	Region 18	75.2	75.2	75.2	26.4	26.4	26.4	5.9	5.9	5.9	64.3	64.3	64.3	29.8	29.8	29.8	2.1	2.1	2.1	1.8	1.8	1.8	26	26	26	81	81	81	65	65	65
2	Region 19	76.9	74.9	78.9	27.0	22.7	31.3	6.9	6.3	7.4	41.3	5.6	77.0	51.9	15.6	88.1	0.5	0.2	0.8	0.5	0.2	0.7	12	5	18	86	86	86	66	66	66
1	Region 21	73.9	73.9	73.9	24.2	24.2	24.2	0.8	0.8	0.8	56.1	56.1	56.1	43.1	43.1	43.1	1.1	1.1	1.1	0.9	0.9	0.9	18	18	18	79	79	79	65	65	65
1	Region 22	75.8	75.8	75.8	29.8	29.8	29.8	2.3	2.3	2.3	58.8	58.8	58.8	38.9	38.9	38.9	2.2	2.2	2.2	1.9	1.9	1.9	21	21	21	76	76	76	64	64	64
1	Region 24	74.0	74.0	74.0	30.3	30.3	30.3	8.1	8.1	8.1	69.8	69.8	69.8	22.1	22.1	22.1	1.4	1.4	1.4	1.2	1.2	1.2	14	14	14	56	56	56	59	59	59
5	Region 25	74.4	73.7	75.4	28.5	26.4	29.8	4.0	1.4	6.7	61.0	47.6	69.0	35.0	24.3	51.0	0.8	0.3	1.2	0.5	0.1	0.9	13	6	26	66	59	72	61	60	63
3	Region 26	74.0	72.7	75.3	27.8	26.1	30.0	6.1	2.3	9.9	64.1	62.6	65.2	29.8	24.9	35.1	2.1	0.9	3.5	1.5	0.7	2.1	20	15	30	69	59	76	62	60	64
4	Region 27	74.6	73.6	75.3	25.2	23.1	26.3	3.2	2.1	5.0	59.5	53.4	64.9	37.3	30.1	44.5	1.0	0.7	1.4	0.9	0.6	1.3	12	7	18	77	76	78	64	64	64
5	Region 28	74.3	72.0	75.2	27.2	23.4	30.6	5.0	1.0	9.6	59.0	49.0	66.2	36.0	24.2	50.0	0.4	0.0	1.0	0.2	0.0	0.4	11	8	15	72	66	79	63	61	64
2	Region 30	76.1	74.9	77.4	34.6	34.1	35.0	10.5	5.6	15.4	72.2	69.9	74.5	17.3	14.7	19.9	0.4	0.0	0.8	0.3	0.0	0.6	15	12	18	77	74	79	64	63	65
1	Region 31	75.3	75.3	75.3	30.5	30.5	30.5	5.9	5.9	5.9	59.2	59.2	59.2	34.9	34.9	34.9	0.2	0.2	0.2	0.2	0.2	0.2	8	8	8	61	61	61	60	60	60
3	Region 32	77.4	76.8	78.0	33.9	31.6	35.9	12.2	5.0	21.8	57.3	39.5	68.1	30.5	10.1	55.5	0.3	0.0	0.7	0.2	0.0	0.6	11	8	13	77	70	86	64	62	66
2	Region 33	76.0	75.4	76.7	31.5	30.7	32.2	3.3	1.0	5.6	65.8	59.0	72.5	31.0	21.9	40.0	0.8	0.7	0.8	0.5	0.5	0.5	10	9	11	73	70	77	63	62	64
2	Region 34	74.8	74.1	75.4	26.9	25.2	28.6	5.2	3.1	7.2	61.1	58.1	64.1	33.8	28.7	38.8	0.6	0.2	1.1	0.5	0.1	0.9	7	5	8	77	72	82	64	63	65
1	Region 36	75.4	75.4	75.4	35.3	35.3	35.3	1.0	1.0	1.0	59.3	59.3	59.3	39.7	39.7	39.7	1.2	1.2	1.2	1.0	1.0	1.0	13	13	13	81	81	81	65	65	65
<b>43</b>	<b>Ave. YM2</b>	<b>75.2</b>			<b>29.4</b>			<b>5.2</b>			<b>61.1</b>			<b>33.8</b>			<b>0.9</b>			<b>0.7</b>			<b>13</b>			<b>73</b>			<b>63</b>		
	<b>Min. YM2</b>	<b>71.6</b>			<b>22.7</b>			<b>0.6</b>			<b>5.6</b>			<b>10.1</b>			<b>0.0</b>			<b>0.0</b>			<b>4</b>			<b>49</b>			<b>57</b>		
	<b>Max. YM2</b>	<b>78.9</b>			<b>37.1</b>			<b>21.8</b>			<b>77.6</b>			<b>88.1</b>			<b>3.5</b>			<b>2.1</b>			<b>30</b>			<b>86</b>			<b>66</b>		

**TABLE 14: PHYSICAL QUALITY CHARACTERISTICS OF YELLOW MAIZE ACCORDING TO GRADE (2020/21)**  
(continue)

Number of samples	Region	Test weight (kg/hl)			100 kernel mass (g)			Kernel size (%)			Breakage susceptibility (%)			Stress cracks (%)			SAGL Milling index 2021			GYA											
		ave.	min.	max.	ave.	min.	max.	Above 10 mm sieve			Above 8 mm sieve			Below 8 mm sieve			ave.	min.	max.	ave.	min.	max.	ave.	min.	max.						
								ave.	min.	max.	ave.	min.	max.	ave.	min.	max.										ave.	min.	max.			
<b>GRADE: YM3</b>																															
1	Region 10	78.9	78.9	78.9	34.9	34.9	34.9	0.9	0.9	0.9	60.4	60.4	60.4	38.7	38.7	38.7	1.4	1.4	1.4	1.1	1.1	1.1	16	16	16	61	61	61	60	60	60
1	Region 21	77.2	77.2	77.2	21.4	21.4	21.4	2.5	2.5	2.5	59.9	59.9	59.9	37.6	37.6	37.6	0.3	0.3	0.3	0.2	0.2	0.2	5	5	5	90	90	90	67	67	67
1	Region 24	71.8	71.8	71.8	28.7	28.7	28.7	2.7	2.7	2.7	67.4	67.4	67.4	29.9	29.9	29.9	2.9	2.9	2.9	1.7	1.7	1.7	29	29	29	58	58	58	59	59	59
1	Region 29	75.1	75.1	75.1	28.1	28.1	28.1	8.7	8.7	8.7	66.8	66.8	66.8	24.5	24.5	24.5	0.7	0.7	0.7	0.7	0.7	0.7	10	10	10	77	77	77	64	64	64
1	Region 30	77.6	77.6	77.6	26.3	26.3	26.3	5.6	5.6	5.6	46.9	46.9	46.9	47.5	47.5	47.5	2.2	2.2	2.2	1.1	1.1	1.1	22	22	22	84	84	84	66	66	66
2	Region 33	72.9	71.3	74.4	34.2	30.1	38.2	6.4	5.8	7.0	66.3	65.1	67.4	27.4	26.8	27.9	0.8	0.7	0.8	0.7	0.6	0.7	11	9	13	68	66	70	62	61	62
1	Region 34	78.2	78.2	78.2	29.7	29.7	29.7	3.7	3.7	3.7	69.2	69.2	69.2	27.1	27.1	27.1	1.0	1.0	1.0	0.6	0.6	0.6	13	13	13	94	94	94	68	68	68
8	Ave. YM3	75.6			29.7			4.6			62.9		32.5			1.3			0.8			15			75			64			
	Min. YM3	71.3			21.4			0.9			46.9		24.5			0.3			0.2			5			58			59			
	Max. YM3	78.9			38.2			8.7			69.2		47.5			2.9			1.7			29			94			68			
<b>CLASS: COM</b>																															
1	12	76.6	76.6	76.6	28.8	28.8	28.8	10.5	10.5	10.5	66.8	66.8	66.8	22.7	22.7	22.7	0.2	0.2	0.2	0.1	0.1	0.1	3	3	3	84	84	84	66	66	66
1	14	75.9	75.9	75.9	26.0	26.0	26.0	6.5	6.5	6.5	68.3	68.3	68.3	25.2	25.2	25.2	1.0	1.0	1.0	0.8	0.8	0.8	15	15	15	80	80	80	65	65	65
1	17	76.8	76.8	76.8	25.9	25.9	25.9	7.1	7.1	7.1	61.7	61.7	61.7	31.2	31.2	31.2	0.2	0.2	0.2	0.1	0.1	0.1	5	5	5	86	86	86	66	66	66
2	18	75.1	74.9	75.2	30.1	27.1	33.1	6.9	6.2	7.6	70.0	68.7	71.2	23.1	22.6	23.7	1.8	0.7	2.8	1.5	0.6	2.3	15	5	25	83	81	85	65	65	66
2	21	75.3	74.5	76.1	27.3	25.1	29.5	6.4	2.4	10.3	64.8	55.6	73.9	28.9	15.8	42.0	1.3	1.0	1.5	0.9	0.8	1.0	15	12	17	81	80	81	65	65	65
1	22	75.9	75.9	75.9	30.0	30.0	30.0	2.9	2.9	2.9	58.1	58.1	58.1	39.0	39.0	39.0	1.5	1.5	1.5	1.5	1.5	1.5	15	15	15	67	67	67	62	62	62
1	25	70.7	70.7	70.7	29.3	29.3	29.3	5.0	5.0	5.0	56.2	56.2	56.2	38.8	38.8	38.8	2.4	2.4	2.4	1.9	1.9	1.9	25	25	25	65	65	65	61	61	61
6	26	74.3	72.2	76.1	28.9	26.3	32.1	7.6	4.4	17.2	66.3	62.4	71.2	26.1	13.4	33.0	1.4	1.1	2.4	1.1	0.9	1.8	19	13	33	70	60	80	62	60	65
2	27	73.8	72.1	75.4	28.3	24.3	32.2	5.2	1.3	9.0	64.6	53.9	75.2	30.3	15.8	44.8	1.4	1.2	1.6	1.3	1.1	1.4	21	14	28	76	75	76	64	64	64
1	28	73.4	73.4	73.4	28.3	28.3	28.3	5.4	5.4	5.4	60.6	60.6	60.6	34.0	34.0	34.0	2.2	2.2	2.2	1.4	1.4	1.4	25	25	25	62	62	62	60	60	60
4	29	76.2	75.0	77.7	33.4	28.9	38.0	7.4	0.6	11.3	73.1	70.2	77.0	19.5	11.7	24.9	1.0	0.4	1.4	0.5	0.4	0.7	20	9	37	78	73	83	64	63	65
3	30	76.3	75.2	76.9	34.6	31.3	37.1	7.3	2.1	10.7	69.3	67.4	71.7	23.3	21.9	26.2	0.7	0.2	1.1	0.5	0.1	0.9	9	4	16	74	72	77	63	63	64
4	31	76.2	74.9	77.1	30.4	26.7	33.6	7.1	4.6	11.3	67.0	60.1	72.3	26.0	19.7	33.9	0.7	0.2	1.5	0.4	0.2	0.7	10	4	22	77	74	79	64	63	64
5	32	75.6	74.1	77.4	30.2	20.9	34.5	7.4	3.9	11.5	65.8	60.1	70.1	26.8	21.7	31.2	0.9	0.2	1.7	0.8	0.2	1.3	9	5	16	75	70	80	64	62	65
5	33	76.0	72.7	79.4	32.8	29.8	36.9	9.3	1.5	23.3	67.3	65.2	68.6	23.4	9.1	30.4	0.5	0.2	0.9	0.4	0.1	0.7	10	6	16	72	66	80	63	61	65




**TABLE 15: PHYSICAL QUALITY CHARACTERISTICS OF YELLOW MAIZE (2020/21)**

Number of samples	Region	Test weight (kg/ht)			100 kernel mass (g)			Above 10 mm sieve			Kernel size (%)			Below 8 mm sieve			Breakage susceptibility (%)			Stress cracks (%)			SAGL Milling index 2021			GYA						
		ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.				
																													ave.	min.	max.	ave.
<b>YELLOW</b>																																
28	Region 10	78.6	73.8	82.6	35.9	32.0	43.8	3.2	0.0	10.1	68.1	48.7	81.0	28.8	9.1	48.8	0.5	0.0	1.8	0.4	0.0	1.4	8	1	20	72	41	84	63	55	66	
15	Region 11	78.4	71.6	80.1	34.0	23.6	38.0	4.9	0.6	20.7	67.2	39.3	78.4	27.9	16.0	60.1	0.8	0.0	2.2	0.6	0.0	1.8	11	3	19	73	51	87	63	58	67	
2	Region 12	75.9	75.1	76.6	32.6	28.8	36.4	8.1	5.6	10.5	66.7	66.6	66.8	25.3	22.7	27.8	0.6	0.2	0.9	0.5	0.1	0.8	7	3	10	80	77	84	65	64	66	
6	Region 13	77.1	74.5	78.4	28.4	25.8	30.2	6.9	4.1	15.6	70.6	65.7	75.4	22.5	12.6	30.0	0.7	0.1	1.3	0.5	0.0	1.1	9	4	14	85	82	88	66	65	67	
8	Region 14	76.9	74.9	79.0	29.9	26.0	34.3	6.4	3.0	13.2	66.3	55.0	78.8	27.3	10.9	41.8	1.2	0.4	2.0	0.9	0.3	1.4	14	6	25	83	71	91	66	63	67	
1	Region 16	76.7	76.7	76.7	28.7	28.7	28.7	3.3	3.3	3.3	64.8	64.8	64.8	31.9	31.9	31.9	0.7	0.7	0.7	0.6	0.6	0.6	6	6	6	71	71	71	63	63	63	
4	Region 17	77.3	75.7	79.8	28.3	25.9	30.7	4.5	2.1	7.1	63.4	59.2	69.9	32.2	24.0	38.3	0.4	0.0	1.2	0.3	0.0	1.0	9	5	15	82	73	86	65	63	66	
9	Region 18	75.7	72.1	77.6	29.3	25.5	34.1	7.1	4.7	9.7	69.7	64.3	77.6	23.2	16.3	29.8	1.5	0.4	2.8	1.2	0.3	2.3	17	5	32	83	80	87	65	65	66	
9	Region 19	74.9	71.2	78.9	27.0	22.7	31.3	9.1	3.1	15.8	62.2	5.6	77.0	28.7	15.2	88.1	0.9	0.2	2.9	0.7	0.2	2.1	10	5	20	85	79	87	66	65	66	
7	Region 20	77.2	76.1	79.1	30.1	27.1	37.1	3.7	1.5	6.8	67.8	60.2	75.5	28.5	22.0	37.3	0.7	0.1	1.5	0.5	0.1	1.3	13	4	33	77	69	82	64	62	65	
7	Region 21	75.8	73.9	78.0	27.3	21.4	32.7	5.6	0.8	13.7	64.0	55.6	73.9	30.4	15.8	43.1	1.0	0.2	2.4	0.7	0.1	1.7	13	2	20	84	79	90	66	65	67	
7	Region 22	77.3	75.8	78.3	28.9	27.6	30.0	5.1	2.3	7.4	65.1	58.1	71.4	29.9	22.7	39.0	1.0	0.4	2.2	0.9	0.3	1.9	10	3	21	81	67	94	65	62	68	
4	Region 23	76.6	75.4	77.8	28.4	27.1	30.0	6.6	2.5	11.6	60.7	46.2	68.8	32.7	19.6	51.3	0.9	0.4	1.4	0.7	0.4	1.0	9	4	15	78	62	90	64	60	67	
6	Region 24	75.8	71.8	78.3	30.2	26.5	33.7	3.6	1.6	8.1	64.7	60.2	69.8	31.8	22.1	37.0	1.2	0.0	2.9	0.9	0.0	1.7	15	5	29	69	56	82	62	59	65	
24	Region 25	74.8	70.7	76.6	29.0	26.3	32.1	5.9	0.8	26.1	63.5	47.6	69.7	30.6	17.6	51.0	0.9	0.0	2.4	0.7	0.0	1.9	15	6	33	68	56	88	62	59	67	
20	Region 26	75.2	72.2	77.2	28.6	25.5	34.2	6.9	1.7	26.8	65.3	60.0	71.2	27.8	6.2	35.1	1.1	0.2	3.5	0.9	0.1	2.1	16	4	33	75	59	100	64	60	70	
10	Region 27	74.4	71.4	76.8	26.0	22.8	32.2	3.6	0.8	9.0	61.2	53.4	75.2	35.2	15.8	44.8	1.0	0.7	1.6	0.9	0.6	1.4	15	7	28	77	75	79	64	64	65	
34	Region 28	75.6	72.0	78.8	29.7	23.4	38.5	6.1	0.1	32.0	64.3	49.0	76.5	29.6	8.2	50.0	0.8	0.0	2.3	0.6	0.0	1.5	13	3	28	73	55	86	63	59	66	
31	Region 29	76.8	74.2	79.4	31.5	27.9	38.0	7.7	0.6	16.8	69.8	58.8	78.7	22.5	10.5	38.2	0.7	0.0	1.6	0.5	0.0	1.3	15	4	49	79	59	89	65	60	67	
45	Region 30	76.3	71.6	78.5	33.4	26.3	39.5	11.1	2.1	32.5	68.2	46.9	80.1	20.7	6.1	47.5	0.5	0.0	2.2	0.4	0.0	1.4	10	3	32	77	60	95	64	60	68	
44	Region 31	77.5	74.3	80.2	32.1	26.7	37.1	8.7	0.4	23.0	67.0	58.9	81.8	24.3	9.8	37.0	0.5	0.0	1.6	0.4	0.0	1.4	9	2	23	77	61	93	64	60	68	
40	Region 32	77.1	73.5	80.3	32.8	20.9	37.2	10.4	0.6	21.8	66.2	39.5	76.1	23.4	9.7	55.5	0.7	0.0	1.8	0.5	0.0	1.4	11	4	21	79	70	89	65	62	67	
29	Region 33	76.1	71.3	79.7	32.3	26.1	38.2	7.4	1.0	23.3	66.1	47.6	73.9	26.4	9.1	50.5	0.8	0.0	2.0	0.6	0.0	1.7	12	4	23	70	59	81	62	60	65	
24	Region 34	76.3	70.9	79.5	29.5	19.9	37.4	8.1	0.1	31.8	62.2	37.3	74.3	29.7	13.0	62.6	0.7	0.0	1.8	0.5	0.0	1.2	11	5	24	82	72	94	65	63	68	
6	Region 35	77.3	73.7	79.9	29.4	22.4	34.1	5.2	1.2	13.2	62.9	51.5	72.1	31.9	20.7	46.8	0.9	0.2	1.8	0.8	0.2	1.5	12	8	15	79	50	89	65	58	67	
20	Region 36	76.3	74.9	79.5	31.7	28.0	36.2	8.4	0.0	28.6	62.2	44.2	71.9	29.4	13.5	55.2	0.9	0.1	3.0	0.6	0.0	2.5	13	4	30	73	49	87	63	57	66	
440	Ave. Yellow	76.5	70.7	82.6	31.2	19.9	43.8	7.4	0.0	32.5	65.9	5.6	81.8	26.7	6.1	88.1	0.8	0.0	3.5	0.6	0.0	2.5	12	1	49	76	41	100	64	55	70	
	Min. Yellow																															
	Max. Yellow																															


**TABLE 16: PHYSICAL QUALITY CHARACTERISTICS OF WHITE AND YELLOW MAIZE  
2011/12 - 2020/21**

Season	Number of samples	Test weight (kg/hl)			100 kernel mass (g)			Kernel size (%)									Breakage susceptibility (%)						Stress cracks (%)				
								Above 10 mm sieve			Above 8 mm sieve			Below 8 mm sieve			< 6.35 mm sieve			< 4.75 mm sieve							
		ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.
<b>White Maize</b>																											
2011/12	577	78.2	71.8	82.0	31.2	17.4	44.4	18.8	0.8	63.3	64.9	26.2	79.7	16.3	2.8	72.4	0.8	0.0	8.6	0.6	0.0	4.9	5	0	25		
2012/13	508	78.2	69.7	82.9	29.6	17.7	46.0	15.1	0.0	59.9	65.0	16.2	80.5	20.0	3.1	83.5	1.0	0.0	6.6	0.7	0.0	4.6	4	0	37		
2013/14	451	77.6	68.7	81.9	34.0	26.0	46.5	24.7	0.7	71.3	64.7	23.4	82.7	10.6	1.1	37.7	1.3	0.0	7.2	1.0	0.0	4.2	7	0	37		
2014/15	485	78.3	70.2	83.1	31.1	20.3	48.3	15.4	0.3	86.7	66.1	13.1	81.8	18.4	0.0	51.5	1.1	0.0	12.1	0.8	0.0	5.6	6	0	61		
2015/16	415	78.1	68.5	83.9	32.4	20.8	40.8	15.2	0.3	99.4	66.7	0.1	89.5	18.2	0.0	63.8	0.9	0.0	7.2	0.4	0.0	4.3	5	0	30		
2016/17	549	77.7	70.0	81.8	35.0	22.8	43.8	22.1	1.8	64.2	64.1	13.6	82.4	13.7	0.7	62.6	1.2	0.0	9.9	0.8	0.0	9.0	8	0	42		
2017/18	451	77.0	63.4	81.7	34.3	18.3	45.8	26.2	1.5	58.7	62.8	39.8	83.7	11.0	1.5	32.1	1.0	0.0	3.8	0.7	0.0	2.7	11	1	30		
2018/19	404	75.9	61.0	83.6	33.4	18.7	51.1	24.6	0.0	93.8	63.4	5.8	79.9	12.0	0.4	46.8	1.6	0.0	7.9	1.2	0.0	5.6	17	1	58		
2019/20	516	75.6	63.4	82.0	34.9	19.3	44.7	27.8	1.0	63.7	62.0	34.8	83.8	10.2	1.2	49.4	1.1	0.0	6.3	0.8	0.0	4.1	16	2	58		
2020/21	560	75.9	68.5	81.1	33.3	18.9	42.6	21.6	0.0	55.1	65.1	41.8	83.6	13.3	1.3	55.7	0.8	0.0	4.3	0.6	0.0	3.3	12	1	41		
<b>Weighted Average</b>		<b>77.3</b>			<b>32.9</b>			<b>21.1</b>			<b>64.5</b>			<b>14.4</b>			<b>1.1</b>			<b>0.8</b>			<b>9</b>				
<b>Minimum</b>		<b>61.0</b>			<b>17.4</b>			<b>0.0</b>			<b>0.1</b>			<b>0.0</b>			<b>0.0</b>			<b>0.0</b>			<b>0</b>				
<b>Maximum</b>		<b>83.9</b>			<b>51.1</b>			<b>99.4</b>			<b>89.5</b>			<b>83.5</b>			<b>12.1</b>			<b>9.0</b>			<b>61</b>				
<b>Yellow Maize</b>																											
2011/12	423	76.1	68.1	81.0	29.4	14.5	40.9	11.3	0.0	38.3	63.9	13.7	79.4	24.8	6.5	86.3	1.3	0.2	15.6	1.0	0.0	8.3	6	0	27		
2012/13	492	76.6	67.8	81.6	28.4	15.2	41.3	9.8	0.0	42.6	61.7	10.1	80.9	28.5	3.4	89.9	1.7	0.1	8.2	1.1	0.0	5.4	5	0	31		
2013/14	479	76.0	56.6	80.9	31.8	18.6	43.1	14.9	0.3	52.7	67.1	21.4	79.7	18.0	2.6	64.8	1.9	0.1	14.5	1.4	0.0	9.9	7	0	53		
2014/15	515	76.3	67.3	83.1	28.6	17.8	38.2	8.8	0.0	30.2	63.4	9.2	78.9	27.8	4.2	90.4	1.3	0.1	6.8	0.9	0.0	4.8	5	0	56		
2015/16	505	76.7	59.8	81.7	31.8	17.1	43.1	11.8	0.3	34.1	66.6	15.6	93.6	21.6	0.2	77.9	1.0	0.0	4.5	0.5	0.0	4.1	5	0	31		
2016/17	451	76.9	67.6	82.4	33.5	18.4	43.3	11.4	0.0	71.9	65.6	13.0	90.9	23.0	1.6	69.7	1.1	0.1	8.6	0.8	0.0	5.3	8	0	50		
2017/18	449	77.2	59.6	82.5	31.5	15.6	40.7	11.1	0.0	72.4	66.7	24.2	82.2	22.2	2.5	62.0	0.9	0.0	4.7	0.7	0.0	3.3	9	1	38		
2018/19	404	76.9	69.0	83.3	30.6	19.5	41.4	8.3	0.0	33.2	66.4	24.2	85.2	25.3	4.7	74.5	1.0	0.0	6.5	0.7	0.0	3.9	13	2	39		
2019/20	374	76.3	63.9	82.4	30.4	15.0	39.3	8.4	0.0	30.1	65.1	30.2	82.1	26.5	6.1	68.5	1.0	0.0	6.0	0.7	0.0	3.7	13	2	48		
2020/21	440	76.5	70.7	82.6	31.2	19.9	43.8	7.4	0.0	32.5	65.9	5.6	81.8	26.7	6.1	88.1	0.8	0.0	3.5	0.6	0.0	2.5	12	1	49		
<b>Weighted Average</b>		<b>76.5</b>			<b>30.7</b>			<b>10.4</b>			<b>65.2</b>			<b>24.4</b>			<b>1.2</b>			<b>0.8</b>			<b>8</b>				
<b>Minimum</b>		<b>56.6</b>			<b>14.5</b>			<b>0.0</b>			<b>5.6</b>			<b>0.2</b>			<b>0.0</b>			<b>0.0</b>			<b>0</b>				
<b>Maximum</b>		<b>83.3</b>			<b>43.8</b>			<b>72.4</b>			<b>93.6</b>			<b>90.4</b>			<b>15.6</b>			<b>9.9</b>			<b>56</b>				
<b>White &amp; Yellow Maize</b>																											
2011/12	1000	77.3	68.1	82.0	30.4	14.5	44.4	15.6	0.0	63.3	64.5	13.7	79.7	19.9	2.8	86.3	1.0	0.0	15.6	0.7	0.0	8.3	6	0	27		
2012/13	1000	77.4	67.8	82.9	29.0	15.2	46.0	12.5	0.0	59.9	63.4	10.1	80.9	24.2	3.1	89.9	1.4	0.0	8.2	0.9	0.0	5.4	5	0	37		
2013/14	930	76.8	56.6	81.9	32.9	18.6	46.5	19.6	0.3	71.3	65.9	23.4	82.7	14.4	1.1	64.8	1.6	0.0	14.5	1.2	0.0	9.9	7	0	53		
2014/15	1000	77.3	67.3	83.1	29.8	17.8	48.3	12.0	0.0	86.7	64.7	9.2	81.8	23.2	0.0	90.4	1.2	0.0	12.1	0.8	0.0	5.6	6	0	61		
2015/16	920	77.3	59.8	83.9	32.1	17.1	43.1	13.3	0.3	99.4	66.7	0.1	93.6	20.0	0.0	77.9	1.0	0.0	7.2	0.5	0.0	4.3	5	0	31		
2016/17	1000	77.3	67.6	82.4	34.3	18.4	43.8	17.3	0.0	71.9	64.8	13.0	90.9	17.9	0.7	69.7	1.2	0.0	9.9	0.8	0.0	9.0	11	0	40		
2017/18	900	77.1	59.6	82.5	32.9	15.6	45.8	18.7	0.0	72.4	64.7	24.2	83.7	16.6	1.5	62.0	1.0	0.0	4.7	0.7	0.0	3.3	10	1	38		
2018/19	808	76.4	61.0	83.6	32.0	18.7	51.1	16.4	0.0	93.8	64.9	5.8	85.2	18.7	0.4	74.5	1.3	0.0	7.9	0.9	0.0	5.6	15	1	58		
2019/20	890	75.9	63.4	82.4	33.0	15.0	44.7	19.7	0.0	63.7	63.3	30.2	83.8	17.1	1.2	68.5	1.1	0.0	6.3	0.8	0.0	4.1	15	2	58		
2020/21	1000	76.2	68.5	82.6	32.4	18.9	43.8	15.4	0.0	55.1	65.5	5.6	83.6	19.2	1.3	88.1	0.8	0.0	4.3	0.6	0.0	3.3	12	1	49		
<b>Weighted Average</b>		<b>76.9</b>			<b>31.8</b>			<b>16.0</b>			<b>64.8</b>			<b>19.2</b>			<b>1.2</b>			<b>0.8</b>			<b>9</b>				
<b>Minimum</b>		<b>56.6</b>			<b>14.5</b>			<b>0.0</b>			<b>0.1</b>			<b>0.0</b>			<b>0.0</b>			<b>0.0</b>			<b>0</b>				
<b>Maximum</b>		<b>83.9</b>			<b>51.1</b>			<b>99.4</b>			<b>93.6</b>			<b>90.4</b>			<b>15.6</b>			<b>9.9</b>			<b>61</b>				


**GRAPH 37: TEST WEIGHT OF WHITE AND YELLOW MAIZE OVER 10 SEASONS**




GRAPH 38: 100 KERNEL MASS OVER 10 SEASONS



GRAPH 39: KERNEL SIZE ABOVE 10 MM SIEVE OVER 10 SEASONS



GRAPH 40: KERNEL SIZE BELOW 8 MM SIEVE OVER 10 SEASONS



<b>TABLE 17: ROFF MILLING AND WHITENESS INDEX OF WHITE MAIZE ACCORDING TO GRADE (2020/21)</b>																									
Number of samples	Region	RoFF Milling										Whiteness index													
		Break 1, %		Break 2, %		Break 3, %		Grits, %		Chop, %		Extraction, % (Total meal)		Whiteness index unsifted		Whiteness index sifted 87:13									
		ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.						
<b>GRADE: WM1</b>																									
8	12	15.0	14.1	15.4	9.2	8.7	9.6	15.8	14.5	17.0	36.3	35.8	37.6	23.7	22.3	26.0	76.3	74.0	77.7	40.3	38.1	43.6	30.7	28.0	36.1
23	13	14.8	11.8	17.0	8.9	8.3	9.8	15.8	14.9	16.6	37.2	33.7	41.2	23.4	20.5	25.7	76.6	74.3	79.5	38.6	31.9	42.5	29.6	21.7	36.4
25	14	15.5	13.9	17.1	9.4	8.8	10.1	16.7	13.6	18.7	36.2	28.1	41.5	22.2	19.1	33.4	77.8	66.6	80.9	40.4	36.9	44.1	31.1	24.5	39.9
6	16	15.5	14.9	16.6	9.1	8.6	9.8	17.5	16.5	19.9	35.0	33.1	37.1	22.9	21.2	24.1	77.1	75.9	78.8	38.6	35.5	40.8	29.4	25.3	33.8
15	17	15.8	14.5	16.9	9.2	8.2	9.7	16.3	13.9	19.3	36.5	31.9	40.0	22.2	20.1	25.9	77.8	74.1	79.9	39.7	36.7	43.0	30.0	26.1	34.8
24	18	14.6	11.7	16.3	9.0	8.5	9.7	16.5	14.9	19.9	37.4	34.6	41.0	22.5	21.2	26.3	77.5	73.7	78.8	38.1	29.1	43.1	28.8	19.7	37.0
17	19	14.3	12.2	16.2	9.0	8.3	9.8	15.7	15.1	16.6	37.5	31.6	41.6	23.6	21.6	26.3	76.4	73.7	78.4	39.5	29.6	47.7	29.6	25.2	35.2
15	20	14.8	12.7	17.9	8.8	8.1	9.5	15.6	13.4	16.7	37.4	32.1	40.1	23.4	21.6	25.5	76.6	74.5	78.4	39.6	32.8	44.1	29.7	25.9	32.7
27	21	15.2	13.0	19.1	9.2	8.4	10.6	17.2	9.6	20.1	36.4	31.5	40.1	21.8	19.0	24.5	78.1	75.5	81.0	37.5	32.4	42.1	27.5	21.0	39.0
9	22	15.4	13.7	17.1	9.5	8.9	10.2	17.7	16.3	20.5	34.8	32.5	36.8	22.5	19.8	24.3	77.5	75.7	80.2	39.4	34.7	43.1	28.8	24.0	32.7
31	23	15.5	13.7	17.5	9.4	8.8	10.5	17.9	16.1	20.4	35.6	30.3	39.4	21.6	19.6	23.8	78.4	76.2	80.4	37.5	32.6	43.8	27.5	21.6	40.8
10	24	15.2	13.3	19.0	9.3	8.6	10.1	17.0	14.8	20.1	36.1	28.0	40.7	22.4	21.0	26.5	77.6	73.5	79.0	36.9	33.4	40.3	27.2	24.3	31.4
5	26	16.2	14.5	17.9	9.6	9.2	10.1	16.4	15.6	17.0	35.9	33.9	39.0	21.9	20.4	23.7	78.1	76.3	79.6	38.9	36.1	41.7	28.8	26.7	31.3
3	27	15.5	14.3	16.6	9.4	9.2	9.7	16.6	14.6	18.8	35.6	33.8	37.0	22.9	20.5	25.8	77.1	74.2	79.5	38.2	32.3	43.3	27.6	22.1	32.6
12	28	13.3	7.0	17.4	8.9	7.0	9.9	17.3	15.6	19.1	39.4	34.5	49.4	21.0	18.7	22.9	79.0	77.1	81.3	32.8	21.1	41.2	24.3	14.3	35.1
26	29	13.3	10.9	15.8	8.7	8.0	9.4	16.9	15.4	19.6	38.3	32.0	41.8	22.7	19.5	24.9	77.3	75.1	80.5	34.5	24.8	40.8	25.3	12.6	37.6
23	30	12.9	10.7	15.5	8.8	8.3	9.6	16.5	15.1	18.7	39.2	35.7	42.9	22.7	20.7	25.8	77.3	74.2	79.3	33.7	28.6	38.2	24.6	18.0	29.0
10	31	13.0	10.4	13.9	8.6	8.0	9.3	17.0	15.5	18.8	39.0	37.2	43.5	22.5	20.2	25.3	77.5	74.7	79.8	32.0	28.8	35.0	21.8	17.3	25.9
13	32	13.6	11.2	16.8	8.8	8.0	9.5	17.0	15.8	18.7	38.5	34.3	42.9	22.0	20.5	23.5	78.0	76.5	79.5	32.8	26.3	41.9	23.2	16.4	33.5
40	33	13.4	11.5	15.7	8.7	8.3	9.9	16.5	14.4	18.6	39.5	35.3	42.2	21.9	18.8	26.4	78.1	73.6	81.2	33.7	28.9	39.1	24.9	20.1	33.7
21	34	13.6	11.9	16.1	8.7	8.2	9.6	16.8	14.2	20.4	38.5	31.7	40.6	22.5	18.6	28.9	77.5	71.1	81.4	36.3	29.4	43.2	28.2	19.1	39.7
6	35	13.0	11.6	15.5	8.6	8.3	9.2	17.7	16.1	20.8	39.9	35.1	43.7	20.8	18.8	23.0	79.2	77.0	81.2	34.2	28.8	38.1	25.0	19.9	29.0
17	36	13.0	11.7	15.1	8.5	7.9	9.4	16.5	15.7	17.8	40.1	37.9	42.7	21.9	19.6	23.9	78.1	76.1	80.4	34.7	30.5	37.2	26.4	19.9	34.5
2	25	16.2	16.1	16.3	9.1	9.0	9.2	16.2	15.2	17.3	35.6	33.9	37.2	22.9	20.2	25.6	77.1	74.4	79.8	35.2	34.1	36.3	24.0	24.0	24.0
<b>388</b>	<b>Ave. WM1</b>	<b>14.3</b>	<b>7.0</b>	<b>19.1</b>	<b>9.0</b>	<b>7.0</b>	<b>10.6</b>	<b>16.7</b>	<b>9.6</b>	<b>20.8</b>	<b>37.6</b>	<b>28.0</b>	<b>49.4</b>	<b>22.4</b>	<b>18.6</b>	<b>33.4</b>	<b>77.6</b>	<b>66.6</b>	<b>81.4</b>	<b>36.7</b>	<b>21.1</b>	<b>47.7</b>	<b>27.3</b>	<b>12.6</b>	<b>40.8</b>
	<b>Min. WM1</b>																								
	<b>Max. WM1</b>																								



**TABLE 17: ROFF MILLING AND WHITENESS INDEX OF WHITE MAIZE ACCORDING TO GRADE (2020/21)**

(continue)

Number of samples	Region	Roff Milling												Whiteness index											
		Break 1, %			Break 2, %			Break 3, %			Grits, %			Chop, %			Extraction, % (Total meal)			Whiteness index unsifted			Whiteness index sifted 87:13		
		ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.
<b>GRADE: WM2</b>																									
1	11	12.9	12.9	12.9	8.4	8.4	8.4	16.7	16.7	16.7	41.4	41.4	41.4	20.5	20.5	20.5	79.5	79.5	79.5	34.0	34.0	34.0	25.9	25.9	25.9
3	12	15.1	14.8	15.5	9.6	8.6	10.4	15.9	15.6	16.4	36.1	35.5	36.9	23.3	22.9	24.1	76.7	75.9	77.1	39.9	38.1	41.5	35.4	29.9	40.4
4	13	14.5	13.3	16.6	9.1	8.5	9.5	16.3	15.3	17.2	37.6	35.1	39.2	22.6	20.5	24.0	77.4	76.0	79.5	39.4	36.4	41.5	31.9	27.7	37.4
9	14	15.2	12.7	16.7	9.2	8.3	9.8	16.6	14.2	17.4	36.4	34.5	38.3	22.6	21.0	26.6	77.4	73.4	79.0	39.9	35.7	43.3	30.0	26.6	34.2
1	16	14.3	14.3	14.3	8.9	8.9	8.9	17.2	17.2	17.2	37.4	37.4	37.4	22.3	22.3	22.3	77.7	77.7	77.7	33.2	33.2	33.2	23.7	23.7	23.7
2	17	15.3	15.1	15.5	9.4	9.0	9.7	16.0	15.4	16.6	36.4	36.3	36.6	22.9	22.0	23.9	77.1	76.1	78.0	38.1	37.6	38.6	29.2	27.5	30.9
3	18	14.7	13.6	15.8	9.0	8.7	9.2	16.3	15.9	16.7	37.4	36.5	38.4	22.5	21.0	23.4	77.5	76.6	79.0	39.5	36.6	43.4	34.7	28.0	42.8
5	19	15.4	13.7	18.5	9.3	8.5	10.2	15.7	15.0	16.7	34.5	31.4	36.5	25.1	23.2	26.5	74.9	73.5	76.8	42.3	40.9	45.1	34.8	31.1	44.3
1	20	16.8	16.8	16.8	9.9	9.9	9.9	15.9	15.9	15.9	34.6	34.6	34.6	22.8	22.8	22.8	77.2	77.2	77.2	37.3	37.3	37.3	26.6	26.6	26.6
3	21	15.2	12.5	16.6	9.6	9.1	9.9	18.3	17.1	19.9	35.8	32.6	39.1	21.2	20.8	21.4	78.8	78.6	79.2	36.9	30.7	40.2	25.4	20.0	28.4
2	22	15.8	15.4	16.2	9.7	9.7	9.7	15.6	15.2	16.0	34.6	34.1	35.2	24.3	24.0	24.6	75.7	75.4	76.0	39.3	38.4	40.2	30.2	30.1	30.3
6	23	15.7	13.8	17.5	9.5	9.3	10.0	17.2	16.2	19.6	35.3	31.9	38.9	22.2	20.9	25.1	77.8	74.9	79.1	38.8	34.4	44.2	29.0	24.7	37.3
2	24	15.8	15.5	16.1	9.6	9.4	9.9	16.9	15.2	18.6	33.2	31.8	34.6	24.5	21.4	27.6	75.5	72.4	78.6	37.3	34.5	40.2	28.2	25.2	31.3
2	26	16.2	16.1	16.3	9.5	9.2	9.9	16.4	15.8	17.0	36.8	36.7	36.8	21.1	20.1	22.1	78.9	77.9	79.9	36.7	33.9	39.4	24.7	20.8	28.6
2	27	15.9	15.7	16.1	9.8	9.8	9.9	16.4	16.3	16.4	35.9	35.8	35.9	22.0	21.8	22.2	78.0	77.8	78.2	42.6	41.9	43.4	32.2	30.6	33.7
1	29	11.8	11.8	11.8	8.6	8.6	8.6	17.2	17.2	17.2	37.8	37.8	37.8	24.6	24.6	24.6	75.4	75.4	75.4	29.8	29.8	29.8	20.9	20.9	20.9
7	30	12.4	10.8	14.4	8.7	8.2	9.1	16.1	15.3	17.0	40.9	38.1	43.2	21.9	20.1	23.7	78.1	76.3	79.9	30.9	28.1	35.3	21.2	16.9	27.8
7	31	12.4	11.9	13.4	8.7	8.2	9.3	16.4	15.2	18.8	40.2	36.9	42.3	22.3	20.4	25.2	77.7	74.8	79.6	31.2	27.7	35.2	22.7	18.9	25.6
8	32	12.8	11.2	15.0	8.5	7.8	9.2	17.7	15.8	20.5	39.7	35.8	42.8	21.2	19.6	22.7	78.8	77.3	80.4	29.0	25.4	32.9	19.2	14.4	24.0
10	33	13.2	11.3	16.4	9.0	8.4	9.8	16.2	15.4	17.9	39.3	35.2	42.9	22.4	21.0	23.5	77.6	76.5	79.0	32.8	30.2	39.6	24.6	20.1	36.1
4	34	13.2	12.2	14.5	8.6	8.4	8.9	16.3	16.1	16.6	39.5	36.8	40.6	22.4	21.4	23.9	77.6	76.1	78.6	35.5	33.1	37.0	28.8	23.0	36.6
1	35	12.9	12.9	12.9	8.2	8.2	8.2	16.6	16.6	16.6	39.4	39.4	39.4	22.9	22.9	22.9	77.1	77.1	77.1	30.1	30.1	30.1	20.9	20.9	20.9
6	36	12.5	11.6	13.1	9.2	8.0	12.4	16.8	16.3	17.4	39.5	38.6	40.2	22.0	19.3	23.1	78.0	76.9	80.7	34.0	31.5	36.5	26.6	21.6	38.0
<b>90</b>	<b>Ave. WM2</b>	<b>14.0</b>			<b>9.1</b>			<b>16.6</b>			<b>37.9</b>			<b>22.5</b>			<b>77.5</b>			<b>35.6</b>			<b>26.9</b>		
	<b>Min. WM2</b>	<b>10.8</b>			<b>7.8</b>			<b>14.2</b>			<b>31.4</b>			<b>19.3</b>			<b>72.4</b>			<b>25.4</b>			<b>14.4</b>		
	<b>Max. WM2</b>	<b>18.5</b>			<b>12.4</b>			<b>20.5</b>			<b>43.2</b>			<b>27.6</b>			<b>80.7</b>			<b>45.1</b>			<b>44.3</b>		

**TABLE 17: ROFF MILLING AND WHITENESS INDEX OF WHITE MAIZE ACCORDING TO GRADE (2020/21)**  
(continue)


Number of samples	Region	Roff Milling												Whiteness index											
		Break 1, %			Break 2, %			Break 3, %			Grits, %			Chop, %			Extraction, % (Total meal)			Whiteness index unsifted			Whiteness index sifted 87:13		
		ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.
<b>GRADE: WM3</b>																									
1	13	15.7	15.7	15.7	9.3	9.3	9.3	15.3	15.3	15.3	32.9	32.9	32.9	26.8	26.8	26.8	73.2	73.2	73.2	42.3	42.3	42.3	30.6	30.6	30.6
3	14	16.0	15.3	16.7	9.5	9.4	9.6	16.5	16.1	16.9	36.4	36.1	36.8	21.7	21.3	22.1	78.3	77.9	78.7	41.2	40.3	41.9	31.8	30.9	33.2
1	16	14.8	14.8	14.8	9.7	9.7	9.7	20.7	20.7	20.7	35.3	35.3	35.3	19.5	19.5	19.5	80.5	80.5	80.5	33.3	33.3	33.3	21.2	21.2	21.2
8	17	14.9	12.5	16.3	9.3	8.6	9.9	16.7	15.3	18.6	36.0	33.1	40.1	23.2	20.9	24.9	76.8	75.1	79.1	36.4	28.7	45.6	27.3	22.8	36.8
1	18	12.6	12.6	12.6	8.9	8.9	8.9	16.2	16.2	16.2	41.9	41.9	41.9	20.5	20.5	20.5	79.5	79.5	79.5	33.4	33.4	33.4	24.9	24.9	24.9
5	19	15.4	14.3	18.6	9.1	8.7	9.7	15.9	15.3	17.0	36.0	32.7	37.6	23.6	22.0	24.9	76.4	75.1	78.0	39.4	37.3	42.2	29.9	29.1	31.2
1	20	13.9	13.9	13.9	8.9	8.9	8.9	15.5	15.5	15.5	37.3	37.3	37.3	24.4	24.4	24.4	75.6	75.6	75.6	38.2	38.2	38.2	31.1	31.1	31.1
3	21	15.8	14.8	16.4	9.4	9.0	9.9	17.7	16.6	19.4	34.5	32.6	36.9	22.5	21.7	23.2	77.5	76.8	78.3	40.4	38.5	42.6	28.6	27.3	31.3
5	23	15.8	15.5	16.1	9.9	9.3	10.4	18.4	16.5	19.7	34.5	32.2	35.4	21.5	19.4	24.1	78.5	75.9	80.6	35.2	30.0	39.4	24.7	18.8	28.4
1	26	14.8	14.8	14.8	8.9	8.9	8.9	15.5	15.5	15.5	36.9	36.9	36.9	23.8	23.8	23.8	76.2	76.2	76.2	35.9	35.9	35.9	24.9	24.9	24.9
2	30	13.7	13.7	13.8	8.8	8.6	9.1	17.5	16.0	19.0	37.4	36.3	38.5	22.5	21.9	23.2	77.5	76.8	78.1	26.7	20.0	33.4	14.6	5.1	24.1
2	31	13.1	11.4	14.8	8.5	8.4	8.5	16.2	16.0	16.4	39.8	37.4	42.2	22.4	22.0	22.9	77.6	77.1	78.0	26.6	17.7	35.5	15.9	6.3	25.5
1	32	14.8	14.8	14.8	9.1	9.1	9.1	17.3	17.3	17.3	35.7	35.7	35.7	23.1	23.1	23.1	76.9	76.9	76.9	32.1	32.1	32.1	21.1	21.1	21.1
1	33	15.1	15.1	15.1	8.7	8.7	8.7	17.8	17.8	17.8	35.7	35.7	35.7	22.7	22.7	22.7	77.3	77.3	77.3	15.6	15.6	15.6	3.1	3.1	3.1
1	34	14.0	14.0	14.0	8.2	8.2	8.2	16.8	16.8	16.8	40.0	40.0	40.0	20.9	20.9	20.9	79.1	79.1	79.1	33.6	33.6	33.6	25.9	25.9	25.9
36	Ave. WM3	15.0			9.2			16.9			36.2			22.6			77.4			35.5			25.5		
	Min. WM3	11.4			8.2			15.3			32.2			19.4			73.2			15.6			3.1		
	Max. WM3	18.6			10.4			20.7			42.2			26.8			80.6			45.6			36.8		

**TABLE 17: ROFF MILLING AND WHITENESS INDEX OF WHITE MAIZE ACCORDING TO GRADE (2020/21)**  
(continue)


Number of samples	Region	Roff Milling												Whiteness index											
		Break 1, %			Break 2, %			Break 3, %			Grits, %			Chop, %			Extraction, % (Total meal)			Whiteness index unsifted			Whiteness index sifted 87:13		
		ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.
<b>CLASS: COM</b>																									
1	11	13.7	13.7	13.7	8.8	8.8	8.8	17.8	17.8	17.8	35.0	35.0	35.0	24.8	24.8	24.8	75.2	75.2	75.2	41.1	41.1	41.1	29.9	29.9	29.9
3	12	14.5	14.0	15.2	8.4	8.1	8.6	16.5	16.2	16.9	38.1	37.4	39.4	22.5	22.0	23.0	77.5	77.0	78.0	37.0	36.3	37.9	30.4	26.3	37.2
1	13	14.1	14.1	14.1	8.2	8.2	8.2	15.4	15.4	15.4	37.8	37.8	37.8	24.5	24.5	24.5	75.5	75.5	75.5	36.9	36.9	36.9	27.3	27.3	27.3
3	14	14.6	12.9	15.5	9.5	9.4	9.5	16.0	15.2	17.1	37.7	36.7	38.6	22.3	20.9	24.7	77.7	75.3	79.1	42.3	40.4	43.5	32.8	29.6	36.3
4	17	16.1	15.0	18.2	9.2	8.7	9.8	15.7	14.2	17.3	34.8	32.0	35.9	24.2	23.0	26.8	75.8	73.2	77.0	40.3	39.0	43.0	29.8	27.3	33.2
1	18	12.9	12.9	12.9	9.0	9.0	9.0	17.2	17.2	17.2	39.4	39.4	39.4	21.5	21.5	21.5	78.5	78.5	78.5	30.5	30.5	30.5	19.7	19.7	19.7
2	19	14.2	12.8	15.6	9.0	8.3	9.7	15.9	15.6	16.2	37.2	34.6	39.7	23.8	23.6	24.0	76.2	76.0	76.4	40.5	39.3	41.8	33.5	30.6	36.4
3	21	14.2	11.9	15.7	9.5	9.3	9.7	17.4	16.6	18.8	36.1	34.0	39.4	22.9	22.3	23.6	77.1	76.4	77.7	39.8	35.9	42.9	29.4	26.1	33.5
2	22	15.6	15.5	15.8	9.2	8.8	9.6	15.5	14.7	16.2	34.0	32.9	35.0	25.7	25.4	26.0	74.3	74.0	74.6	37.7	36.8	38.6	26.8	25.3	28.3
5	23	15.9	14.8	17.6	9.9	9.2	10.1	19.0	17.0	20.0	33.7	32.6	36.3	21.5	21.2	21.8	78.5	78.2	78.8	35.8	34.6	37.8	25.6	23.6	26.5
1	24	14.9	14.9	14.9	8.9	8.9	8.9	17.0	17.0	17.0	37.0	37.0	37.0	22.2	22.2	22.2	77.8	77.8	77.8	35.8	35.8	35.8	25.5	25.5	25.5
1	28	15.1	15.1	15.1	8.5	8.5	8.5	17.2	17.2	17.2	37.6	37.6	37.6	21.7	21.7	21.7	78.3	78.3	78.3	31.5	31.5	31.5	22.8	22.8	22.8
2	30	14.5	14.3	14.8	9.2	9.0	9.5	16.6	16.4	16.8	37.6	37.4	37.9	22.0	21.1	23.0	78.0	77.0	78.9	34.9	33.6	36.2	25.0	24.7	25.2
1	31	16.1	16.1	16.1	10.5	10.5	10.5	18.5	18.5	18.5	33.6	33.6	33.6	21.3	21.3	21.3	78.7	78.7	78.7	36.5	36.5	36.5	26.9	26.9	26.9
4	32	13.3	11.2	14.5	8.7	8.0	9.1	17.0	16.2	18.8	39.1	35.4	42.6	21.9	20.0	24.4	78.1	75.6	80.0	32.1	29.3	34.3	22.6	19.1	27.5
3	33	10.5	9.1	11.6	8.8	8.2	9.3	15.1	14.5	15.6	42.9	41.2	45.1	22.8	21.9	23.5	77.2	76.5	78.1	20.8	7.9	35.0	11.0	-5.0	26.6
4	34	13.6	12.0	15.1	8.5	8.1	9.1	16.4	15.8	17.0	38.5	34.7	40.1	23.1	21.4	25.3	76.9	74.7	78.6	33.8	30.0	39.4	25.5	20.7	31.9
4	36	13.5	12.5	14.5	8.6	8.3	9.0	16.1	15.7	16.5	38.4	36.9	40.2	23.3	22.8	23.7	76.7	76.3	77.2	33.5	30.6	35.8	23.2	18.8	26.3
1	25	19.2	19.2	19.2	10.7	10.7	10.7	16.4	16.4	16.4	31.6	31.6	31.6	22.2	22.2	22.2	77.8	77.8	77.8	38.1	38.1	38.1	28.4	28.4	28.4
46	Ave. COM	14.4			9.1			16.7			37.1			22.8			77.2			35.5			26.0		
	Min. COM	9.1			8.0			14.2			31.6			20.0			73.2			7.9			-5.0		
	Max. COM	19.2			10.7			20.0			45.1			26.8			80.0			43.5			37.2		
560	Ave. WM	14.3			9.0			16.7			37.5			22.4			77.6			36.3			27.0		
	Min. WM	7.0			7.0			9.6			28.0			18.6			66.6			7.9			-5.0		
	Max. WM	19.2			12.4			20.8			49.4			33.4			81.4			47.7			44.3		

TABLE 18: ROFF MILLING AND WHITENESS INDEX OF WHITE MAIZE (2020/21)																									
Number of samples	Region	Roff Milling										Whiteness index													
		Break 1, %		Break 2, %		Break 3, %		Grits, %		Chop, %		Extraction, % (Total meal)		Whiteness index unsifted		Whiteness index sifted 87:13									
		ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.						
2	Region 11	13.3	12.9	13.7	8.6	8.4	8.8	17.3	16.7	17.8	38.2	35.0	41.4	22.6	20.5	24.8	77.4	75.2	79.5	37.6	34.0	41.1	27.9	25.9	29.9
14	Region 12	14.9	14.0	15.5	9.1	8.1	10.4	16.0	14.5	17.0	36.7	35.5	39.4	23.3	22.0	26.0	76.7	74.0	78.0	39.5	36.3	43.6	31.6	26.3	40.4
29	Region 13	14.7	11.8	17.0	8.9	8.2	9.8	15.8	14.9	17.2	37.1	32.9	41.2	23.4	20.5	26.8	76.6	73.2	79.5	38.7	31.9	42.5	29.9	21.7	37.4
40	Region 14	15.4	12.7	17.1	9.4	8.3	10.1	16.6	13.6	18.7	36.4	28.1	41.5	22.2	19.1	33.4	77.8	66.6	80.9	40.5	35.7	44.1	31.1	24.5	39.9
8	Region 16	15.2	14.3	16.6	9.2	8.6	9.8	17.8	16.5	20.7	35.4	33.1	37.4	22.4	19.5	24.1	77.6	75.9	80.5	37.3	33.2	40.8	27.6	21.2	33.8
29	Region 17	15.5	12.5	18.2	9.2	8.2	9.9	16.3	13.9	19.3	36.1	31.9	40.1	22.8	20.1	26.8	77.2	73.2	79.9	38.7	28.7	45.6	29.2	22.8	36.8
29	Region 18	14.5	11.7	16.3	9.0	8.5	9.7	16.5	14.9	19.9	37.6	34.6	41.9	22.4	20.5	26.3	77.6	73.7	79.5	37.8	29.1	43.4	29.0	19.7	42.8
29	Region 19	14.7	12.2	18.6	9.1	8.3	10.2	15.7	15.0	17.0	36.7	31.4	41.6	23.9	21.6	26.5	76.1	73.5	78.4	40.1	29.6	47.7	30.8	25.2	44.3
17	Region 20	14.8	12.7	17.9	8.9	8.1	9.9	15.6	13.4	16.7	37.2	32.1	40.1	23.4	21.6	25.5	76.6	74.5	78.4	39.4	32.8	44.1	29.6	25.9	32.7
36	Region 21	15.2	11.9	19.1	9.3	8.4	10.6	17.4	9.6	20.1	36.2	31.5	40.1	21.9	19.0	24.5	78.0	75.5	81.0	37.9	30.7	42.9	27.6	20.0	39.0
13	Region 22	15.5	13.7	17.1	9.5	8.8	10.2	17.0	14.7	20.5	34.6	32.5	36.8	23.3	19.8	26.0	76.7	74.0	80.2	39.1	34.7	43.1	28.7	24.0	32.7
47	Region 23	15.6	13.7	17.6	9.5	8.8	10.5	18.0	16.1	20.4	35.3	30.3	39.4	21.7	19.4	25.1	78.3	74.9	80.6	37.3	30.0	44.2	27.2	18.8	40.8
13	Region 24	15.3	13.3	19.0	9.3	8.6	10.1	17.0	14.8	20.1	35.7	28.0	40.7	22.7	21.0	27.6	77.3	72.4	79.0	36.9	33.4	40.3	27.2	24.3	31.4
3	Region 25	17.2	16.1	19.2	9.6	9.0	10.7	16.3	15.2	17.3	34.2	31.6	37.2	22.7	20.2	25.6	77.3	74.4	79.8	36.2	34.1	38.1	25.5	24.0	28.4
8	Region 26	16.1	14.5	17.9	9.5	8.9	10.1	16.3	15.5	17.0	36.2	33.9	39.0	22.0	20.1	23.8	78.0	76.2	79.9	38.0	33.9	41.7	27.3	20.8	31.3
5	Region 27	15.7	14.3	16.6	9.6	9.2	9.9	16.5	14.6	18.8	35.7	33.8	37.0	22.5	20.5	25.8	77.5	74.2	79.5	40.0	32.3	43.4	29.5	22.1	33.7
13	Region 28	13.4	7.0	17.4	8.9	7.0	9.9	17.3	15.6	19.1	39.3	34.5	49.4	21.1	18.7	22.9	78.9	77.1	81.3	32.7	21.1	41.2	24.2	14.3	35.1
27	Region 29	13.2	10.9	15.8	8.7	8.0	9.4	16.9	15.4	19.6	38.3	32.0	41.8	22.8	19.5	24.9	77.2	75.1	80.5	34.3	24.8	40.8	25.2	12.6	37.6
34	Region 30	12.9	10.7	15.5	8.8	8.2	9.6	16.5	15.1	19.0	39.3	35.7	43.2	22.5	20.1	25.8	77.5	74.2	79.9	32.8	20.0	38.2	23.4	5.1	29.0
20	Region 31	12.9	10.4	16.1	8.7	8.0	10.5	16.8	15.2	18.8	39.2	33.6	43.5	22.4	20.2	25.3	77.6	74.7	79.8	31.4	17.7	36.5	21.8	6.3	26.9
26	Region 32	13.4	11.2	16.8	8.7	7.8	9.5	17.3	15.8	20.5	38.9	34.3	42.9	21.8	19.6	24.4	78.2	75.6	80.4	31.5	25.4	41.9	21.8	14.4	33.5
54	Region 33	13.2	9.1	16.4	8.8	8.2	9.9	16.4	14.4	18.6	39.6	35.2	45.1	22.1	18.8	26.4	77.9	73.6	81.2	32.5	7.9	39.6	23.6	-5.0	36.1
30	Region 34	13.5	11.9	16.1	8.7	8.1	9.6	16.6	14.2	20.4	38.6	31.7	40.6	22.5	18.6	28.9	77.5	71.1	81.4	35.8	29.4	43.2	27.8	19.1	39.7
7	Region 35	13.0	11.6	15.5	8.6	8.2	9.2	17.5	16.1	20.8	39.9	35.1	43.7	21.1	18.8	23.0	78.9	77.0	81.2	33.7	28.8	38.1	24.4	19.9	29.0
27	Region 36	13.0	11.6	15.1	8.7	7.9	12.4	16.5	15.7	17.8	39.7	36.9	42.7	22.1	19.3	23.9	77.9	76.1	80.7	34.3	30.5	37.2	26.0	18.8	38.0
560	Ave. White	14.3			9.0			16.7			37.5			22.4			77.6			36.3			27.0		
	Min. White		7.0		7.0			9.6			28.0			18.6			66.6			7.9			-5.0		
	Max. White			19.2			12.4	20.8			49.4			33.4			81.4			47.7			44.3		


GRAPH 41: ROFF MILL BREAK 1 MEAL FRACTION %  
SINCE 2012/13




GRAPH 42: ROFF MILL BREAK 2 MEAL FRACTION %  
SINCE 2012/13




GRAPH 43: ROFF MILL BREAK 3 MEAL FRACTION %  
SINCE 2012/13



GRAPH 44: ROFF MILL GRITS FRACTION %  
SINCE 2012/13



GRAPH 45: ROFF MILL CHOP FRACTION %  
SINCE 2012/13



GRAPH 46: ROFF MILL TOTAL MEAL EXTRACTION %  
SINCE 2012/13




TABLE 19: NUTRITIONAL VALUES OF WHITE MAIZE ACCORDING TO GRADE (2020/21)												TABLE 20: NUTRITIONAL VALUES OF YELLOW MAIZE ACCORDING TO GRADE (2020/21)																
Number of samples	Region	Fat % (db)			Protein % (db)			Starch % (db)			Crude Fibre % (db)			Number of samples	Region	Fat % (db)			Protein % (db)			Starch % (db)			Crude Fibre % (db)			
		ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.			ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	
<b>GRADE: WM1</b>																												
-	Region 10	-	-	-	-	-	-	-	-	-	-	-	-	24	Region 10	3.8	3.2	4.2	8.5	7.9	9.1	75.4	74.1	76.4	2.2	2.0	2.4	
-	Region 11	-	-	-	-	-	-	-	-	-	-	-	-	13	Region 11	3.9	3.5	4.2	8.4	7.8	8.9	75.4	74.5	76.1	2.2	2.1	2.3	
8	Region 12	4.1	3.9	4.8	8.3	7.9	8.7	75.3	74.9	75.7	2.3	2.3	2.4	-	Region 12	-	-	-	-	-	-	-	-	-	-	-	-	
23	Region 13	4.1	3.6	4.3	8.1	7.2	8.9	75.6	74.5	76.6	2.3	2.2	2.4	5	Region 13	3.9	3.6	4.5	8.5	8.0	8.8	75.8	75.5	76.2	2.2	2.2	2.3	
25	Region 14	4.0	3.4	4.5	7.7	7.1	8.6	76.1	75.2	76.8	2.2	2.2	2.4	6	Region 14	4.0	3.7	4.0	8.6	8.3	8.8	75.2	74.6	75.6	2.3	2.2	2.3	
-	Region 15	-	-	-	-	-	-	-	-	-	-	-	-	-	Region 15	-	-	-	-	-	-	-	-	-	-	-	-	
6	Region 16	4.1	3.9	4.3	7.5	6.9	8.0	76.2	75.4	76.8	2.2	2.2	2.3	-	Region 16	-	-	-	-	-	-	-	-	-	-	-	-	
15	Region 17	3.9	3.6	4.3	7.9	7.0	8.6	76.0	74.9	76.6	2.2	2.2	2.3	3	Region 17	4.0	3.9	4.1	8.8	7.7	9.7	75.0	73.7	76.3	2.3	2.2	2.3	
24	Region 18	4.0	3.4	4.7	8.4	7.5	10.1	75.6	74.1	76.3	2.2	2.2	2.3	6	Region 18	3.9	3.6	4.2	8.8	8.3	9.2	75.0	74.5	76.0	2.3	2.2	2.4	
17	Region 19	4.0	3.5	4.3	8.3	7.5	8.9	75.5	75.1	76.0	2.2	2.1	2.4	7	Region 19	4.0	3.8	4.1	9.2	8.9	9.5	74.9	74.2	75.4	2.3	2.2	2.4	
15	Region 20	4.0	3.5	4.3	8.1	6.8	8.8	75.5	74.7	76.2	2.2	2.1	2.3	7	Region 20	3.9	3.7	4.5	8.4	7.9	8.8	75.2	74.2	75.9	2.3	2.1	2.4	
27	Region 21	4.0	3.6	4.4	8.1	7.6	9.0	75.7	74.2	76.3	2.2	2.2	2.3	3	Region 21	4.1	4.0	4.2	8.9	8.7	9.2	74.8	74.5	75.2	2.3	2.2	2.3	
9	Region 22	3.9	3.6	4.2	7.8	7.1	8.8	76.0	75.7	76.5	2.3	2.3	2.4	5	Region 22	4.3	4.2	4.5	8.6	8.2	9.3	74.9	74.3	75.5	2.3	2.3	2.4	
31	Region 23	4.0	3.6	4.4	7.8	7.1	8.6	76.0	75.2	76.5	2.3	2.2	2.4	4	Region 23	4.0	3.6	4.4	8.5	8.2	8.9	75.3	75.0	75.5	2.3	2.2	2.4	
10	Region 24	4.1	3.8	4.8	7.8	6.1	8.5	75.9	75.5	76.4	2.3	2.2	2.4	4	Region 24	3.8	3.6	4.0	8.3	7.3	8.8	75.5	75.1	76.2	2.2	2.2	2.4	
2	Region 25	4.2	4.0	4.5	7.5	7.0	7.9	75.5	75.3	75.8	2.3	2.2	2.3	18	Region 25	3.9	3.6	4.4	8.2	7.3	9.6	74.9	73.6	76.3	2.3	2.2	2.5	
5	Region 26	4.1	4.0	4.2	7.5	7.3	7.9	76.1	76.1	76.2	2.3	2.2	2.4	11	Region 26	4.0	3.6	4.7	8.7	8.3	9.5	74.8	73.8	75.4	2.3	2.2	2.4	
3	Region 27	4.2	3.8	4.4	7.3	7.0	7.9	75.9	75.6	76.5	2.3	2.2	2.3	4	Region 27	3.8	3.7	3.9	9.1	8.5	10.3	74.6	73.2	75.4	2.3	2.2	2.4	
12	Region 28	4.0	3.6	4.8	8.8	7.2	11.3	74.8	71.4	76.1	2.3	2.2	2.4	28	Region 28	3.9	3.4	4.3	8.5	7.4	9.6	74.8	73.5	76.1	2.3	2.2	2.5	
26	Region 29	3.8	3.4	4.2	8.8	7.3	9.8	75.3	74.0	76.7	2.2	2.1	2.3	26	Region 29	4.0	3.6	4.8	9.0	7.4	9.9	74.3	72.6	75.8	2.3	2.2	2.5	
23	Region 30	4.0	3.7	4.3	8.6	7.5	9.3	75.3	74.4	76.4	2.2	2.1	2.3	39	Region 30	4.1	3.7	5.1	8.9	8.1	9.6	74.3	72.4	75.4	2.3	2.2	2.5	
10	Region 31	4.0	3.8	4.8	8.7	8.0	9.5	75.0	74.5	75.9	2.3	2.2	2.4	39	Region 31	3.9	3.5	4.4	8.9	7.7	10.0	74.4	72.8	76.1	2.3	2.2	2.5	
13	Region 32	3.9	3.5	4.2	8.7	7.6	9.4	75.2	74.1	75.9	2.3	2.2	2.4	32	Region 32	4.2	3.6	4.8	9.2	7.7	10.1	73.7	71.9	76.5	2.4	2.2	2.6	
40	Region 33	4.1	3.8	4.5	8.4	7.8	9.1	75.2	73.8	76.5	2.3	2.2	2.5	20	Region 33	3.9	3.3	4.3	8.5	7.4	9.5	74.7	73.7	75.7	2.4	2.2	2.5	
21	Region 34	3.9	3.5	4.6	8.4	7.1	8.9	75.5	74.7	75.9	2.2	2.1	2.4	16	Region 34	3.9	3.6	4.7	8.9	8.4	10.0	74.8	72.5	75.9	2.3	2.1	2.5	
6	Region 35	4.3	3.9	4.8	8.9	8.4	9.4	75.0	74.3	75.7	2.3	2.2	2.3	6	Region 35	4.0	3.7	4.3	8.7	8.0	9.7	74.8	73.6	75.9	2.4	2.3	2.4	
17	Region 36	4.0	3.9	4.4	8.7	8.0	9.2	75.3	74.4	76.2	2.3	2.2	2.3	17	Region 36	3.9	3.5	4.6	8.5	7.6	9.4	74.9	73.2	76.4	2.3	2.3	2.5	
<b>388</b>	<b>Ave. WM1</b>	<b>4.0</b>	<b>3.4</b>	<b>4.8</b>	<b>8.3</b>	<b>6.1</b>	<b>11.3</b>	<b>75.6</b>	<b>71.4</b>	<b>76.8</b>	<b>2.3</b>	<b>2.1</b>	<b>2.5</b>	<b>343</b>	<b>Ave. YM1</b>	<b>4.0</b>	<b>3.2</b>	<b>5.1</b>	<b>8.7</b>	<b>7.3</b>	<b>10.3</b>	<b>74.7</b>	<b>71.9</b>	<b>76.5</b>	<b>2.3</b>	<b>2.0</b>	<b>2.6</b>	
	<b>Min. WM1</b>													<b>Min. YM1</b>														
	<b>Max. WM1</b>													<b>Max. YM1</b>														

TABLE 19: NUTRITIONAL VALUES OF WHITE MAIZE ACCORDING TO GRADE (2020/21) ( continue)												TABLE 20: NUTRITIONAL VALUES OF YELLOW MAIZE ACCORDING TO GRADE (2020/21) ( continue)																
Number of samples	Region	Fat % (db)			Protein % (db)			Starch % (db)			Crude Fibre % (db)			Number of samples	Region	Fat % (db)			Protein % (db)			Starch % (db)			Crude Fibre % (db)			
		ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.			ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	
GRADE: WM2												GRADE: YM2																
-	Region 10	-	-	-	-	-	-	-	-	-	-	-	-	3	Region 10	3.8	3.4	4.1	8.7	8.3	9.0	74.9	74.7	75.1	2.3	2.2	2.4	
1	Region 11	3.8	3.8	3.8	9.8	9.8	75.0	75.0	75.0	75.0	75.0	75.0	2.2	2.2	2	Region 11	3.7	3.6	3.8	8.4	7.6	9.2	75.5	74.6	76.4	2.2	2.2	2.2
3	Region 12	4.1	4.0	4.3	8.2	7.4	8.7	75.5	74.7	76.1	75.5	76.1	2.3	2.3	1	Region 12	4.1	4.1	4.1	8.4	8.4	8.4	75.4	75.4	75.4	2.2	2.2	2.2
4	Region 13	4.0	3.9	4.2	8.0	7.7	8.4	76.1	75.4	77.3	76.1	75.4	2.2	2.2	1	Region 13	4.2	4.2	4.2	7.9	7.9	7.9	75.8	75.8	75.8	2.4	2.4	2.4
9	Region 14	4.0	3.5	4.5	7.9	7.5	8.2	76.0	75.0	76.7	76.0	75.0	2.3	2.3	1	Region 14	3.9	3.9	3.9	8.1	8.1	8.1	75.3	75.3	75.3	2.2	2.2	2.2
-	Region 15	-	-	-	-	-	-	-	-	-	-	-	-	-	-	Region 15	-	-	-	-	-	-	-	-	-	-	-	
1	Region 16	4.2	4.2	4.2	8.1	8.1	8.1	76.3	76.3	76.3	76.3	76.3	2.2	2.2	1	Region 16	4.1	4.1	4.1	7.9	7.9	7.9	75.3	75.3	75.3	2.4	2.4	2.4
2	Region 17	4.0	4.0	4.1	8.1	7.7	8.4	76.1	75.9	76.3	76.1	75.9	2.2	2.2	-	Region 17	-	-	-	-	-	-	-	-	-	-	-	
3	Region 18	4.0	3.9	4.1	8.0	7.8	8.1	75.8	75.4	76.1	75.8	75.4	2.2	2.2	1	Region 18	4.1	4.1	4.1	8.1	8.1	8.1	76.0	76.0	76.0	2.3	2.3	2.3
5	Region 19	4.0	3.8	4.2	8.0	7.1	8.8	75.7	74.9	76.2	75.7	74.9	2.3	2.3	2	Region 19	4.2	4.2	4.3	9.1	9.0	9.1	74.7	74.4	75.1	2.3	2.3	2.3
1	Region 20	3.9	3.9	3.9	7.4	7.4	7.4	75.8	75.8	75.8	75.8	75.8	2.3	2.3	-	Region 20	-	-	-	-	-	-	-	-	-	-	-	
3	Region 21	4.0	3.7	4.3	8.5	7.7	9.2	75.3	74.3	75.9	75.3	74.3	2.4	2.3	1	Region 21	3.9	3.9	3.9	7.9	7.9	7.9	76.0	76.0	76.0	2.3	2.3	2.3
2	Region 22	3.8	3.8	3.9	7.7	7.7	7.7	76.1	75.7	76.5	76.1	75.7	2.4	2.4	1	Region 22	3.9	3.9	3.9	8.4	8.4	8.4	75.4	75.4	75.4	2.3	2.3	2.3
6	Region 23	4.1	3.6	4.5	8.1	7.6	8.8	75.8	74.8	76.3	75.8	74.8	2.3	2.2	-	Region 23	-	-	-	-	-	-	-	-	-	-	-	
2	Region 24	4.4	3.8	5.0	7.6	7.3	8.0	75.4	75.1	75.7	75.4	75.1	2.4	2.3	1	Region 24	3.8	3.8	3.8	7.6	7.6	7.6	74.4	74.4	74.4	2.5	2.5	2.5
-	Region 25	-	-	-	-	-	-	-	-	-	-	-	-	-	5	Region 25	3.9	3.7	4.2	8.1	7.7	8.5	74.8	73.8	75.7	2.3	2.2	2.4
2	Region 26	4.0	3.8	4.1	7.8	7.6	8.0	75.9	75.7	76.2	75.9	75.7	2.2	2.2	3	Region 26	3.9	3.7	4.1	8.3	8.0	8.8	74.9	74.1	75.7	2.4	2.3	2.4
2	Region 27	4.0	3.9	4.1	7.3	7.2	7.4	76.7	76.4	77.0	76.7	76.4	2.2	2.2	4	Region 27	3.8	3.7	4.0	8.6	8.2	8.8	75.3	75.2	75.5	2.3	2.2	2.3
-	Region 28	-	-	-	-	-	-	-	-	-	-	-	-	-	5	Region 28	3.8	3.6	4.0	8.7	8.0	9.9	74.7	73.4	75.7	2.3	2.3	2.4
1	Region 29	4.0	4.0	4.0	9.5	9.5	9.5	73.7	73.7	73.7	73.7	73.7	2.4	2.4	-	Region 29	-	-	-	-	-	-	-	-	-	-	-	
7	Region 30	3.9	3.6	4.6	8.8	8.4	9.2	75.2	73.6	76.3	75.2	73.6	2.3	2.1	2	Region 30	4.5	4.0	5.1	8.6	8.2	8.9	74.6	74.6	74.6	2.3	2.3	2.4
7	Region 31	3.9	3.7	4.0	8.7	8.2	9.1	75.1	74.2	75.9	75.1	74.2	2.3	2.2	1	Region 31	3.9	3.9	3.9	7.8	7.8	7.8	75.1	75.1	75.1	2.5	2.5	2.5
8	Region 32	3.9	3.7	4.1	8.8	8.0	9.4	74.9	73.7	76.2	74.9	73.7	2.3	2.2	3	Region 32	4.1	3.6	4.6	9.1	8.8	9.8	74.2	73.1	75.4	2.4	2.3	2.5
10	Region 33	4.0	3.7	4.2	8.6	7.9	9.2	75.3	74.3	76.2	75.3	74.3	2.3	2.2	2	Region 33	3.7	3.4	3.9	8.9	8.5	9.3	74.5	74.5	74.5	2.3	2.2	2.4
4	Region 34	3.8	3.6	4.1	8.6	7.6	9.5	75.6	74.6	76.1	75.6	74.6	2.2	2.2	2	Region 34	4.0	3.9	4.0	8.6	8.6	8.6	74.7	74.4	74.9	2.3	2.2	2.4
1	Region 35	3.9	3.9	3.9	8.1	8.1	8.1	75.8	75.8	75.8	75.8	75.8	2.2	2.2	-	Region 35	-	-	-	-	-	-	-	-	-	-	-	
6	Region 36	4.0	3.7	4.2	8.7	8.0	9.3	75.0	74.4	75.6	75.0	74.4	2.3	2.2	1	Region 36	4.0	4.0	4.0	8.9	8.9	8.9	74.8	74.8	74.8	2.3	2.3	2.3
90	Ave. WM2	4.0	3.5	5.0	8.3	7.1	9.8	75.5	73.6	77.3	75.5	73.6	2.3	2.1	43	Ave. YM2	3.9	3.4	5.1	8.5	7.6	9.9	74.9	73.1	76.4	2.3	2.2	2.5
	Min. WM2															Min. YM2												
	Max. WM2															Max. YM2												

TABLE 19: NUTRITIONAL VALUES OF WHITE MAIZE ACCORDING TO GRADE (2020/21) ( continue)												TABLE 20: NUTRITIONAL VALUES OF YELLOW MAIZE ACCORDING TO GRADE (2020/21) ( continue)															
Number of samples	Region	Fat % (db)			Protein % (db)			Starch % (db)			Crude Fibre % (db)			Number of samples	Region	Fat % (db)			Protein % (db)			Starch % (db)			Crude Fibre % (db)		
		ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.			ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.
<b>GRADE: WM3</b>												<b>GRADE: YM3</b>															
-	Region 10	-	-	-	-	-	-	-	-	-	-	-	-	1	Region 10	3.8	3.8	3.8	7.5	7.5	7.5	75.9	75.9	75.9	2.2	2.2	2.2
-	Region 11	-	-	-	-	-	-	-	-	-	-	-	-	-	Region 11	-	-	-	-	-	-	-	-	-	-	-	
-	Region 12	-	-	-	-	-	-	-	-	-	-	-	-	-	Region 12	-	-	-	-	-	-	-	-	-	-	-	
1	Region 13	4.1	4.1	4.1	7.4	7.4	7.4	75.7	75.7	75.7	2.3	2.3	2.3	-	Region 13	-	-	-	-	-	-	-	-	-	-	-	
3	Region 14	3.8	3.6	3.9	7.5	7.4	7.6	76.4	75.8	76.8	2.2	2.2	2.3	-	Region 14	-	-	-	-	-	-	-	-	-	-	-	
-	Region 15	-	-	-	-	-	-	-	-	-	-	-	-	-	Region 15	-	-	-	-	-	-	-	-	-	-	-	
1	Region 16	3.7	3.7	3.7	8.1	8.1	8.1	75.8	75.8	75.8	2.3	2.3	2.3	-	Region 16	-	-	-	-	-	-	-	-	-	-	-	
8	Region 17	4.0	3.7	4.3	8.1	7.4	8.8	75.5	74.4	76.6	2.3	2.2	2.4	-	Region 17	-	-	-	-	-	-	-	-	-	-	-	
1	Region 18	3.7	3.7	3.7	9.0	9.0	9.0	75.5	75.5	75.5	2.3	2.3	2.3	-	Region 18	-	-	-	-	-	-	-	-	-	-	-	
5	Region 19	3.9	3.8	4.2	7.9	7.3	8.5	76.1	75.2	76.9	2.3	2.2	2.3	-	Region 19	-	-	-	-	-	-	-	-	-	-	-	
1	Region 20	3.8	3.8	3.8	8.4	8.4	8.4	75.3	75.3	75.3	2.2	2.2	2.2	-	Region 20	-	-	-	-	-	-	-	-	-	-	-	
3	Region 21	3.9	3.7	4.1	8.3	7.9	8.5	75.5	75.4	75.6	2.3	2.2	2.4	1	Region 21	4.1	4.1	4.1	9.6	9.6	9.6	73.0	73.0	73.0	2.2	2.2	2.2
-	Region 22	-	-	-	-	-	-	-	-	-	-	-	-	-	Region 22	-	-	-	-	-	-	-	-	-	-	-	
5	Region 23	3.9	3.8	4.0	8.0	7.8	8.3	75.5	75.3	75.7	2.3	2.3	2.4	-	Region 23	-	-	-	-	-	-	-	-	-	-	-	
-	Region 24	-	-	-	-	-	-	-	-	-	-	-	-	1	Region 24	3.5	3.5	3.5	8.0	8.0	8.0	75.0	75.0	75.0	2.5	2.5	2.5
-	Region 25	-	-	-	-	-	-	-	-	-	-	-	-	-	Region 25	-	-	-	-	-	-	-	-	-	-	-	
1	Region 26	4.1	4.1	4.1	7.8	7.8	7.8	75.7	75.7	75.7	2.3	2.3	2.3	-	Region 26	-	-	-	-	-	-	-	-	-	-	-	
-	Region 27	-	-	-	-	-	-	-	-	-	-	-	-	-	Region 27	-	-	-	-	-	-	-	-	-	-	-	
-	Region 28	-	-	-	-	-	-	-	-	-	-	-	-	-	Region 28	-	-	-	-	-	-	-	-	-	-	-	
-	Region 29	-	-	-	-	-	-	-	-	-	-	-	-	1	Region 29	3.9	3.9	3.9	8.6	8.6	8.6	74.9	74.9	74.9	2.3	2.3	2.3
2	Region 30	3.7	3.6	3.9	8.2	8.0	8.4	75.4	74.5	76.2	2.2	2.1	2.3	1	Region 30	3.9	3.9	3.9	9.5	9.5	9.5	74.2	74.2	74.2	2.3	2.3	2.3
2	Region 31	3.9	3.9	4.0	9.0	8.7	9.3	74.6	74.1	75.1	2.3	2.3	2.4	-	Region 31	-	-	-	-	-	-	-	-	-	-	-	
1	Region 32	3.7	3.7	3.7	7.8	7.8	7.8	75.7	75.7	75.7	2.3	2.3	2.3	-	Region 32	-	-	-	-	-	-	-	-	-	-	-	
1	Region 33	4.1	4.1	4.1	8.1	8.1	8.1	75.8	75.8	75.8	2.3	2.3	2.3	2	Region 33	4.0	3.8	4.2	8.8	8.5	9.2	74.4	73.4	75.4	2.5	2.4	2.6
1	Region 34	4.1	4.1	4.1	8.3	8.3	8.3	76.2	76.2	76.2	2.2	2.2	2.2	1	Region 34	3.9	3.9	3.9	10.1	10.1	10.1	73.3	73.3	73.3	2.3	2.3	2.3
-	Region 35	-	-	-	-	-	-	-	-	-	-	-	-	-	Region 35	-	-	-	-	-	-	-	-	-	-	-	
-	Region 36	-	-	-	-	-	-	-	-	-	-	-	-	-	Region 36	-	-	-	-	-	-	-	-	-	-	-	
36	Ave. WM3	3.9	3.6	4.3	8.1	7.3	9.2	75.6	74.1	76.9	2.3	2.1	2.4	8	Ave. YM3	3.9	3.5	4.2	8.9	7.5	10.1	74.4	73.0	75.9	2.3	2.2	2.6
-	Min. WM3	-	-	-	-	-	-	-	-	-	-	-	-	-	Min. YM3	-	-	-	-	-	-	-	-	-	-	-	
-	Max. WM3	-	-	-	-	-	-	-	-	-	-	-	-	-	Max. YM3	-	-	-	-	-	-	-	-	-	-	-	



TABLE 19: NUTRITIONAL VALUES OF WHITE MAIZE ACCORDING TO GRADE (2020/21) ( continue)												TABLE 20: NUTRITIONAL VALUES OF YELLOW MAIZE ACCORDING TO GRADE (2020/21) ( continue)															
Number of samples	Region	Fat % (db)			Protein % (db)			Starch % (db)			Crude Fibre % (db)			Number of samples	Region	Fat % (db)			Protein % (db)			Starch % (db)			Crude Fibre % (db)		
		ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.			ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.
<b>CLASS: COM</b>																											
1	Region 11	3.6	3.6	3.6	8.9	8.9	8.9	75.7	75.7	75.7	2.2	2.2	2.2	-	Region 11	4.0	4.0	4.0	8.1	8.1	8.1	75.8	75.8	75.8	2.3	2.3	2.3
3	Region 12	4.0	3.8	4.1	8.6	8.3	8.8	75.4	75.0	75.7	2.3	2.3	2.3	1	Region 12	-	-	-	-	-	-	-	-	-	-	-	-
1	Region 13	4.2	4.2	4.2	8.1	8.1	8.1	75.7	75.7	75.7	2.2	2.2	2.2	-	Region 13	3.9	3.9	3.9	8.0	8.0	8.0	76.1	76.1	76.1	2.3	2.3	2.3
3	Region 14	3.8	3.6	4.0	7.8	7.4	8.1	76.6	76.4	76.8	2.2	2.2	2.3	1	Region 14	4.0	4.0	4.0	8.0	8.0	8.0	76.0	76.0	76.0	2.3	2.3	2.3
4	Region 17	4.0	3.9	4.2	7.8	6.8	8.3	75.7	75.5	76.0	2.3	2.3	2.3	1	Region 17	3.9	3.8	4.1	8.4	7.9	8.9	75.5	74.9	76.1	2.4	2.3	2.4
1	Region 18	4.3	4.3	4.3	8.3	8.3	8.3	75.3	75.3	75.3	2.2	2.2	2.2	2	Region 18	-	-	-	-	-	-	-	-	-	-	-	-
2	Region 19	3.7	3.6	3.8	8.2	7.7	8.6	76.2	76.2	76.2	2.2	2.1	2.3	-	Region 19	4.0	3.8	4.2	9.0	9.0	9.1	74.9	74.6	75.2	2.4	2.4	2.4
3	Region 21	3.9	3.7	4.2	8.9	8.1	10.0	75.1	74.2	75.7	2.3	2.2	2.3	2	Region 21	3.8	3.8	3.8	8.1	8.1	8.1	75.6	75.6	75.6	2.3	2.3	2.3
2	Region 22	4.0	3.8	4.1	7.5	7.4	7.6	76.1	76.0	76.2	2.3	2.3	2.4	1	Region 22	-	-	-	-	-	-	-	-	-	-	-	-
5	Region 23	4.0	3.8	4.2	8.1	7.8	8.4	75.6	75.3	75.9	2.3	2.2	2.4	-	Region 23	3.6	3.6	3.6	9.3	9.3	9.3	73.7	73.7	73.7	2.5	2.5	2.5
1	Region 24	3.9	3.9	3.9	7.7	7.7	7.7	76.0	76.0	76.0	2.3	2.3	2.3	6	Region 24	3.9	3.7	4.1	8.4	7.1	9.7	74.3	73.0	75.6	2.4	2.3	2.5
1	Region 25	5.1	5.1	5.1	6.0	6.0	6.0	76.2	76.2	76.2	2.4	2.4	2.4	2	Region 25	4.0	3.7	4.3	9.7	9.6	9.7	73.9	73.8	74.1	2.4	2.4	2.5
-	Region 26	-	-	-	-	-	-	-	-	-	-	-	-	1	Region 26	3.5	3.5	3.5	8.5	8.5	8.5	75.7	75.7	75.7	2.4	2.4	2.4
-	Region 27	-	-	-	-	-	-	-	-	-	-	-	-	4	Region 27	4.0	3.7	4.3	9.3	8.9	10.0	74.0	73.2	75.0	2.4	2.3	2.4
1	Region 28	4.0	4.0	4.0	7.9	7.9	7.9	75.0	75.0	75.0	2.3	2.3	2.3	3	Region 28	3.9	3.8	4.2	8.9	8.7	9.2	74.5	74.0	74.9	2.4	2.3	2.4
-	Region 29	-	-	-	-	-	-	-	-	-	-	-	-	4	Region 29	4.1	3.9	4.6	9.0	8.2	9.3	74.1	73.4	74.7	2.4	2.4	2.4
2	Region 30	3.9	3.7	4.1	8.2	7.9	8.5	75.5	74.6	76.4	2.3	2.3	2.4	5	Region 30	4.0	3.8	4.2	8.9	8.5	9.6	74.2	73.2	74.7	2.4	2.3	2.6
1	Region 31	3.7	3.7	3.7	8.7	8.7	8.7	74.3	74.3	74.3	2.5	2.5	2.5	5	Region 31	4.2	3.7	4.7	9.0	8.6	9.5	74.1	73.3	74.6	2.4	2.4	2.6
4	Region 32	4.1	3.8	4.6	8.6	8.3	9.0	75.1	75.1	75.2	2.4	2.3	2.4	5	Region 32	4.0	3.9	4.2	9.1	8.6	9.6	74.2	73.4	75.0	2.4	2.3	2.5
3	Region 33	3.9	3.7	4.1	9.4	9.1	9.5	74.4	74.2	74.8	2.3	2.2	2.4	2	Region 33	3.5	3.2	3.9	8.6	8.0	9.3	75.4	74.3	76.5	2.3	2.2	2.3
4	Region 34	4.1	3.7	4.6	8.6	7.9	9.1	75.0	74.7	75.4	2.3	2.2	2.4	46	Region 34	4.0	4.0	4.0	8.8	7.1	10.0	74.5	73.0	76.5	2.4	2.2	2.6
4	Region 36	4.2	4.1	4.7	7.8	7.0	8.6	75.4	74.8	75.7	2.3	2.2	2.4	440	Region 36	4.0	3.2	4.7	8.7	7.1	10.3	74.7	71.9	76.5	2.3	2.0	2.6
46	Ave. COM	4.0	3.6	5.1	8.2	6.0	10.0	75.5	74.2	76.8	2.3	2.1	2.5	1000	Ave. COM	4.0	3.2	5.1	8.5	6.0	11.3	75.2	71.4	77.3	2.3	2.0	2.6
560	Ave. White	4.0	3.4	5.1	8.3	6.0	11.3	75.5	71.4	77.3	2.3	2.1	2.5	560	Min. COM	4.0	3.2	5.1	8.5	6.0	11.3	75.2	71.4	77.3	2.3	2.0	2.6
1000	Ave. Maize	4.0	3.2	5.1	8.5	6.0	11.3	75.2	71.4	77.3	2.3	2.0	2.6	1000	Max. COM	4.0	3.2	5.1	8.5	6.0	11.3	75.2	71.4	77.3	2.3	2.0	2.6
	Min. White														Min. Yellow												
	Max. White														Max. Yellow												
	Min. Maize														Ave. Maize												
	Max. Maize														Min. Maize												
															Max. Maize												

**TABLE 21: NUTRITIONAL VALUES OF WHITE AND YELLOW MAIZE (2020/21)**

Number of samples	Region	Fat % (db)			Protein % (db)			Starch % (db)			Crude Fibre % (db)		
		ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.
		<b>WHITE</b>											
2	Region 11	3.7	3.6	3.8	9.3	8.9	9.8	75.3	75.0	75.7	2.2	2.2	2.2
14	Region 12	4.1	3.8	4.8	8.3	7.4	8.8	75.4	74.7	76.1	2.3	2.3	2.4
29	Region 13	4.1	3.6	4.3	8.0	7.2	8.9	75.7	74.5	77.3	2.3	2.2	2.4
40	Region 14	3.9	3.4	4.5	7.8	7.1	8.6	76.1	75.0	76.8	2.3	2.2	2.4
8	Region 16	4.0	3.7	4.3	7.6	6.9	8.1	76.2	75.4	76.8	2.2	2.2	2.3
29	Region 17	4.0	3.6	4.3	8.0	6.8	8.8	75.8	74.4	76.6	2.2	2.2	2.4
29	Region 18	4.0	3.4	4.7	8.3	7.5	10.1	75.6	74.1	76.3	2.2	2.2	2.3
29	Region 19	4.0	3.5	4.3	8.2	7.1	8.9	75.7	74.9	76.9	2.3	2.1	2.4
17	Region 20	4.0	3.5	4.3	8.1	6.8	8.8	75.5	74.7	76.2	2.2	2.1	2.3
36	Region 21	4.0	3.6	4.4	8.2	7.6	10.0	75.6	74.2	76.3	2.3	2.2	2.4
13	Region 22	3.9	3.6	4.2	7.8	7.1	8.8	76.0	75.7	76.5	2.3	2.3	2.5
47	Region 23	4.0	3.6	4.5	7.9	7.1	8.8	75.9	74.8	76.5	2.3	2.2	2.4
13	Region 24	4.1	3.8	5.0	7.8	6.1	8.5	75.8	75.1	76.4	2.3	2.2	2.4
3	Region 25	4.5	4.0	5.1	7.0	6.0	7.9	75.7	75.3	76.2	2.3	2.2	2.4
8	Region 26	4.0	3.8	4.2	7.6	7.3	8.0	76.0	75.7	76.2	2.3	2.2	2.4
5	Region 27	4.1	3.8	4.4	7.3	7.0	7.9	76.2	75.6	77.0	2.2	2.2	2.3
13	Region 28	4.0	3.6	4.8	8.7	7.2	11.3	74.8	71.4	76.1	2.3	2.2	2.4
27	Region 29	3.8	3.4	4.2	8.8	7.3	9.8	75.3	73.7	76.7	2.2	2.1	2.4
34	Region 30	3.9	3.6	4.6	8.6	7.5	9.3	75.3	73.6	76.4	2.2	2.1	2.4
20	Region 31	3.9	3.7	4.8	8.7	8.0	9.5	75.0	74.1	75.9	2.3	2.2	2.5
26	Region 32	3.9	3.5	4.6	8.6	7.6	9.4	75.1	73.7	76.2	2.3	2.2	2.4
54	Region 33	4.1	3.7	4.5	8.5	7.8	9.5	75.2	73.8	76.5	2.3	2.2	2.5
30	Region 34	3.9	3.5	4.6	8.4	7.1	9.5	75.5	74.6	76.2	2.2	2.1	2.4
7	Region 35	4.3	3.9	4.8	8.8	8.1	9.4	75.1	74.3	75.8	2.3	2.2	2.3
27	Region 36	4.1	3.7	4.7	8.6	7.0	9.3	75.2	74.4	76.2	2.3	2.2	2.4
<b>560</b>	<b>Ave. white</b>	<b>4.0</b>			<b>8.3</b>			<b>75.5</b>			<b>2.3</b>		
	<b>Min. white</b>	<b>3.4</b>			<b>6.0</b>			<b>71.4</b>			<b>2.1</b>		
	<b>Max. white</b>	<b>5.1</b>			<b>11.3</b>			<b>77.3</b>			<b>2.5</b>		
<b>YELLOW</b>													
28	Region 10	3.8	3.2	4.2	8.5	7.5	9.1	75.3	74.1	76.4	2.2	2.0	2.4
15	Region 11	3.8	3.5	4.2	8.4	7.6	9.2	75.4	74.5	76.4	2.2	2.1	2.3
2	Region 12	4.0	4.0	4.1	8.2	8.1	8.4	75.6	75.4	75.8	2.2	2.2	2.3
6	Region 13	4.0	3.6	4.5	8.4	7.9	8.8	75.8	75.5	76.2	2.2	2.2	2.4
8	Region 14	3.9	3.7	4.0	8.5	8.0	8.8	75.3	74.6	76.1	2.3	2.2	2.3
1	Region 16	4.1	4.1	4.1	7.9	7.9	7.9	75.3	75.3	75.3	2.4	2.4	2.4
4	Region 17	4.0	3.9	4.1	8.6	7.7	9.7	75.3	73.7	76.3	2.3	2.2	2.3
9	Region 18	4.0	3.6	4.2	8.6	7.9	9.2	75.2	74.5	76.1	2.3	2.2	2.4
9	Region 19	4.0	3.8	4.3	9.2	8.9	9.5	74.8	74.2	75.4	2.3	2.2	2.4
7	Region 20	3.9	3.7	4.5	8.4	7.9	8.8	75.2	74.2	75.9	2.3	2.1	2.4
7	Region 21	4.0	3.8	4.2	8.9	7.9	9.6	74.7	73.0	76.0	2.3	2.2	2.4
7	Region 22	4.2	3.8	4.5	8.5	8.1	9.3	75.0	74.3	75.6	2.3	2.3	2.4
4	Region 23	4.0	3.6	4.4	8.5	8.2	8.9	75.3	75.0	75.5	2.3	2.2	2.4
6	Region 24	3.8	3.5	4.0	8.1	7.3	8.8	75.2	74.4	76.2	2.3	2.2	2.5
24	Region 25	3.9	3.6	4.4	8.2	7.3	9.6	74.8	73.6	76.3	2.3	2.2	2.5


**TABLE 21: NUTRITIONAL VALUES OF WHITE AND YELLOW MAIZE (2020/21)**  
**(continue)**

Number of samples	Region	Fat % (db)			Protein % (db)			Starch % (db)			Crude Fibre % (db)		
		ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.
<b>YELLOW</b>													
20	Region 26	4.0	3.6	4.7	8.5	7.1	9.7	74.7	73.0	75.7	2.3	2.2	2.5
10	Region 27	3.8	3.7	4.3	9.0	8.2	10.3	74.8	73.2	75.5	2.3	2.2	2.5
34	Region 28	3.9	3.4	4.3	8.5	7.4	9.9	74.8	73.4	76.1	2.3	2.2	2.5
31	Region 29	4.0	3.6	4.8	9.0	7.4	10.0	74.3	72.6	75.8	2.3	2.2	2.5
45	Region 30	4.1	3.7	5.1	8.9	8.1	9.6	74.3	72.4	75.4	2.3	2.2	2.5
44	Region 31	4.0	3.5	4.6	8.9	7.7	10.0	74.4	72.8	76.1	2.3	2.2	2.5
40	Region 32	4.2	3.6	4.8	9.2	7.7	10.1	73.8	71.9	76.5	2.4	2.2	2.6
29	Region 33	3.9	3.3	4.7	8.6	7.4	9.5	74.6	73.3	75.7	2.4	2.2	2.6
24	Region 34	3.9	3.6	4.7	9.0	8.4	10.1	74.6	72.5	75.9	2.3	2.1	2.5
6	Region 35	4.0	3.7	4.3	8.7	8.0	9.7	74.8	73.6	75.9	2.4	2.3	2.4
20	Region 36	3.9	3.2	4.6	8.6	7.6	9.4	74.9	73.2	76.5	2.3	2.2	2.5
<b>440</b>	<b>Ave. yellow</b>	<b>4.0</b>			<b>8.7</b>			<b>74.7</b>			<b>2.3</b>		
	<b>Min. yellow</b>		<b>3.2</b>			<b>7.1</b>			<b>71.9</b>			<b>2.0</b>	
	<b>Max. yellow</b>			<b>5.1</b>			<b>10.3</b>			<b>76.5</b>			<b>2.6</b>
<b>WHITE AND YELLOW</b>													
28	Region 10	3.8	3.2	4.2	8.5	7.5	9.1	75.3	74.1	76.4	2.2	2.0	2.4
17	Region 11	3.8	3.5	4.2	8.5	7.6	9.8	75.4	74.5	76.4	2.2	2.1	2.3
16	Region 12	4.1	3.8	4.8	8.3	7.4	8.8	75.4	74.7	76.1	2.3	2.2	2.4
35	Region 13	4.1	3.6	4.5	8.1	7.2	8.9	75.7	74.5	77.3	2.3	2.2	2.4
48	Region 14	3.9	3.4	4.5	7.9	7.1	8.8	76.0	74.6	76.8	2.3	2.2	2.4
9	Region 16	4.0	3.7	4.3	7.7	6.9	8.1	76.1	75.3	76.8	2.3	2.2	2.4
33	Region 17	4.0	3.6	4.3	8.0	6.8	9.7	75.7	73.7	76.6	2.3	2.2	2.4
38	Region 18	4.0	3.4	4.7	8.4	7.5	10.1	75.5	74.1	76.3	2.3	2.2	2.4
38	Region 19	4.0	3.5	4.3	8.4	7.1	9.5	75.5	74.2	76.9	2.3	2.1	2.4
24	Region 20	4.0	3.5	4.5	8.2	6.8	8.8	75.4	74.2	76.2	2.2	2.1	2.4
43	Region 21	4.0	3.6	4.4	8.3	7.6	10.0	75.5	73.0	76.3	2.3	2.2	2.4
20	Region 22	4.0	3.6	4.5	8.0	7.1	9.3	75.7	74.3	76.5	2.3	2.3	2.5
51	Region 23	4.0	3.6	4.5	8.0	7.1	8.9	75.8	74.8	76.5	2.3	2.2	2.4
19	Region 24	4.0	3.5	5.0	7.9	6.1	8.8	75.7	74.4	76.4	2.3	2.2	2.5
27	Region 25	3.9	3.6	5.1	8.1	6.0	9.6	74.9	73.6	76.3	2.3	2.2	2.5
28	Region 26	4.0	3.6	4.7	8.3	7.1	9.7	75.1	73.0	76.2	2.3	2.2	2.5
15	Region 27	3.9	3.7	4.4	8.4	7.0	10.3	75.2	73.2	77.0	2.3	2.2	2.5
47	Region 28	3.9	3.4	4.8	8.6	7.2	11.3	74.8	71.4	76.1	2.3	2.2	2.5
58	Region 29	3.9	3.4	4.8	8.9	7.3	10.0	74.8	72.6	76.7	2.3	2.1	2.5
79	Region 30	4.1	3.6	5.1	8.8	7.5	9.6	74.7	72.4	76.4	2.3	2.1	2.5
64	Region 31	4.0	3.5	4.8	8.8	7.7	10.0	74.6	72.8	76.1	2.3	2.2	2.5
66	Region 32	4.1	3.5	4.8	9.0	7.6	10.1	74.3	71.9	76.5	2.4	2.2	2.6
83	Region 33	4.0	3.3	4.7	8.6	7.4	9.5	75.0	73.3	76.5	2.3	2.2	2.6
54	Region 34	3.9	3.5	4.7	8.7	7.1	10.1	75.1	72.5	76.2	2.3	2.1	2.5
13	Region 35	4.2	3.7	4.8	8.8	8.0	9.7	75.0	73.6	75.9	2.3	2.2	2.4
47	Region 36	4.0	3.2	4.7	8.6	7.0	9.4	75.1	73.2	76.5	2.3	2.2	2.5
<b>1000</b>	<b>Ave. W &amp; Y</b>	<b>4.0</b>			<b>8.5</b>			<b>75.2</b>			<b>2.3</b>		
	<b>Min. W &amp; Y</b>		<b>3.2</b>			<b>6.0</b>			<b>71.4</b>			<b>2.0</b>	
	<b>Max. W &amp; Y</b>			<b>5.1</b>			<b>11.3</b>			<b>77.3</b>			<b>2.6</b>


**TABLE 22: NUTRITIONAL VALUES OF SOUTH AFRICAN WHITE AND YELLOW MAIZE 2011/12 - 2020/21**

Season	Number of samples	Fat % (db)			Protein % (db)			Starch % (db)			Crude Fibre % (db)		
		ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.
<b>White Maize</b>													
2011/12	577	4.1	3.3	4.7	8.6	6.3	11.2	72.6	70.6	74.3	-	-	-
2012/13	508	4.1	3.3	5.3	9.2	6.4	11.5	71.4	68.5	73.6	-	-	-
2013/14	451	4.0	3.4	5.0	8.6	6.7	10.1	72.9	70.9	75.1	-	-	-
2014/15	485	4.2	3.3	5.8	9.4	6.3	11.2	72.6	69.8	74.9	-	-	-
2015/16	415	4.1	3.5	5.2	9.7	7.5	12.2	72.6	69.8	76.3	-	-	-
2016/17	549	4.2	3.4	5.3	8.7	6.8	11.5	74.1	69.8	75.9	-	-	-
2017/18	451	4.1	3.4	5.0	8.7	6.8	11.2	73.2	69.4	74.8	-	-	-
2018/19	404	4.0	3.0	5.3	9.0	7.4	12.3	73.6	69.7	75.4	1.9	1.7	2.8
2019/20	516	4.0	3.4	5.0	8.4	6.7	11.7	73.2	69.6	74.9	1.9	1.6	2.6
2020/21	560	4.0	3.4	5.1	8.3	6.0	11.3	75.5	71.4	77.3	2.3	2.1	2.5
<b>Weighted Average</b>		<b>4.1</b>			<b>8.8</b>			<b>73.2</b>			<b>2.1</b>		
<b>Minimum</b>			<b>3.0</b>			<b>6.0</b>			<b>68.5</b>			<b>1.6</b>	
<b>Maximum</b>				<b>5.8</b>		<b>12.3</b>			<b>77.3</b>				<b>2.8</b>
<b>Yellow Maize</b>													
2011/12	423	3.8	3.0	4.6	8.9	7.0	11.3	73.0	71.0	75.0	-	-	-
2012/13	492	3.9	2.9	4.7	9.2	7.1	12.8	71.9	69.4	73.9	-	-	-
2013/14	479	3.8	3.0	4.8	8.6	6.0	11.3	73.1	70.8	75.7	-	-	-
2014/15	515	4.0	3.1	5.1	9.5	7.1	11.9	72.9	70.8	75.2	-	-	-
2015/16	505	4.0	3.3	5.1	9.7	7.7	12.6	72.3	70.0	75.3	-	-	-
2016/17	451	4.1	3.3	5.1	8.9	7.1	10.8	73.7	71.3	76.2	-	-	-
2017/18	449	4.0	3.2	5.0	9.0	6.7	11.3	72.0	68.5	74.6	-	-	-
2018/19	404	4.0	3.3	5.0	9.2	6.7	11.4	72.7	70.2	75.2	2.0	1.6	2.3
2019/20	374	3.9	3.3	4.6	9.0	7.0	10.9	72.3	68.3	74.9	1.9	1.6	2.5
2020/21	440	4.0	3.2	5.1	8.7	7.1	10.3	74.7	71.9	76.5	2.3	2.0	2.6
<b>Weighted Average</b>		<b>4.0</b>			<b>9.1</b>			<b>72.9</b>			<b>2.1</b>		
<b>Minimum</b>			<b>2.9</b>			<b>6.0</b>			<b>68.3</b>			<b>1.6</b>	
<b>Maximum</b>				<b>5.1</b>		<b>12.8</b>			<b>76.5</b>				<b>2.6</b>
<b>White and Yellow Maize</b>													
2011/12	1000	4.0	3.0	4.7	8.7	6.3	11.3	72.8	70.6	75.0	-	-	-
2012/13	1000	4.0	2.9	5.3	9.2	6.4	12.8	71.6	68.5	73.9	-	-	-
2013/14	930	3.9	3.0	5.0	8.6	6.0	11.3	73.0	70.8	75.7	-	-	-
2014/15	1000	4.1	3.1	5.8	9.4	6.3	11.9	72.8	69.8	75.2	-	-	-
2015/16	920	4.1	3.3	5.2	9.7	7.5	12.6	72.4	69.8	76.3	-	-	-
2016/17	1000	4.1	3.3	5.3	8.8	6.8	11.5	73.9	69.8	76.2	-	-	-
2017/18	900	4.0	3.2	5.0	8.8	6.7	11.3	72.6	68.5	74.8	-	-	-
2018/19	808	4.0	3.0	5.3	9.1	6.7	12.3	73.2	69.7	75.4	1.9	1.6	2.8
2019/20	890	4.0	3.3	5.0	8.7	6.7	11.7	72.8	68.3	74.9	1.9	1.6	2.6
2020/21	1000	4.0	3.2	5.1	8.5	6.0	11.3	75.2	71.4	77.3	2.3	2.0	2.6
<b>Weighted Average</b>		<b>4.0</b>			<b>8.9</b>			<b>73.0</b>			<b>2.0</b>		
<b>Minimum</b>			<b>2.9</b>			<b>6.0</b>			<b>68.3</b>			<b>1.6</b>	
<b>Maximum</b>				<b>5.8</b>		<b>12.8</b>			<b>77.3</b>				<b>2.8</b>


GRAPH 47: FAT CONTENT OF WHITE AND YELLOW MAIZE OVER 10 SEASONS



GRAPH 48: PROTEIN CONTENT OF WHITE AND YELLOW MAIZE OVER 10 SEASONS



GRAPH 49: STARCH CONTENT OF WHITE AND YELLOW MAIZE OVER 10 SEASONS



**TABLE 23: PRESENCE OF GENETICALLY MODIFIED MAIZE (2020/21)**

REGION	W/Y	Cry1Ab % (LOD: 0.4%)	Cry2Ab % (LOD: 0.5%)	CP4 EPSPS % (LOD: 0.25%)	REGION	W/Y	Cry1Ab % (LOD: 0.4%)	Cry2Ab % (LOD: 0.5%)	CP4 EPSPS % (LOD: 0.25%)	REGION	W/Y	Cry1Ab % (LOD: 0.4%)	Cry2Ab % (LOD: 0.5%)	CP4 EPSPS % (LOD: 0.25%)
10	Y	>5.0	>5.0	>5.0	23	W	>5.0	>5.0	>5.0	30	W	>5.0	>5.0	>5.0
11	Y	>5.0	>5.0	>5.0	23	W	>5.0	>5.0	>5.0	31	Y	>5.0	>5.0	>5.0
12	W	>5.0	>5.0	>5.0	23	Y	>5.0	>5.0	>5.0	31	W	>5.0	>5.0	>5.0
13	Y	2.4	>5.0	>5.0	23	W	>5.0	>5.0	>5.0	31	Y	>5.0	>5.0	>5.0
13	Y	1.00	1.6	>5.0	24	Y	>5.0	>5.0	>5.0	31	W	<0.4	<0.5	<0.25
13	W	>5.0	>5.0	>5.0	25	Y	>5.0	>5.0	>5.0	32	W	>5.0	>5.0	>5.0
13	W	>5.0	>5.0	>5.0	25	W	>5.0	>5.0	>5.0	32	W	<0.4	0.40	0.51
14	Y	>5.0	>5.0	>5.0	25	Y	>5.0	>5.0	>5.0	32	Y	>5.0	>5.0	>5.0
14	W	>5.0	>5.0	>5.0	26	W	>5.0	>5.0	>5.0	32	W	>5.0	>5.0	>5.0
14	Y	>5.0	>5.0	>5.0	26	Y	>5.0	>5.0	>5.0	32	W	<0.4	<0.5	<0.25
14	W	>5.0	>5.0	>5.0	26	Y	>5.0	>5.0	>5.0	32	Y	<0.4	0.51	0.6
14	W	>5.0	>5.0	>5.0	27	W	>5.0	>5.0	>5.0	32	Y	>5.0	>5.0	>5.0
16	W	>5.0	>5.0	>5.0	27	Y	>5.0	>5.0	>5.0	33	W	>5.0	>5.0	>5.0
17	W	>5.0	>5.0	>5.0	28	Y	>5.0	>5.0	>5.0	33	Y	>5.0	>5.0	>5.0
17	Y	>5.0	>5.0	>5.0	28	W	2.9	>5.0	>5.0	33	Y	>5.0	>5.0	>5.0
17	W	>5.0	>5.0	>5.0	28	W	>5.0	>5.0	1.7	33	Y	3.9	<0.5	>5.0
18	Y	>5.0	>5.0	>5.0	28	W	>5.0	>5.0	>5.0	33	Y	2.6	>5.0	>5.0
18	W	>5.0	>5.0	>5.0	28	Y	>5.0	>5.0	>5.0	33	W	>5.0	>5.0	>5.0
18	Y	>5.0	>5.0	>5.0	29	Y	>5.0	>5.0	>5.0	33	W	>5.0	>5.0	>5.0
18	W	>5.0	>5.0	>5.0	29	Y	>5.0	>5.0	>5.0	33	W	>5.0	>5.0	>5.0
19	Y	>5.0	>5.0	>5.0	29	W	<0.4	0.83	0.34	34	Y	0.43	0.94	3.7
19	W	>5.0	>5.0	>5.0	29	Y	3.3	>5.0	>5.0	34	Y	>5.0	>5.0	>5.0
19	Y	>5.0	>5.0	>5.0	29	W	>5.0	>5.0	>5.0	34	W	>5.0	>5.0	>5.0
19	W	>5.0	>5.0	>5.0	29	W	>5.0	>5.0	>5.0	34	W	<0.4	0.29	0.34
20	Y	>5.0	>5.0	>5.0	29	Y	<0.4	0.22	0.46	34	W	>5.0	>5.0	>5.0
20	W	>5.0	>5.0	>5.0	29	W	>5.0	>5.0	>5.0	35	Y	>5.0	>5.0	>5.0
20	W	>5.0	>5.0	>5.0	29	W	>5.0	>5.0	>5.0	35	W	>5.0	>5.0	>5.0
21	Y	>5.0	>5.0	>5.0	30	Y	>5.0	>5.0	>5.0	36	W	>5.0	>5.0	>5.0
21	W	>5.0	>5.0	>5.0	30	Y	>5.0	>5.0	>5.0	36	W	>5.0	>5.0	>5.0
21	Y	>5.0	>5.0	>5.0	30	Y	>5.0	>5.0	>5.0	36	Y	>5.0	>5.0	>5.0
21	W	>5.0	>5.0	>5.0	30	W	<0.4	0.32	>5.0	36	Y	>5.0	>5.0	>5.0
22	Y	0.60	1.2	1.5	30	Y	<0.4	<0.5	0.35	36	W	0.27	1.2	>5.0
22	W	>5.0	>5.0	>5.0	30	W	4.4	>5.0	>5.0	36	W			>5.0
23	Y	>5.0	>5.0	>5.0	30	W	<0.4	>5.0	0.21					
n	Season	% Samples positive for Cry1Ab			n	Season	% Samples positive for Cry2Ab			n	Season	% Samples positive for CP4 EPSPS		
100	2020/21	90	86	90	2020/21	100	96	96	96	100	2020/21	98	98	98
70	2019/20	86	86	86	2019/20	70	97	97	97	70	2019/20	91	91	91
70	2018/19	91	91	91	2018/19	70	83	83	83	70	2018/19	96	96	96
100	2017/18	91	91	91	2017/18	100	83	83	83	100	2017/18	100	100	100
100	2016/17	98	98	98	2016/17	100	84	84	84	100	2016/17	100	100	100


LOD: Limit of Detection

# Mycotoxins

The South African commercial maize producers are privileged to have at their disposal, an excellent overview of the occurrence of the most important mycotoxins in maize, over an eleven season period and from all maize production regions in South Africa. During each of these seasons, 350 samples were selected from the maize crop survey samples, representing approximately 35 to 40% of the survey samples each season. The samples were always representatively selected for white and yellow maize from all the production regions. It must be noted that the grading defects of the maize samples were not considered in this selection, since as can be expected, no correlations were found between the visual inspection results (grading of the maize) and mycotoxin content. Thirteen mycotoxins including Aflatoxin B<sub>1</sub>, B<sub>2</sub>, G<sub>1</sub> and G<sub>2</sub>, Fumonisin (FUM) B<sub>1</sub>, B<sub>2</sub> and B<sub>3</sub>, Deoxynivalenol (DON), 15-acetyl-deoxynivalenol (15-ADON), Ochratoxin A, T2-toxin, H-T2 toxin and Zearalenone (ZON) were analysed with the LC-MS/MS instrument.

This season, 69% of the samples (240 samples) contained one or more mycotoxin, mainly DON (64%), FUM (22%) and ZON (3%). This was a notable decrease compared to the 90% occurrence in the previous season. The presence of DON and ZON, as illustrated in Graph 50, decreased for the first time in six years. The FUM prevalence decreased over the same period, from 57% in 2015/16 to 22% this season.

GRAPH 50: PERCENTAGE WHITE AND YELLOW MAIZE SAMPLES THAT TESTED POSITIVE FOR MYCOTOXINS OVER ELEVEN SEASONS



In Limpopo province, all the samples tested contained mycotoxins, as did more than 90% of samples in the Northern Cape and KwaZulu-Natal provinces. A decrease in mycotoxin occurrence was found in the other four provinces this season, ranging from an approximate 10% decrease to decreases of 84% and 83% in Mpumalanga and Gauteng respectively. In North West province, 57% of samples contained mycotoxins compared to the 83% during the previous season. In the Free State, samples containing mycotoxins decreased from 91% to 46%.

The results of the 13 mycotoxins obtained, including the range of concentration levels and notable trends in the mean concentration levels in white and yellow maize and in the different provinces, are summarised as follows:

## AFLATOXINS

No aflatoxins were found. Previously, aflatoxins were found in one yellow maize sample collected in North West province in 2019/20, in one white maize sample collected in the Free State in 2018/19 and in 2014/15 in three white maize samples collected in the North West province.

## OCHRATOXIN A, T2- TOXIN AND HT-2 TOXIN

None were reported in locally produced commercial maize since the survey began in 2010/11.


## DON AND 15-ADON

- Approximately 63% of white maize and 66% of yellow maize samples contained DON this season, a decrease compared to the previous season, but similar to the 2018/19 results. The DON concentration ranges are summarised in Graph 51 (white maize) and Graph 52 (yellow maize). The eleven-year mean DON concentrations in the seven provinces are illustrated in Graphs 53 and 54.

GRAPH 51: DON CONCENTRATION RANGE IN WHITE MAIZE SAMPLES OVER EIGHT SEASONS



GRAPH 52: DON CONCENTRATION RANGE IN YELLOW MAIZE SAMPLES OVER EIGHT SEASONS




- One percent of the white maize samples (collected in Mpumalanga) contained more than 2000 µg/kg DON, the national regulated maximum allowable level in unprocessed maize for human consumption. This is a decrease of 7% compared to the previous season when white maize samples with DON levels exceeding 2 000 µg/kg were collected in five of the production provinces.




- The white maize samples with the highest DON concentrations were collected in the Free State (1 159 µg/kg), Mpumalanga (3 256 µg/kg), Gauteng (1 599 µg/kg) and KwaZulu- Natal provinces (1 198 µg/kg).
- In yellow maize, similar to the previous two seasons, the highest DON concentration (2 169 µg/kg) was found in a sample from North West.
- Lower mean DON concentrations of the positive maize samples, compared to the previous season, were observed in white maize in five provinces and in all seven provinces in yellow maize. In Limpopo, DON was reported in white maize but not in yellow maize.

GRAPHS 53: WHITE MAIZE DON MEAN CONCENTRATION (µg/kg) PER PROVINCE OVER ELEVEN SEASONS



GRAPH 54: YELLOW MAIZE DON MEAN CONCENTRATION (µg/kg) PER PROVINCE OVER ELEVEN SEASONS




- When 15-ADON is found to be present in a sample, the sample also contains DON, mostly when the DON concentration level exceeds 500 µg/kg. This season, 82% of the samples did not contain 15-ADON, a 17% decrease in occurrence compared to the previous season. The mean 15-ADON concentration of the positive samples was 128 µg/kg.


**FUMONISINS ( $FUM\ TOTAL = FB1 + FB2 + FB3$ )**

- This season, only 18% of the white maize samples contained fumonisins, showing a continual decrease in contamination from the 62% in 2015/16. The yellow maize samples that contained fumonisins decreased from approximately 43% in the previous 4 seasons to 26% this season.
- Although the number of samples containing fumonisins decreased, FUM in the range of 4000 µg/kg (the SA regulated maximum allowable level in unprocessed maize for human consumption) was found in white maize from the Northern Cape (5 373 µg/kg) and Limpopo (3 952 µg/kg). The highest FUM (2 648 µg/kg) in a yellow maize sample was found in Mpumalanga. The concentration ranges of the samples with FUM over the past eight seasons are summarised in Graph 55 (white maize) and Graph 56 (yellow maize).

**GRAPH 55: FUM CONCENTRATION RANGE IN WHITE MAIZE SAMPLES OVER EIGHT SEASONS**




**GRAPH 56: FUM CONCENTRATION RANGE IN YELLOW MAIZE SAMPLES OVER EIGHT SEASONS**




- The white maize mean FUM concentrations were low in four production provinces, but increased in the Northern Cape, Limpopo and KwaZulu-Natal. The eleven-year FUM mean concentration variations in white maize in the seven provinces are illustrated in Graph 57.
- The mean FUM concentration in yellow maize ranged from 152 µg/kg in the Free State to 884 µg/kg in Limpopo. These trends are illustrated in Graph 58.

GRAPH 57: WHITE MAIZE FUM (TOTAL) MEAN CONCENTRATION ( $\mu\text{g}/\text{kg}$ ) PER PROVINCE OVER ELEVEN SEASONS




GRAPH 58: YELLOW MAIZE FUM (TOTAL) MEAN CONCENTRATION ( $\mu\text{g}/\text{kg}$ ) PER PROVINCE OVER ELEVEN SEASONS




## ZEARALENONE

- No zearalenone was found in white maize produced in the Northern Cape, North West, Gauteng and Limpopo nor in yellow maize produced in North West, Mpumalanga, Gauteng, Limpopo and KwaZulu-Natal. The ZON occurrences in the seven provinces are illustrated in Graphs 59 and 60 by reporting the mean ZON concentrations found in white and yellow maize.
- The highest concentration ZON in an individual white maize sample ( $101 \mu\text{g}/\text{kg}$ ) was reported in Mpumalanga this season.
- The multi-mycotoxin results over eleven consecutive seasons, provide an excellent South African perspective of commercially produced maize. The variation in occurrence and concentration levels of DON, 15-ADON, FUM and ZON confirmed that the mycotoxin risk varies significantly between production seasons in the different production regions and also maize class.

GRAPH 59: WHITE MAIZE ZON MEAN CONCENTRATION (µg/kg) PER PROVINCE OVER ELEVEN SEASONS



GRAPH 60: YELLOW MAIZE ZON MEAN CONCENTRATION (µg/kg) PER PROVINCE OVER ELEVEN SEASONS



The multi-mycotoxin results over eleven consecutive seasons provide an excellent South African perspective of the commercially produced maize. The variation in occurrence and concentration levels of DON, 15-ADON, FUM and ZON confirmed that the mycotoxin risk varies significantly between production seasons in the different production regions and maize class.

## International mycotoxin regulations

Information with regards to mycotoxin regulations per region and country, can be obtained from the Mycotoxins.info webpage supported by Biomin (<http://www.mycotoxins.info/regulations>).

# National mycotoxin regulations

According to the Foodstuffs, Cosmetics and Disinfectants Act (Act 54 of 1972) and regulations published in Government Notice No. R. 1145, dated 8 October 2004, all foodstuffs, ready for human consumption, may not contain more than 10 µg/kg of aflatoxin, of which aflatoxin B<sub>1</sub> may not exceed 5 µg/kg.

Amendments to Government Notice No. R. 1145, dated 8 October 2004, published in Government Notice No. 987 of 05 September 2016, specify that

- Cereal grains (wheat, maize and barley) intended for further processing, may not contain more than 2 000 µg/kg of Deoxynivalenol.
- Flour, meal, semolina and flakes derived from wheat, maize or barley, ready for human consumption, may not contain more than 1 000 µg/kg of Deoxynivalenol.
- Raw maize grain, intended for further processing, may not contain more than 4 000 µg/kg of Fumonisin (B<sub>1</sub> + B<sub>2</sub>), the whole commodity.
- Maize flour and maize meal, ready for human consumption, may not contain more than 2 000 µg/kg of Fumonisin (B<sub>1</sub> + B<sub>2</sub>), the whole commodity.

Further processing means any other treatment or processing method that has been proven to reduce levels of fungus produced toxins in foodstuffs intended for human consumption.

According to the Fertilizers, Farm Feeds, Agricultural Remedies and Stock Remedies Act (Act 36 of 1947) as well as amendments published in Government Notices No. R. 70 of 12 February 2010 and R. 789 of 10 September 2010, the maximum allowable levels of mycotoxins in animal feeds, are as follows:

Substance, Products	Farm Feeds	MAXIMUM CONTENT IN mg/kg (ppm) relative to a farm feed with a moisture content of 120 g/kg	MAXIMUM CONTENT IN µg/kg (ppb) relative to a farm feed with a moisture content of 120 g/kg
<b>Aflatoxin B<sub>1</sub></b>	Feed ingredients with the exception of:	0.05	50
	groundnut, copra, palm-kernel cotton seed, maize and products derived from the processing thereof	0.02	20
	Complete farm feeds for cattle, sheep and goats with the exception of:	0.05	50
	dairy cattle	0.005	5
	calves and lambs	0.01	10
	complete feeds for pigs and poultry (except young animals)	0.02	20
	other complete farm feeds (including pets)	0.01	10
	maize products intended for feedlot	0.3	300
<b>Deoxynivalenol (DON)</b>	supplement/concentrates for cattle, sheep and goats (except for dairy animals, calves and lambs)	0.05	50
	Feeding stuffs on a full ration basis for:		
	Pigs	1	1 000
	cattle	5	5 000
	calves up to 4 months	2	2 000
	dairy cattle	3	3 000
	poultry	4	4 000
<b>Fumonisin B<sub>1</sub></b>	pets	1	1 000
	Horses and pets	5	5 000
	Pigs	10	10 000
	Beef and poultry	50	50 000
<b>Ochratoxin A</b>	Fish	10	10 000
	Feeding stuffs on full ration basis for:		
	Pigs	0.05	50
<b>Zearalenone</b>	poultry	0.2	200
	Feeding stuffs on full ration basis for:		
	sows and pigs	5	5 000
<b>Zearalenone</b>	piglets	3	3 000
	calves and dairy cattle	0.5	500

**TABLE 24: MYCOTOXIN RESULTS - MAIZE CROP QUALITY 2020/21**

Region	Grade	Aflatoxin µg/kg						Fumonisin µg/kg			DON µg/kg	15-ADON µg/kg	Ochratoxin A µg/kg	Zearalenone µg/kg	HT-2 µg/kg	T-2 µg/kg	
		B <sub>1</sub> LOQ: 5 µg/kg	B <sub>2</sub> LOQ: 5 µg/kg	G <sub>1</sub> LOQ: 5 µg/kg	G <sub>2</sub> LOQ: 5 µg/kg	Total	B <sub>1</sub> LOQ: 20 µg/kg	B <sub>2</sub> LOQ: 20 µg/kg	B <sub>3</sub> LOQ: 20 µg/kg	Total							
10	YM1	ND	ND	ND	ND	ND	ND	ND	ND	278	68	30	376	196	ND	ND	ND
10	YM2	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	516	ND	ND	ND
10	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	108	ND	ND	ND
10	YM2	ND	ND	ND	ND	ND	ND	ND	ND	115	20	ND	135	401	ND	ND	ND
10	YM3	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	452	ND	ND	ND
10	YM1	ND	ND	ND	ND	ND	ND	ND	ND	112	ND	ND	112	291	ND	ND	ND
10	YM1	ND	ND	ND	ND	ND	ND	ND	ND	76	38	ND	114	265	ND	ND	ND
11	YM1	ND	ND	ND	ND	ND	ND	ND	ND	27	ND	ND	27	ND	ND	ND	ND
11	COM	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
11	WM2	ND	ND	ND	ND	ND	ND	ND	ND	3 610	1 459	304	5373	632	ND	ND	ND
12	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
12	WM1	ND	ND	ND	ND	ND	ND	ND	ND	55	ND	ND	55	198	ND	ND	ND
12	COM	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
12	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	108	ND	ND	ND
12	WM2	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	113	ND	ND	ND
12	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
13	WM1	ND	ND	ND	ND	ND	ND	ND	ND	127	44	ND	171	ND	ND	ND	ND
13	YM1	ND	ND	ND	ND	ND	ND	ND	ND	32	ND	ND	32	251	ND	ND	ND
13	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	520	ND	ND	ND
13	YM2	ND	ND	ND	ND	ND	ND	ND	ND	139	39	ND	178	ND	ND	ND	ND
13	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	244	ND	ND	ND
13	WM1	ND	ND	ND	ND	ND	ND	ND	ND	154	89	ND	243	239	ND	ND	ND
13	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
13	WM3	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
13	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	117	ND	ND	ND
13	WM2	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
13	COM	ND	ND	ND	ND	ND	ND	ND	ND	259	54	38	351	206	ND	ND	ND
14	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
14	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	183	ND	ND	ND
14	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND

**TABLE 24: MYCOTOXIN RESULTS - MAIZE CROP QUALITY 2020/21 (continue)**

Region	Grade	Aflatoxin µg/kg					Fumonisin µg/kg			DON µg/kg	15-ADON µg/kg	Ochratoxin A µg/kg	Zearalenone µg/kg	HT-2 µg/kg	T-2 µg/kg
		B <sub>1</sub> LOQ: 5 µg/kg	B <sub>2</sub> LOQ: 5 µg/kg	G <sub>1</sub> LOQ: 5 µg/kg	G <sub>2</sub> LOQ: 5 µg/kg	Total	B <sub>1</sub> LOQ: 20 µg/kg	B <sub>2</sub> LOQ: 20 µg/kg	B <sub>3</sub> LOQ: 20 µg/kg						
14	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
14	WM1	ND	ND	ND	ND	211	23	ND	234	267	ND	ND	ND	ND	ND
14	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
14	COM	ND	ND	ND	ND	22	ND	ND	22	182	ND	ND	ND	ND	ND
14	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
14	YM2	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
14	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
14	YM1	ND	ND	ND	ND	ND	ND	ND	ND	139	ND	ND	ND	ND	ND
14	COM	ND	ND	ND	ND	ND	ND	ND	ND	528	120	ND	ND	ND	ND
14	COM	ND	ND	ND	ND	ND	ND	ND	ND	167	ND	ND	ND	ND	ND
14	WM2	ND	ND	ND	ND	ND	ND	ND	ND	149	ND	ND	ND	ND	ND
14	WM1	ND	ND	ND	ND	ND	ND	ND	ND	250	ND	ND	ND	ND	ND
14	WM3	ND	ND	ND	ND	ND	ND	ND	ND	265	ND	ND	ND	ND	ND
16	YM2	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
16	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
16	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
17	WM3	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
17	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
17	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
17	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
17	WM1	ND	ND	ND	ND	ND	ND	ND	ND	107	ND	ND	ND	ND	ND
17	YM1	ND	ND	ND	ND	ND	ND	ND	ND	210	ND	ND	ND	ND	ND
17	COM	ND	ND	ND	ND	ND	ND	ND	ND	177	ND	ND	ND	ND	ND
17	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
17	YM1	ND	ND	ND	ND	232	78	ND	310	1 141	221	ND	ND	ND	ND
17	COM	ND	ND	ND	ND	ND	ND	ND	ND	761	127	ND	ND	ND	ND
17	WM2	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
17	WM1	ND	ND	ND	ND	ND	ND	ND	ND	662	121	ND	ND	ND	ND
18	YM1	ND	ND	ND	ND	ND	ND	ND	ND	104	ND	ND	ND	ND	ND
18	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND

**TABLE 24: MYCOTOXIN RESULTS - MAIZE CROP QUALITY 2020/21 (continue)**

Region	Grade	Aflatoxin µg/kg						Fumonisin µg/kg			DON µg/kg	15-ADON µg/kg	Ochratoxin A µg/kg	Zearalenone µg/kg	HT-2 µg/kg	T-2 µg/kg
		B <sub>1</sub> LOQ: 5 µg/kg	B <sub>2</sub> LOQ: 5 µg/kg	G <sub>1</sub> LOQ: 5 µg/kg	G <sub>2</sub> LOQ: 5 µg/kg	Total	B <sub>1</sub> LOQ: 20 µg/kg	B <sub>2</sub> LOQ: 20 µg/kg	B <sub>3</sub> LOQ: 20 µg/kg	Total						
18	YM1	ND	ND	ND	ND	ND	95	38	ND	133	ND	ND	ND	ND	ND	
18	COM	ND	ND	ND	ND	ND	74	ND	ND	74	2 169	237	ND	ND	ND	
18	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	465	ND	ND	ND	ND	
18	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	115	ND	ND	ND	ND	
18	WM1	ND	ND	ND	ND	ND	29	ND	ND	29	ND	ND	ND	ND	ND	
18	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	146	ND	ND	ND	ND	
18	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	
18	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	
18	YM2	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	
18	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	
18	WM2	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	
18	WM2	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	
19	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	
19	WM3	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	
19	YM2	ND	ND	ND	ND	ND	179	51	ND	230	ND	ND	ND	ND	ND	
19	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	
19	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	105	ND	ND	ND	ND	
19	YM1	ND	ND	ND	ND	ND	232	74	22	328	196	ND	ND	ND	ND	
19	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	166	ND	ND	ND	ND	
19	WM2	ND	ND	ND	ND	ND	91	28	ND	119	577	ND	ND	ND	ND	
19	WM2	ND	ND	ND	ND	ND	ND	ND	ND	ND	163	ND	ND	ND	ND	
19	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	
19	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	379	ND	ND	ND	ND	
19	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	
19	WM3	ND	ND	ND	ND	ND	ND	ND	ND	ND	325	ND	ND	ND	ND	
20	YM1	ND	ND	ND	ND	ND	126	33	ND	159	ND	ND	ND	ND	ND	
20	YM1	ND	ND	ND	ND	ND	432	235	24	691	ND	ND	ND	ND	ND	
20	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	331	129	ND	ND	ND	
20	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	651	147	ND	ND	ND	
20	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	491	ND	ND	ND	ND	



**TABLE 24: MYCOTOXIN RESULTS - MAIZE CROP QUALITY 2020/21 (continue)**

Region	Grade	Aflatoxin µg/kg					Fumonisin µg/kg			DON µg/kg LOQ: 100 µg/kg	15-ADON µg/kg LOQ: 100 µg/kg	Ochratoxin A µg/kg LOQ: 5 µg/kg	Zearalenone µg/kg LOQ: 20 µg/kg	HT-2 µg/kg LOD: 20 µg/kg	T-2 µg/kg LOQ: 20 µg/kg
		B <sub>1</sub> LOQ: 5 µg/kg	B <sub>2</sub> LOQ: 5 µg/kg	G <sub>1</sub> LOQ: 5 µg/kg	G <sub>2</sub> LOQ: 5 µg/kg	Total	B <sub>1</sub> LOQ: 20 µg/kg	B <sub>2</sub> LOQ: 20 µg/kg	B <sub>3</sub> LOQ: 20 µg/kg						
20	YM1	ND	ND	ND	ND	ND	ND	ND	ND	1 165	208	ND	ND	ND	ND
20	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
20	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
20	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
20	WM1	ND	ND	ND	ND	ND	ND	ND	810	151	ND	ND	ND	ND	ND
20	WM1	ND	ND	ND	ND	ND	ND	ND	119	ND	ND	ND	ND	ND	ND
21	WM1	ND	ND	ND	ND	ND	ND	ND	439	105	ND	ND	ND	ND	ND
21	YM1	ND	ND	ND	ND	77	33	ND	284	ND	ND	ND	ND	ND	ND
21	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
21	WM2	ND	ND	ND	ND	ND	ND	ND	270	ND	ND	ND	ND	ND	ND
21	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
21	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
21	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
21	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
21	COM	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
21	WM2	ND	ND	ND	ND	ND	ND	ND	1 159	315	ND	ND	ND	ND	ND
21	YM2	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
21	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
21	YM3	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
21	COM	ND	ND	ND	ND	ND	ND	ND	109	ND	ND	ND	ND	ND	ND
21	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
21	WM3	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
22	WM1	ND	ND	ND	ND	ND	ND	ND	111	ND	ND	ND	ND	ND	ND
22	YM1	ND	ND	ND	ND	ND	ND	ND	330	ND	ND	ND	ND	ND	ND
22	WM1	ND	ND	ND	ND	ND	ND	ND	170	ND	ND	ND	ND	ND	ND
22	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
22	YM2	ND	ND	ND	ND	ND	ND	ND	307	ND	ND	ND	ND	ND	ND
22	COM	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
22	WM1	ND	ND	ND	ND	ND	ND	ND	185	ND	ND	ND	ND	ND	ND
22	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND

TABLE 24: MYCOTOXIN RESULTS - MAIZE CROP QUALITY 2020/21 (continue)															
Region	Grade	Aflatoxin µg/kg				Total	Fumonisin µg/kg			DON µg/kg LOQ: 100 µg/kg	15-ADON µg/kg LOQ: 100 µg/kg	Ochratoxin A µg/kg LOQ: 5 µg/kg	Zearalenone µg/kg LOQ: 20 µg/kg	HT-2 µg/kg LOQ: 20 µg/kg	T-2 µg/kg LOQ: 20 µg/kg
		B <sub>1</sub> LOQ: 5 µg/kg	B <sub>2</sub> LOQ: 5 µg/kg	G <sub>1</sub> LOQ: 5 µg/kg	G <sub>2</sub> LOQ: 5 µg/kg		B <sub>1</sub> LOQ: 20 µg/kg	B <sub>2</sub> LOQ: 20 µg/kg	B <sub>3</sub> LOQ: 20 µg/kg						
20	YM1	ND	ND	ND	ND	ND	ND	ND	ND	1 165	208	ND	ND	ND	ND
20	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
20	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
20	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
20	WM1	ND	ND	ND	ND	ND	ND	ND	810	151	ND	ND	ND	ND	ND
20	WM1	ND	ND	ND	ND	ND	ND	ND	119	ND	ND	ND	ND	ND	ND
21	WM1	ND	ND	ND	ND	ND	ND	ND	439	105	ND	ND	ND	ND	ND
21	YM1	ND	ND	ND	ND	ND	77	33	284	ND	ND	ND	ND	ND	ND
21	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
21	WM2	ND	ND	ND	ND	ND	ND	ND	270	ND	ND	ND	ND	ND	ND
21	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
21	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
21	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
21	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
21	COM	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
21	WM2	ND	ND	ND	ND	ND	ND	ND	1 159	315	ND	ND	ND	ND	ND
21	YM2	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
21	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
21	YM3	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
21	COM	ND	ND	ND	ND	ND	ND	ND	109	ND	ND	ND	ND	ND	ND
21	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
21	WM3	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
22	WM1	ND	ND	ND	ND	ND	ND	ND	111	ND	ND	ND	ND	ND	ND
22	YM1	ND	ND	ND	ND	ND	ND	ND	330	ND	ND	ND	ND	ND	ND
22	WM1	ND	ND	ND	ND	ND	ND	ND	170	ND	ND	ND	ND	ND	ND
22	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
22	YM2	ND	ND	ND	ND	ND	ND	ND	307	ND	ND	ND	ND	ND	ND
22	COM	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
22	WM1	ND	ND	ND	ND	ND	ND	ND	185	ND	ND	ND	ND	ND	ND
22	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND

TABLE 24: MYCOTOXIN RESULTS - MAIZE CROP QUALITY 2020/21 (continue)															
Region	Grade	Aflatoxin µg/kg				Total	Fumonisin µg/kg			DON µg/kg LOQ: 100 µg/kg	15-ADON µg/kg LOQ: 100 µg/kg	Ochratoxin A µg/kg LOQ: 5 µg/kg	Zearalenone µg/kg LOQ: 20 µg/kg	HT-2 µg/kg LOQ: 20 µg/kg	T-2 µg/kg LOQ: 20 µg/kg
		B <sub>1</sub> LOQ: 5 µg/kg	B <sub>2</sub> LOQ: 5 µg/kg	G <sub>1</sub> LOQ: 5 µg/kg	G <sub>2</sub> LOQ: 5 µg/kg		B <sub>1</sub> LOQ: 20 µg/kg	B <sub>2</sub> LOQ: 20 µg/kg	B <sub>3</sub> LOQ: 20 µg/kg						
23	WM1	ND	ND	ND	ND	ND	ND	ND	ND	154	ND	ND	ND	ND	ND
23	WM1	ND	ND	ND	ND	ND	ND	ND	ND	114	ND	ND	ND	ND	ND
23	WM2	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
23	YM1	ND	ND	ND	ND	ND	178	50	ND	ND	ND	ND	ND	ND	ND
23	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
23	COM	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
23	YM1	ND	ND	ND	ND	ND	94	23	117	124	ND	ND	ND	ND	ND
23	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
23	WM3	ND	ND	ND	ND	ND	ND	ND	ND	315	ND	ND	ND	ND	ND
23	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
23	YM1	ND	ND	ND	ND	ND	ND	ND	ND	208	ND	ND	ND	ND	ND
23	WM2	ND	ND	ND	ND	ND	ND	ND	ND	516	ND	ND	ND	ND	ND
23	WM2	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
23	COM	ND	ND	ND	ND	ND	193	46	239	269	ND	ND	ND	ND	ND
23	WM1	ND	ND	ND	ND	ND	ND	ND	ND	144	ND	ND	ND	ND	ND
23	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
24	WM1	ND	ND	ND	ND	ND	ND	ND	ND	181	ND	ND	ND	ND	ND
24	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
24	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
24	YM2	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
24	YM1	ND	ND	ND	ND	ND	ND	ND	ND	531	105	52	ND	ND	ND
24	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
25	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
25	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
25	COM	ND	ND	ND	ND	ND	ND	ND	ND	353	ND	ND	ND	ND	ND
25	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
25	WM1	ND	ND	ND	ND	ND	ND	ND	ND	331	ND	ND	ND	ND	ND
25	WM1	ND	ND	ND	ND	ND	ND	ND	ND	122	ND	ND	ND	ND	ND
25	YM1	ND	ND	ND	ND	ND	ND	ND	ND	173	ND	ND	ND	ND	ND
25	COM	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND

TABLE 24: MYCOTOXIN RESULTS - MAIZE CROP QUALITY 2020/21 (continue)															
Region	Grade	Aflatoxin µg/kg				Total	Fumonisin µg/kg			DON µg/kg LOQ: 100 µg/kg	15-ADON µg/kg LOQ: 100 µg/kg	Ochratoxin A µg/kg LOQ: 5 µg/kg	Zearalenone µg/kg LOQ: 20 µg/kg	HT-2 µg/kg LOQ: 20 µg/kg	T-2 µg/kg LOQ: 20 µg/kg
		B <sub>1</sub> LOQ: 5 µg/kg	B <sub>2</sub> LOQ: 5 µg/kg	G <sub>1</sub> LOQ: 5 µg/kg	G <sub>2</sub> LOQ: 5 µg/kg		B <sub>1</sub> LOQ: 20 µg/kg	B <sub>2</sub> LOQ: 20 µg/kg	B <sub>3</sub> LOQ: 20 µg/kg						
25	YM2	ND	ND	ND	ND	ND	ND	ND	ND	199	ND	ND	ND	ND	ND
26	COM	ND	ND	ND	ND	ND	ND	ND	124	ND	ND	ND	ND	ND	ND
26	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
26	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
26	WM2	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
26	WM3	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
26	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
26	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
26	YM1	ND	ND	ND	ND	ND	ND	ND	150	ND	ND	ND	ND	ND	ND
26	WM2	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
26	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
26	COM	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
26	YM1	ND	ND	ND	ND	ND	ND	ND	222	ND	ND	ND	ND	ND	ND
26	YM2	ND	ND	ND	ND	ND	ND	ND	129	ND	ND	ND	ND	ND	ND
27	WM2	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
27	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
27	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
28	YM2	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
28	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
28	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
28	WM1	ND	ND	ND	ND	ND	ND	ND	113	ND	ND	ND	ND	ND	ND
28	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
28	WM1	ND	ND	ND	ND	ND	ND	ND	306	ND	ND	ND	ND	ND	ND
28	YM1	ND	ND	ND	ND	ND	ND	ND	188	ND	ND	ND	ND	ND	ND
28	WM1	ND	ND	ND	ND	ND	ND	ND	244	ND	ND	ND	ND	ND	ND
28	COM	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
28	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
28	WM1	ND	ND	ND	ND	ND	ND	62	133	466	118	ND	ND	ND	ND
28	WM1	ND	ND	ND	ND	ND	ND	ND	796	155	ND	ND	ND	ND	ND
28	YM2	ND	ND	ND	ND	ND	ND	ND	356	ND	ND	ND	ND	ND	ND

TABLE 24: MYCOTOXIN RESULTS - MAIZE CROP QUALITY 2020/21 (continue)															
Region	Grade	Aflatoxin µg/kg				Total	Fumonisin µg/kg			DON µg/kg LOQ: 100 µg/kg	15-ADON µg/kg LOQ: 100 µg/kg	Ochratoxin A µg/kg LOQ: 5 µg/kg	Zearalenone µg/kg LOQ: 20 µg/kg	HT-2 µg/kg LOQ: 20 µg/kg	T-2 µg/kg LOQ: 20 µg/kg
		B <sub>1</sub> LOQ: 5 µg/kg	B <sub>2</sub> LOQ: 5 µg/kg	G <sub>1</sub> LOQ: 5 µg/kg	G <sub>2</sub> LOQ: 5 µg/kg		B <sub>1</sub> LOQ: 20 µg/kg	B <sub>2</sub> LOQ: 20 µg/kg	B <sub>3</sub> LOQ: 20 µg/kg						
28	WM1	ND	ND	ND	ND	ND	ND	ND	ND	153	ND	ND	ND	ND	ND
28	WM1	ND	ND	ND	ND	ND	ND	ND	ND	312	ND	ND	27	ND	ND
28	YM2	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
28	YM1	ND	ND	ND	ND	ND	ND	ND	ND	170	ND	ND	ND	ND	ND
28	YM1	ND	ND	ND	ND	ND	ND	ND	ND	290	ND	ND	ND	ND	ND
29	YM3	ND	ND	ND	ND	ND	ND	ND	ND	263	101	ND	ND	ND	ND
29	YM1	ND	ND	ND	ND	ND	ND	98	45	20	163	ND	ND	ND	ND
29	WM1	ND	ND	ND	ND	ND	ND	ND	ND	1 132	262	ND	ND	ND	ND
29	YM1	ND	ND	ND	ND	ND	ND	ND	ND	194	ND	ND	ND	ND	ND
29	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
29	WM1	ND	ND	ND	ND	ND	ND	ND	ND	431	ND	ND	ND	ND	ND
29	WM1	ND	ND	ND	ND	ND	ND	ND	ND	139	ND	ND	ND	ND	ND
29	YM1	ND	ND	ND	ND	ND	ND	ND	ND	609	164	ND	ND	ND	ND
29	COM	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
29	WM1	ND	ND	ND	ND	ND	ND	ND	ND	615	179	ND	ND	ND	ND
29	YM1	ND	ND	ND	ND	ND	ND	412	111	32	555	ND	ND	ND	ND
29	WM2	ND	ND	ND	ND	ND	ND	1 646	589	121	2356	184	101	ND	ND
29	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
29	YM1	ND	ND	ND	ND	ND	ND	ND	ND	168	ND	ND	ND	ND	ND
29	YM1	ND	ND	ND	ND	ND	ND	88	32	100	120	ND	ND	ND	ND
29	WM1	ND	ND	ND	ND	ND	ND	ND	ND	178	ND	ND	ND	ND	ND
29	WM1	ND	ND	ND	ND	ND	ND	ND	ND	171	ND	ND	ND	ND	ND
29	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
29	WM1	ND	ND	ND	ND	ND	ND	ND	ND	720	111	ND	ND	ND	ND
29	WM1	ND	ND	ND	ND	ND	ND	315	137	29	481	ND	ND	ND	ND
30	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
30	YM1	ND	ND	ND	ND	ND	ND	ND	ND	527	111	ND	ND	ND	ND
30	YM1	ND	ND	ND	ND	ND	ND	ND	ND	283	ND	ND	ND	ND	ND
30	WM3	ND	ND	ND	ND	ND	ND	ND	ND	594	105	ND	ND	ND	ND
30	WM1	ND	ND	ND	ND	ND	ND	ND	ND	887	ND	ND	ND	ND	ND

TABLE 24: MYCOTOXIN RESULTS - MAIZE CROP QUALITY 2020/21 (continue)															
Region	Grade	Aflatoxin µg/kg				Total	Fumonisin µg/kg			DON µg/kg LOQ: 100 µg/kg	15-ADON µg/kg LOQ: 100 µg/kg	Ochratoxin A µg/kg LOQ: 5 µg/kg	Zearalenone µg/kg LOQ: 20 µg/kg	HT-2 µg/kg LOQ: 20 µg/kg	T-2 µg/kg LOQ: 20 µg/kg
		B <sub>1</sub> LOQ: 5 µg/kg	B <sub>2</sub> LOQ: 5 µg/kg	G <sub>1</sub> LOQ: 5 µg/kg	G <sub>2</sub> LOQ: 5 µg/kg		B <sub>1</sub> LOQ: 20 µg/kg	B <sub>2</sub> LOQ: 20 µg/kg	B <sub>3</sub> LOQ: 20 µg/kg						
30	COM	ND	ND	ND	ND	ND	ND	ND	ND	487	100	ND	ND	ND	ND
30	WM1	ND	ND	ND	ND	ND	264	93	22	995	184	ND	ND	ND	ND
30	YM1	ND	ND	ND	ND	ND	ND	ND	ND	510	ND	ND	ND	ND	ND
30	YM1	ND	ND	ND	ND	ND	ND	ND	ND	277	ND	ND	ND	ND	ND
30	WM1	ND	ND	ND	ND	ND	ND	ND	ND	1 606	359	ND	ND	ND	ND
30	WM1	ND	ND	ND	ND	ND	ND	ND	ND	142	ND	ND	ND	ND	ND
30	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
30	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
30	YM1	ND	ND	ND	ND	ND	ND	ND	ND	111	ND	ND	ND	ND	ND
30	COM	ND	ND	ND	ND	ND	ND	ND	ND	391	ND	ND	ND	ND	ND
30	YM3	ND	ND	ND	ND	ND	ND	ND	ND	474	ND	ND	ND	ND	ND
30	WM1	ND	ND	ND	ND	ND	76	38	ND	470	ND	ND	20	ND	ND
30	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
30	WM2	ND	ND	ND	ND	ND	ND	ND	ND	294	163	ND	ND	ND	ND
30	COM	ND	ND	ND	ND	ND	ND	ND	ND	715	ND	ND	ND	ND	ND
30	WM2	ND	ND	ND	ND	ND	35	ND	ND	950	162	ND	ND	ND	ND
30	WM1	ND	ND	ND	ND	ND	ND	ND	ND	750	ND	ND	ND	ND	ND
30	YM1	ND	ND	ND	ND	ND	268	130	23	291	ND	ND	ND	ND	ND
30	YM2	ND	ND	ND	ND	ND	ND	ND	ND	573	111	ND	ND	ND	ND
30	WM2	ND	ND	ND	ND	ND	ND	ND	ND	359	ND	ND	ND	ND	ND
31	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
31	YM1	ND	ND	ND	ND	ND	ND	ND	ND	243	ND	ND	ND	ND	ND
31	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
31	YM1	ND	ND	ND	ND	ND	ND	ND	ND	123	ND	ND	ND	ND	ND
31	YM1	ND	ND	ND	ND	ND	44	ND	ND	204	ND	ND	ND	ND	ND
31	COM	ND	ND	ND	ND	ND	483	183	40	100	ND	ND	ND	ND	ND
31	YM2	ND	ND	ND	ND	ND	143	90	ND	327	ND	ND	ND	ND	ND
31	COM	ND	ND	ND	ND	ND	ND	ND	ND	432	122	ND	ND	ND	ND
31	COM	ND	ND	ND	ND	ND	50	26	ND	732	160	ND	ND	ND	ND
31	WM1	ND	ND	ND	ND	ND	ND	ND	ND	593	ND	ND	ND	ND	ND

TABLE 24: MYCOTOXIN RESULTS - MAIZE CROP QUALITY 2020/21 (continue)																
Region	Grade	Aflatoxin µg/kg				Total	Fumonisin µg/kg				DON µg/kg LOQ: 100 µg/kg	15-ADON µg/kg LOQ: 100 µg/kg	Ochratoxin A µg/kg LOQ: 5 µg/kg	Zearalenone µg/kg LOQ: 20 µg/kg	HT-2 µg/kg LOQ: 20 µg/kg	T-2 µg/kg LOQ: 20 µg/kg
		B <sub>1</sub> LOQ: 5 µg/kg	B <sub>2</sub> LOQ: 5 µg/kg	G <sub>1</sub> LOQ: 5 µg/kg	G <sub>2</sub> LOQ: 5 µg/kg		B <sub>1</sub> LOQ: 20 µg/kg	B <sub>2</sub> LOQ: 20 µg/kg	B <sub>3</sub> LOQ: 20 µg/kg	Total						
31	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	184	ND	ND	ND	ND	ND
31	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	151	ND	ND	ND	ND	ND
31	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
31	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	1 248	ND	ND	ND	ND	ND
31	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	199	ND	ND	ND	ND	ND
31	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	342	ND	ND	ND	ND	ND
31	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	227	ND	ND	ND	ND	ND
31	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	340	ND	ND	ND	ND	ND
31	WM2	ND	ND	ND	ND	ND	ND	ND	ND	ND	507	ND	ND	ND	ND	ND
31	WM2	ND	ND	ND	ND	ND	ND	ND	ND	ND	2 945	ND	ND	ND	ND	ND
31	WM3	ND	ND	ND	ND	ND	ND	ND	ND	ND	128	ND	ND	ND	ND	ND
32	WM2	ND	ND	ND	ND	ND	ND	ND	ND	ND	1 953	ND	ND	ND	ND	ND
32	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	585	ND	ND	ND	ND	ND
32	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	252	ND	ND	ND	ND	ND
32	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	243	ND	ND	ND	ND	ND
32	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	323	ND	ND	ND	ND	ND
32	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	234	ND	ND	ND	ND	ND
32	COM	ND	ND	ND	ND	ND	ND	ND	ND	ND	313	ND	ND	ND	ND	ND
32	COM	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
32	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	267	ND	ND	ND	ND	ND
32	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	696	ND	ND	ND	ND	ND
32	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	178	ND	ND	ND	ND	ND
32	WM2	ND	ND	ND	ND	ND	ND	ND	ND	ND	3 256	ND	ND	ND	ND	ND
32	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	331	ND	ND	ND	ND	ND
32	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	296	ND	ND	ND	ND	ND
32	WM2	ND	ND	ND	ND	ND	ND	ND	ND	ND	1 068	ND	ND	ND	ND	ND
32	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	128	ND	ND	ND	ND	ND
32	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	646	ND	ND	ND	ND	ND
32	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	735	ND	ND	ND	ND	ND
32	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	386	ND	ND	ND	ND	ND

TABLE 24: MYCOTOXIN RESULTS - MAIZE CROP QUALITY 2020/21 (continue)																
Region	Grade	Aflatoxin µg/kg				Total	Fumonisin µg/kg				DON µg/kg LOQ: 100 µg/kg	15-ADON µg/kg LOQ: 100 µg/kg	Ochratoxin A µg/kg LOQ: 5 µg/kg	Zearalenone µg/kg LOQ: 20 µg/kg	HT-2 µg/kg LOQ: 20 µg/kg	T-2 µg/kg LOQ: 20 µg/kg
		B <sub>1</sub> LOQ: 5 µg/kg	B <sub>2</sub> LOQ: 5 µg/kg	G <sub>1</sub> LOQ: 5 µg/kg	G <sub>2</sub> LOQ: 5 µg/kg		B <sub>1</sub> LOQ: 20 µg/kg	B <sub>2</sub> LOQ: 20 µg/kg	B <sub>3</sub> LOQ: 20 µg/kg	Total						
32	YM1	ND	ND	ND	ND	ND	277	78	ND	355	124	ND	ND	ND	ND	ND
32	YM2	ND	ND	ND	ND	ND	496	198	41	735	ND	ND	ND	ND	ND	ND
32	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
32	WM2	ND	ND	ND	ND	ND	126	53	ND	179	1 392	244	ND	ND	ND	ND
32	WM2	ND	ND	ND	ND	ND	ND	ND	ND	179	1 217	158	ND	ND	ND	ND
32	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	862	103	20	ND	ND	ND
33	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	461	129	ND	ND	ND	ND
33	YM1	ND	ND	ND	ND	ND	434	142	29	605	776	159	ND	ND	ND	ND
33	WM1	ND	ND	ND	ND	ND	99	39	ND	138	ND	ND	ND	ND	ND	ND
33	YM1	ND	ND	ND	ND	ND	42	ND	ND	42	283	ND	ND	ND	ND	ND
33	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	501	100	ND	ND	ND	ND
33	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	478	107	ND	ND	ND	ND
33	YM3	ND	ND	ND	ND	ND	475	180	32	687	139	ND	ND	ND	ND	ND
33	WM3	ND	ND	ND	ND	ND	ND	ND	ND	ND	179	ND	ND	ND	ND	ND
33	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	223	ND	ND	ND	ND	ND
33	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	327	ND	ND	ND	ND	ND
33	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
33	YM1	ND	ND	ND	ND	ND	1 945	514	189	2 648	ND	ND	ND	ND	ND	ND
33	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	800	162	ND	ND	ND	ND
33	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	152	ND	ND	ND	ND	ND
33	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	331	ND	ND	ND	ND	ND
33	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
33	COM	ND	ND	ND	ND	ND	ND	ND	ND	ND	114	ND	ND	ND	ND	ND
33	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	600	103	ND	ND	ND	ND
33	COM	ND	ND	ND	ND	ND	ND	ND	ND	ND	424	159	ND	ND	ND	ND
33	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
33	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND
33	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	316	ND	ND	ND	ND	ND
33	WM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	149	ND	ND	ND	ND	ND
33	WM2	ND	ND	ND	ND	ND	ND	ND	ND	ND	562	118	ND	ND	ND	ND





TABLE 24: MYCOTOXIN RESULTS - MAIZE CROP QUALITY 2020/21 (continue)														
Region	Grade	Aflatoxin µg/kg			Fumonisin µg/kg			DON µg/kg	15-ADON µg/kg	Ochratoxin A µg/kg	Zearalenone µg/kg	HT-2 µg/kg	T-2 µg/kg	
		B <sub>1</sub> LOQ: 5 µg/kg	B <sub>2</sub> LOQ: 5 µg/kg	G <sub>1</sub> LOQ: 5 µg/kg	G <sub>2</sub> LOQ: 5 µg/kg	Total	B <sub>1</sub> LOQ: 20 µg/kg							B <sub>2</sub> LOQ: 20 µg/kg
35	YM1	ND	ND	ND	ND	1 246	343	119	1 708	ND	ND	ND	ND	
35	WM1	ND	ND	ND	ND	41	ND	ND	41	ND	ND	ND	ND	
36	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	
36	WM2	ND	ND	ND	ND	244	67	ND	311	123	ND	ND	ND	
36	WM1	ND	ND	ND	ND	ND	ND	ND	ND	425	ND	ND	ND	
36	YM1	ND	ND	ND	ND	93	26	ND	119	151	ND	ND	ND	
36	WM1	ND	ND	ND	ND	ND	ND	ND	ND	164	ND	ND	ND	
36	COM	ND	ND	ND	ND	ND	ND	ND	ND	247	ND	ND	ND	
36	YM1	ND	ND	ND	ND	ND	ND	ND	ND	251	ND	ND	ND	
36	COM	ND	ND	ND	ND	ND	ND	ND	ND	138	ND	ND	ND	
36	YM1	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	
36	WM2	ND	ND	ND	ND	567	224	61	852	186	22	ND	ND	
36	YM1	ND	ND	ND	ND	652	255	48	955	414	ND	ND	ND	
36	YM1	ND	ND	ND	ND	ND	ND	ND	ND	391	ND	ND	ND	
36	WM1	ND	ND	ND	ND	ND	ND	ND	ND	510	ND	ND	ND	
36	YM1	ND	ND	ND	ND	38	ND	ND	38	442	104	ND	ND	
36	WM1	ND	ND	ND	ND	ND	ND	ND	ND	1 095	210	ND	ND	
36	YM2	ND	ND	ND	ND	ND	ND	ND	ND	151	ND	ND	ND	
36	YM1	ND	ND	ND	ND	ND	ND	ND	ND	360	ND	ND	ND	
36	WM1	ND	ND	ND	ND	299	87	27	413	574	ND	ND	ND	
<b>Total number of samples</b>		<b>350</b>	<b>350</b>	<b>350</b>	<b>350</b>	<b>350</b>	<b>350</b>	<b>350</b>	<b>350</b>	<b>350</b>	<b>350</b>	<b>350</b>	<b>350</b>	
<b>Average of total number of samples</b>		<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>71</b>	<b>24</b>	<b>5</b>	<b>100</b>	<b>279</b>	<b>32</b>	<b>0</b>	<b>0</b>	
<b>Number of positive results</b>		<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>76</b>	<b>55</b>	<b>29</b>	<b>76</b>	<b>225</b>	<b>63</b>	<b>0</b>	<b>0</b>	
<b>Average of positive results</b>		<b>-</b>	<b>-</b>	<b>-</b>	<b>-</b>	<b>325</b>	<b>154</b>	<b>60</b>	<b>459</b>	<b>434</b>	<b>176</b>	<b>-</b>	<b>-</b>	
<b>Maximum of positive results</b>		<b>-</b>	<b>-</b>	<b>-</b>	<b>-</b>	<b>3 610</b>	<b>1 459</b>	<b>304</b>	<b>5 373</b>	<b>3 256</b>	<b>571</b>	<b>-</b>	<b>-</b>	

**Note:**

Limit of quantitation (LOQ) means the lowest concentration level that can be quantified with acceptable precision and accuracy by the LC-MS/MS.

A concentration measured below the LOQ is reported as <LOQ.

Limit of detection (LOD) is the lowest concentration level that can be detected but not quantified and is 50% of the LOQ of each mycotoxin.

A concentration measured below the LOD is reported as not detected (ND).

µg/kg = ppb (parts per billion)

**TABLE 25: MYCOTOXIN RESULTS - SUMMARY OF SEASONS 2011/12 TO 2020/21**

Season	Total Number of samples received	Number of samples tested for mycotoxins	Aflatoxin µg/kg			Fumonisin µg/kg			Deoxynivalenol µg/kg			Zearalenone µg/kg			Ochratoxin A µg/kg			T-2 Toxin µg/kg		
			ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.	ave.	min.	max.
**2011/12	1 000	350	0	0	0	383	0	11 297	146	0	911	33	0	297	0	0	0	0	0	0
**2012/13	1 000	350	0	0	0	530	0	11 243	186	0	1 175	30	0	426	0	0	2	0	232	0
**2013/14	930	350	0	0	0	451	0	5 357	243	0	6 134	38	0	445	0	0	0	0	0	0
**2014/15	1 000	350	2	0	48	357	0	3 382	397	0	9 736	36	0	337	0	0	0	0	0	0
**2015/16	920	350	0	0	0	444	0	11 347	175	0	1 585	16	0	127	0	0	0	0	0	0
**2016/17	1 000	350	0	0	0	471	0	6 059	513	0	7 698	36	0	399	0	0	0	0	0	0
**2017/18	900	350	0	0	0	991	0	8 356	656	0	3 510	51	0	361	0	0	0	0	0	0
**2018/19	808	350	10	0	143	666	0	34 740	550	0	11 181	64	0	957	0	0	0	0	0	0
**2019/20	890	350	1	0	10	361	0	5 928	898	0	7 700	43	0	539	0	0	0	0	0	0
**2020/21	1 000	350	0	0	0	724	0	5 373	321	0	3 256	12	0	101	0	0	0	0	0	0
<b>Total</b>	<b>9 448</b>	<b>3 500</b>																		
	<b>Min.</b>		<b>0</b>				<b>0</b>			<b>0</b>					<b>0</b>				<b>0</b>	
	<b>Max.</b>				<b>143</b>			<b>34 740</b>			<b>11 181</b>			<b>957</b>			<b>0</b>			<b>232</b>

\*\* Sum of Aflatoxin (B<sub>1</sub>; B<sub>2</sub>; G<sub>1</sub>; G<sub>2</sub>) and sum of Fumonisin (B<sub>1</sub>; B<sub>2</sub>; B<sub>3</sub>)   RSA averages calculated from averages per province.

**Mycotoxin methodology**

During 2010 SAGL implemented a multi-mycotoxin screening method using UPLC-MS/MS. The following limit of detection applies for each toxin:

Mycotoxin	LOQ for maize µg/kg	LOD for maize µg/kg
Aflatoxin B <sub>1</sub>	5	2.5
Aflatoxin B <sub>2</sub>	5	2.5
Aflatoxin G <sub>1</sub>	5	2.5
Aflatoxin G <sub>2</sub>	5	2.5
Fumonisin B <sub>1</sub>	20	10
Fumonisin B <sub>2</sub>	20	10
Fumonisin B <sub>3</sub>	20	10
Deoxynivalenol	100	50
Zearalenone	20	10
Ochratoxin A	5	2.5
T - 2 Toxin	20	10

**Notes:**

- Limit of detection (LOD) means the lowest level that can be detected accurately by the technique.
- Limit of quantitation (LOQ) means the lowest level that can be quantified accurately by the technique.
- A result above zero but lower than the limit of detection/quantitation, is reported as <LOD/<LOQ.

µg/kg = ppb (parts per billion)

# Methods

## SAMPLING PROCEDURE

A working group determined the process to be followed to ensure that the crop quality samples sent to the SAGL by the various grain silo owners/ agricultural businesses, are representative of the total crop.

Each delivery is sampled as per the grading regulations for grading purposes.

Once grading has been completed, a sub-sample of each of these grading samples are placed in separate containers according to class and grade, per silo bin at each silo.

After 80% of the expected harvest has been received, the content of each container was divided with a multi slot divider in order to obtain a 3 kg sample (this should be done for each class and grade separately).

If there was more than one container per class and grade per silo bin, the combined contents of the containers were mixed thoroughly before dividing it with a multi slot divider to obtain the required 3 kg sample.

The samples were marked clearly with the name of the depot, the bin/bag/bunker/dam number(s) represented by each individual sample as well as the class and grade and were then forwarded to the SAGL.

## RSA GRADING

RSA grading was done in accordance with the Grading Regulations for maize, as published in the Government Gazette No. 32190, Government Notice No. R. 473 of 8 May 2009 and amended by Industry-Wide Dispensation REF NO: 20/4/14/1, dated 15 April 2010.

Description of deviations relating to RSA grading:

### DEFECTIVE MAIZE KERNELS

*The following definition of Defective maize kernels is quoted from the Grading Regulations:*

“Defective maize kernels” means maize kernels and pieces of maize kernels –

- (a) that are shrivelled, obviously immature, frost-damaged, heat damaged, water damaged, mouldy or chalky;
- (b) that are discoloured by external factors such

as water and sun: Provided that discoloration on both sides of the maize kernel limited to less than a quarter from the bottom tip of the maize kernel shall not be considered as defective, oxidation stained maize kernels, coffee stained maize kernels and pinked maize kernels shall not be considered as defective;

- (c) that have sprouted, including kernels which the shoot (plumule) in the germ is visibly discoloured;
- (d) that have cavities in the germ or endosperm caused by insects or rodents;
- (e) that are visibly soiled (smeared) or contaminated by smut, fire, soil, smoke or coal-dust;
- (f) all matter that can pass through the 6.35 mm round-hole sieve; and
- (g) that are of subspecies other than *Zea mays indentata* or *Zea mays indurata*.

Provided that –

- (i) irregularity of shape and size of maize kernels shall not affect the grading thereof;
- (ii) chipped or cracked maize kernels or pieces of maize kernels which are in a sound condition and which appear in a sample of maize, but which do not pass through a 6.35 mm round-hole sieve, shall not be regarded as defective maize kernels under these regulations.”

### FOREIGN MATTER

The term “foreign matter” means all matter above the sieve other than maize, glass, stones, coal, dung or metal.

### OTHER COLOUR

“Other colour maize kernels” in relation to -

- (a) white maize, means maize kernels or pieces of maize kernels of which the endosperm as a result of genetic (characteristics) composition have another colour than white, excluding pinked maize kernels;
- (b) yellow maize, means maize kernels or pieces of maize kernels of which the endosperm as a result of genetic (characteristics) composition have another colour than yellow.

## COMBINED DEVIATION

The term “combined deviation” means the sum of defective kernels (above and below the 6.35 mm sieve), foreign matter and other colour kernels.

## PINKED KERNELS

The term “pinked maize kernels” means kernels and pieces of kernels of white maize of which the pericarp or part thereof is shaded red or pink in colour. The specification, according to the Grading Regulations for classes 1 to 3 of white maize is a maximum of 12%. No specification for pinked kernels in yellow maize according to the Grading Regulations.

## FUNGAL INFECTION

Kernels which are mouldy (fungi infected) are reported as defective kernels according to the grading regulations.

“Mouldy” means kernels and pieces of kernels that –

- (a) are visibly infected by fungi and are characterised by black, blue, green, yellow or white fungi growth anywhere on the kernel, or are characterised by fungi growth underneath the bran layer of the kernel;
- (b) are infected by ear-rot and are characterised by red, pink or brown discolorations. The kernel are partially to completely infected.

All samples were also inspected for the visual symptoms of *Diplodia* and *Fusarium* infection, which were reported separately.

*Fusarium spp* infections are localized on the cob with discoloured maize kernels, which become reddish (light pink to lilac).

*Diplodia maydis* normally rots the entire maize cob and infected maize kernels are recognized by a light ash colour to black colour that appears at the germ and can infest the whole kernel.

The % Cobrot reported, represents the percentage maize kernels that are both *Fusarium* and *Diplodia* infected.

## WATER DAMAGED MAIZE KERNELS

“Water damaged maize kernels” means maize kernels with a light yellow shine from the tip cap in a band around the maize kernel.

## USA GRADING

USA grading was determined in accordance with the American Grading Regulations (United States Department of Agriculture).

The US grading system makes provision for three

classes of maize/corn based on colour, namely Class White corn, Class Yellow corn and Class Mixed corn. Each class is divided into five U.S. numerical grades (Nos. 1 to 5) and U.S. Sample Grade. US No.1 is the most desirable grade followed by No. 2 down to sample grade.

Description of deviations relating to USA grading:

## DAMAGED KERNELS

Kernels and pieces of corn kernels that are badly ground-damaged, badly weather-damaged, diseased, frost-damaged, germ-damaged, heat-damaged, insect-bored, mould-damaged, sprout-damaged or otherwise materially damaged.

## HEAT DAMAGED KERNELS

Kernels and pieces of kernels which are materially discoloured and damaged by heat, including material discolouration caused by artificial drying methods.

## BROKEN CORN AND FOREIGN MATERIAL

Broken corn is all matter that passes readily through a 12/64-inch (4.76 mm) round-hole sieve and over a 6/64-inch (2.38 mm) round-hole sieve according to procedures prescribed in Federal Grain Inspection Service (FGIS) instructions.

Foreign material is all matter that passes readily through a 6/64-inch round-hole sieve and all matter other than corn that remains on top of the 12/64-inch round-hole sieve after sieving according to procedures prescribed in FGIS instructions.

Broken corn and foreign material is all matter that passes readily through a 12/64-inch round-hole sieve and all matter other than corn that remains in the sieved sample after sieving according to procedures prescribed in FGIS instructions.

## BUSHEL WEIGHT

The specific mass (or grain density) of maize/corn (expressed as test weight or bushel weight) is a quality characteristic which is important to some maize consumers and is applied as a grading factor in the USA grading regulations.

Test weight is the weight per Winchester bushel (2,150.42 cubic inches) as determined using an approved device according to procedures prescribed in FGIS instructions.

## OTHER COLOUR

Maize samples are deemed to be Class mixed corn when maize kernels of another colour exceeds 2% for white maize and 5% for yellow maize.

## PHYSICAL CHARACTERISTICS

### TEST WEIGHT

Bushel weight in pounds (lbs) was determined on the maize crop samples and the results converted to test weight reported in kilogram/hectoliter (kg/hl), by multiplication with a factor of 1.2872.

The specific mass (or grain density) of maize expressed as test weight is influenced by amongst other, factors like cultivar, moisture content, foreign matter, other grain and damaged kernels like insect damaged and immature kernels.

### HUNDRED (100) KERNEL MASS - INDUSTRY ACCEPTED METHOD 001

100 kernel mass is the weight in grams of one hundred whole maize kernels and provides a measure of grain size and density. The results are reported on an "as is" basis.

### KERNEL SIZE - INDUSTRY ACCEPTED METHOD 017

Kernel size is important to the sophisticated starch manufacturing industry. Kernels that are too small hamper the separation of kernel fractions in the wet milling process. The result is a lower starch yield. A mixture of small and large kernels causes additional problems, as homogeneous steeping cannot be achieved. On the other hand, very large kernels can also cause problems since the ratio between volume and mass is unfavourable to proper steeping.

The dry milling industry also prefers fairly larger maize kernels. However, uniform kernel size is of particular importance to this industry, since too large kernels create problems especially when mixed with smaller kernels.

Kernel size is less important to the animal feed manufacturing industry. Larger kernels are nevertheless preferred, as small kernels are easily lost during the screening stage of processing. The determination of kernel size comprises the sieving of a 100 g representative whole maize sample through both 8 mm and 10 mm round-hole grading sieves, routinely used in the seed industry.

### BREAKAGE SUSCEPTIBILITY - INDUSTRY ACCEPTED METHOD 007

Maize is normally cleaned before processing. In the cleaning process, broken kernels are removed together with other impurities, causing losses. Broken kernels are further broken during handling, resulting in excessive grain dust being generated. This creates the potential for dust explosions, health hazards, hygiene problems, etc. Maize containing

a high percentage of broken kernels is more prone to insect infestation and is subject to general deterioration.

In the modern dry milling industry, maize is cleaned first and then conditioned by dampening before the germ is removed. Broken kernels cause many problems during these stages of processing. Broken kernels can also lead to a lower extraction of the so-called high-quality products, like samp and maize grits. The presence of many broken kernels causes problems with the fibre and fat content of maize products (e.g. the various grades of maize meal), because the quantity of germ required to be returned to the milled endosperm cannot be determined accurately.

In the wet milling process broken kernels steep more rapidly than whole kernels and by the time the whole kernels have been sufficiently steeped, the broken kernels have been over-steeped, causing an ineffective separation of protein and starch.

In the livestock feed industry breakability is not an important quality characteristic, except for dust and hygiene reasons.

All samples were subjected to a breakage susceptibility test. After the sample of whole maize kernels was propelled in a Stein Breakage tester for 4 minutes, the fractions below the 6.35 mm and 4.75 mm sieves were collected and the percentages broken kernels smaller than (<) 6.35 mm and < 4.75 mm determined.

### STRESS CRACKS - INDUSTRY ACCEPTED METHOD 006

Stress cracks are determined by visual inspection of a certain amount of whole maize kernels examined on top of a light box for small internal cracks in the endosperm. Some kernels may even have two or more internal cracks. Any form of stress may cause internal cracks, for example rapid moisture loss in the field, during harvest or during drying.

### MILLING INDEX - INDUSTRY ACCEPTED METHOD 015

Milling Index is an indication of the milling abilities and milling quality of maize kernels where a higher milling index means a higher extraction of the high-grade and most profitable products like samp, maize rice and maize grits (degermed products) that are manufactured from the corneous part of the endosperm.

The milling index is a model developed on the Foss NIT Infratec 1241- Generation 3 Standard Version Grain Analyser where the NIT spectra were

modelled against the Roff milling fractions. In the previous seasons (until 2015/16) the Milling index of the samples were determined with the calibration model developed by the Grain Crops Institute of the ARC. With this model, the average milling index of a sample with good milling characteristics is about 95 with a variation of about 55 (low milling quality) to about 115 (very good milling quality).

The SAGL was tasked by the Maize Trust to develop a new model for Milling Index using samples from maize cultivar trials supplied by the ARC-GCI and by commercial seed breeders over four seasons (from 2012/13 onward). The trials included a range of hardness levels. The New Milling Index (NMI) that was developed is similar to the original ARC formula but on a 14% moisture basis, and with the constants removed. The NMI model has improved precision compared to the older version, due to the almost tenfold increase in the number of samples used to build the calibration model.

During the fifth year, samples of commercial hybrids, selected imported maize samples and outlier samples from the 2014/15 and 2015/16 seasons were included to develop a robust model with the assistance of Foss to produce accurate results. The latest version of the improved new model, SAGL Milling Index 2021, includes two parameters, SAGL Milling Index (SAGL MI) as well as Grit Yield All (GYA).

SAGL MI indicates the relative ratio of total hard endosperm products (B2 grits, B3 fine grits and B3 coarse grits) to offal products (B1 fine flour and total chop/bran) as determined on a Roff mill and used for calibration of the NIT. It is expressed as a dimensionless index value according to the following scale:

SAGL MI	<40	40-60	60-80	80-100	>100
Description	Soft	Medium	Moderately hard	Hard	Very hard

GYA is defined as the sum of the mass fractions of the Roff B2 grits, B3 fine grits and B3 coarse grits fractions expressed as a mass percentage of the total mass of the whole maize before milling. GYA is linearly correlated with the SAGL MI and indicates the true amount of total hard endosperm that can be extracted from the maize during Roff milling. The NIT calibration value for GYA provides this estimate directly from the whole maize without need for further milling tests. GYA is also reported on a 14% moisture base.

The 2020/21 season maize samples were measured with the NIT on the SAGL Milling Index 2021 model.

## MILLING OF MAIZE ON ROFF MAIZE MILL - INDUSTRY ACCEPTED METHOD 013

The Roff 150 Series maize mill is used to mill representative samples of 500 g. The mill is pre-set to the following specifications: Break 1 roll nip - 0.3 mm, Break 2 roll nip - 0.18 mm and Break 3 roll nip - 0.08 mm. These settings are according to the specifications in the method developed by the ARC Grain Crops Institute. Every mill has three separations, namely chop, grits and maize meal. The grits from Break 1 are transferred to the Break 2 rolls and the grits from Break 2 are transferred to Break 3 rolls.

The following fractions are weighed and determined as percentages: Break 1 meal, Break 2 meal, Break 3 meal and Break 3 grits. Break 1, 2 and 3 chop are combined and then weighed for determination of % Chop. Break 3 grits are weighed for determination of % Grits. The percentage extraction total meal is determined as the sum of the percentages Break 1, 2 and 3 meal as well as the % Grits.

## WHITENESS INDEX - INDUSTRY ACCEPTED METHOD 004

Whiteness index of white maize meal was determined with the HunterLab ColorFlex 45°/0°. Whiteness is associated with a region or volume in colour space in which objects are recognized as white. The degree of whiteness is measured by the degree of departure of the object from a perfect white. The higher the whiteness index value, the whiter the sample.

Whiteness index was done on unsifted and sifted maize meal obtained from Break 2 and 3 of the Roff mill. The sifted samples were obtained by sieving the unsifted samples through a 300 µm sieve. The fractions on top and below the sieve were then combined to result in sifted samples that contain 87% of maize meal > 300 µm and 13% of maize meal < 300 µm.

## NUTRITIONAL VALUES

The moisture, fat, protein, starch and crude fibre contents are measured with an Infratec 1241 - Generation 3 Standard Version Whole Grain Analyser. The measurements are based on the fact that the constituents to be measured in the grain, absorb electromagnetic radiation in the near-infrared region of the spectrum. Since the Infratec 1241 Grain Analyser uses transmission absorption, the test is done on intact maize kernels. The results are reported on a dry (moisture free) basis.

With the assistance of Foss, a calibration for crude fibre content was developed during the 2017/18 season. The calibration on the Infratec 1241 Grain Analyser (NIT), is updated annually by Foss using NIT spectra and international primary chemical method results of maize crop quality samples from the specific season under discussion, provided by SAGL.

The chemical methods used to check the calibration are:

- Moisture on whole maize kernels: Sample is dried in an oven for 72 hours at 103 °C (AACCI 44-15.02)
- Moisture on milled maize: Sample is dried in an oven for 1 hour at 130 °C (AACCI 44-15.02)
- Crude fat: Petroleum ether extraction (Soxhlet) method (In house method 024)
- Crude protein: Dumas (Leco) method (AACCI 46-30.01)
- Starch: Hydrochloric Acid dissolution method (Polarimeter) (In house method 019)
- Crude fibre: In-House method 031, based on AACCI method 32-10.01 using the Velp FIWE Advance fibre AutoExtractor.

The results obtained by the Infratec 1241 Grain Analyser (NIT) on the 2020/21 season's samples, were checked by analysing every tenth sample by means of the primary methods.

## MYCOTOXIN ANALYSES

Mycotoxins are fungal metabolites, toxic to animals and humans, that are produced by moulds commonly found in almost all types of grain.

350 of the 1 000 maize crop samples were tested for Aflatoxin B<sub>1</sub>, B<sub>2</sub>, G<sub>1</sub>, G<sub>2</sub>, Fumonisin B<sub>1</sub>, B<sub>2</sub> and B<sub>3</sub>, Deoxynivalenol, 15-ADON, HT-2 Toxin, T-2 Toxin, Zearalenone and Ochratoxin A by means of a multi-mycotoxin screening method (In house method 026) using UPLC - MS/MS.

Limit of quantitation (LOQ) means the lowest concentration level that can be quantified with acceptable precision and accuracy by the mass spectrometer. A concentration measured below the LOQ is reported as <LOQ.

Limit of detection (LOD) is the lowest concentration level that can be detected but not quantified and is 50% of the LOQ of each mycotoxin. A concentration measured below the LOD is reported as not detected (ND).

## GMO (GENETICALLY MODIFIED ORGANISMS)

The EnviroLogix QuickComb kit for bulk grain was used to quantitatively determine the presence of genetically modified maize. The kit is designed to extract and detect the presence of certain proteins at the levels typically expressed in genetically modified bulk maize grain. The procedure prescribed in the EnviroLogix - QuickScan Instruction Manual, latest edition was followed. Results were scanned and interpreted quantitatively with the EnviroLogix QuickScan system.

100 crop samples were tested for Cry1Ab, Cry2Ab and CP4 EPSPS modified maize. Cry1Ab protein in maize is produced from a gene derived from *Bacillus thuringiensis* (Bt).

Cry1Ab	MON810	Tradename/Brand
Cry1Ab	MON810 Cry1A.105 Bt11	YieldGard®
Cry2Ab	MON89034	<i>in</i> SmartStax™
CP4 EPSPS	NK603	Roundup Ready®







## CERTIFICATE OF ACCREDITATION

*In terms of section 22(2) (b) of the Accreditation for Conformity Assessment, Calibration and Good Laboratory Practice Act, 2006 (Act 19 of 2006), read with sections 23(1), (2) and (3) of the said Act, I hereby certify that:-*

### SOUTHERN AFRICAN GRAIN LABORATORY NPC

Co. Reg. No.: 1997/018518/08

Facility Accreditation Number: **T0116**

is a South African National Accreditation System accredited facility provided that all conditions and requirements are complied with

This certificate is valid as per the scope as stated in the accompanying schedule of accreditation, Annexure "A", bearing the above accreditation number for

### CHEMICAL AND PHYSICAL ANALYSIS

The facility is accredited in accordance with the recognised International Standard

**ISO/IEC 17025:2017**

The accreditation demonstrates technical competency for a defined scope and the operation of a quality management system

While this certificate remains valid, the Accredited Facility named above is authorised to use the relevant accreditation symbol to issue facility reports and/or certificates



---

Mr R Josias

Chief Executive Officer

Effective Date: 01 November 2019  
Certificate Expires: 31 October 2024



**ANNEXURE A**  
**SCHEDULE OF ACCREDITATION**

Facility Number: **T0116**

**Permanent Address of Laboratory:**

Southern African Grain Laboratories NPC  
Agri-Hub Office Park - Grain Building  
477 Witherite Road  
The Willows  
Pretoria  
0040

**Technical Signatories:**

Ms J Nortje (All Methods excl. In-house method 029)  
Ms M Bothma (All Chemical Methods)  
Ms A de Jager (Nutrients & Contaminants Methods)  
Ms W Louw (In-house Methods 001, 002, 003, 010 & 026)  
Ms D Moleke (Rheological Methods)  
Mrs H Meyer (All Chemical, Nutrients and Contaminants & Grading Methods)  
Ms J Kruger (All Chemical Methods)  
Ms M Motlanthe (In-house Methods 001, 003 & 026)  
Mr B van Der Linde (Grading)  
Ms M Ramare (All Chemical Methods Excl. In-House Method 012 and SOP MC23)  
Ms T de Beer (Rheological Methods)

**Postal Address:**

Postnet Suite # 391  
Private Bag X1  
The Willows  
0041

**Nominated Representative:**

Mrs H Meyer

**Tel:** (012) 807-4019

**Issue No.:** 32

**Fax:** N/A

**Date of Issue:** 19 November 2021

**E-mail:** hannelien.meyer@sagl.co.za

**Expiry Date:** 31 October 2024

<b>Material or Products Tested</b>	<b>Type of Tests / Properties Measured, Range of Measurement</b>	<b>Standard Specifications, Techniques / Equipment Used</b>
<b>CHEMICAL</b>		
Ground Barley	Moisture (Oven Method)	Analytical EBC Method 3.2, latest Edition (2 hour; 130 <sup>0</sup> C)
Cereal and cereal products specifically-wheat, rice, (hulled paddy), barley, millet, rye, and oats as grain, semolina and flour	Moisture (Oven Method)	ICC Std No.110/1, Latest Edition (90 min; 130 <sup>0</sup> C) (2 hour; 130 <sup>0</sup> C)
Flour, semolina, bread, all kind of grains and cereal products and food products (except those that are sugar coated)	Moisture (Oven Method)	AACCI 44-15.02, Latest Edition (1 hour; 130 <sup>0</sup> C) (72 hour; 103 <sup>0</sup> C)

Facility Number: T0116

Maize Grits	Moisture (Oven Method)	Analytical EBC Method 6.2.2, latest edition (4 hours, 130 <sup>0</sup> C)
Animal feed, Plant tissue and Sunflower (Milled)	Moisture (Oven Method)	AgriLASA 2.1, Latest Edition (5 hours, 105 <sup>0</sup> C)
All flours, cereal grains, oilseeds and animal feeds	Nitrogen and protein (Combustion method - Dumas)	AACCI 46-30.01, Latest Edition
Cereal based food stuff	Dietary fibres (Total)	In-house method 012
Food stuff and feeds	Carbohydrates (by difference) (calculation) Energy value (calculation) Total digestible nutritional value (calculation)	SOP MC 23
Food Stuff and feeds	Determination of Ash	In-house method 011
Wheat Kernels	Moisture (Oven Method)	Government Gazette Wheat Regulation, Latest Edition (72 hour, 103 <sup>0</sup> C)
Flours of grains e.g. barley, oats, triticale, maize, rye, sorghum and wheat; oilseeds like soybeans and sunflower, feeds and mixed feeds and foodstuffs	Crude fat (Ether extraction by Soxhlet)	In-house method 024
Meal and flour of wheat, rye, barley, other grains, starch containing and malted products	Falling number	ICC Std 107/1, Latest Edition
<b>NUTRIENTS AND CONTAMINANTS</b>		
Vitamin fortified food and feed products and fortification mixes grain based	Vitamin A as all trans Retinol (Saponification) (HPLC)	In-house method 001
	Thiamine Mononitrate (HPLC) Riboflavin (HPLC) Nicotinamide (HPLC) Pyridoxine Hydrochloride (HPLC)	In-house method 002
	Folic Acid (HPLC)	In-house method 003
	Total Sodium (Na) Total Iron (Fe) Total Zinc (Zn)	In-house method 010

Facility Number: T0116

Yeast and Bread	Vitamin D <sub>2</sub> (HPLC)	In-House method 029
Food and feed	Multi-Mycotoxin: -Aflatoxin G <sub>1</sub> , B <sub>1</sub> , G <sub>2</sub> , B <sub>2</sub> and total -Deoxynivalenol (DON), 15-ADON -Fumonisin B <sub>1</sub> , B <sub>2</sub> , B <sub>3</sub> -Ochratoxin A -T2, HT-2 - Zearalenone	In-house method 026
<b>GRADING</b>		
Maize	Defective kernels (White maize/ yellow maize)	Government Gazette Maize Regulation, Latest Edition
Cereal as grains (Wheat, barley, rye and oats)	Hectolitre mass (Kern222)	ISO 7971-3, Latest edition
Wheat	Screenings	Government Gazette Wheat Grading Regulation, Latest Edition
<b>RHEOLOGICAL</b>		
Wheat flour	Alveograph (Rheological properties)	ICC Std.121, Latest Edition
Flours	Farinograph (Rheological properties)	AACCI 54.02, Latest Edition (Rheological behaviour of flour Farinograph: Constant Flour Weight procedure)
Hard, soft and durum wheat (flour and whole wheat flour)	Mixograph (Rheological properties)	Industry accepted method 020 (Based on AACCI 54-40.02, Latest Edition Mixograph Method)

Original Date of Accreditation: 01 November 1999

ISSUED BY THE SOUTH AFRICAN NATIONAL ACCREDITATION SYSTEM

  
\_\_\_\_\_  
Accreditation Manager

**GOVERNMENT NOTICES  
GOEWERMENTSKENNISGEWINGS**

**DEPARTMENT OF AGRICULTURE  
DEPARTEMENT VAN LANDBOU**

No. R. 473

8 May 2009

AGRICULTURAL PRODUCT STANDARDS ACT, 1990  
(ACT No. 119 OF 1990)

**REGULATIONS RELATING TO THE GRADING, PACKING AND MARKING OF MAIZE  
INTENDED FOR SALE IN THE REPUBLIC OF SOUTH AFRICA**

The Minister of Agriculture, acting under section 15 of the Agricultural Product Standards Act, 1990 (Act No. 119 of 1990),

- (a) made the regulations in the Schedule; and
- (b) determined that the said regulations shall come into operation on date of publication.

**SCHEDULE**

**Definitions**

1. In these regulations any word or expression to which a meaning has been assigned in the Act shall have that meaning and, unless the context otherwise indicates -

"bag" means a bag manufactured from -

- (a) jute or phormium or a mixture of jute and phormium; or
- (b) polypropylene that complies with SABS specification CKS632;

"bulk container" means any vehicle or container in which bulk maize is stored or transported-

"consignment" means -

- (a) a quantity of maize of the same class, which belongs to the same owner, delivered at any one time under cover of the same consignment note, delivery note or receipt note, or delivered by the same vehicle or bulk container, or loaded from the same bin of a grain elevator or from a ship's hold; or
- (b) in the case where a quantity referred to in paragraph (a), is subdivided into different grades, each such quantity of each of the different grades;

"coffee stained maize kernels" means maize kernels with a shiny brown colour that occurs anywhere on the pericarp of the maize kernel;

"container" means a bag or a bulk container;

"defective maize kernels" means maize kernels and pieces of maize kernels-

- (a) that are shrivelled, obviously immature, frost-damaged, heat damaged, water damaged, mouldy or chalky;
- (b) that are discoloured by external factors such as water and sun: Provided that discoloration on both sides of the maize kernel limited to less than a quarter from the bottom tip of the maize kernel shall not be considered as defective; oxidation stained maize kernels; coffee stained maize kernels; and pinked maize kernels shall not be considered as defective;
- (c) that have sprouted, including kernels of which the shoot (plumule) in the germ is visibly discoloured;

- (d) that have cavities in the germ or endosperm caused by insects or rodents;
- (e) that are visibly soiled (smeared) or contaminated by smut, fire, soil, smoke or coal-dust;
- (f) all matter that can pass through the 6,35 mm round-hole sieve; and
- (g) that are of subspecies other than *Zea mays indentata* or *Zea mays indurata*.

Provided that:

- (i) irregularity of shape and size of maize kernels shall not affect the grading thereof;
- (ii) chipped or cracked maize kernels or pieces of maize kernels which are in a sound condition and which appear in a sample of maize, but which do not pass through a 6,35 mm round-hole sieve, shall not be regarded as defective maize kernels under these regulations;

**"discoloured maize kernels"** means maize kernels that are as a result of environmental conditions more than 25% discoloured on both sides of the kernel, excluding coffee stained maize kernels, oxidation stained maize kernels and pinked maize kernels;

**"foreign matter"** means all matter above the sieve other than maize, glass, stone, coal, dung or metal;

**"frost damaged"** means maize kernels that are covered with wrinkles on both sides of the kernel to the crown and have a pearl-like appearance. Maize kernels of which the bran is flaking is considered frost damaged if signs of frost damage are present;

**"heat damaged"** means kernels that are as a result of external heat or internal fermentation affected with excess moisture and have at least one of the following characteristics:

- (a) Kernels or pieces of kernels that are amber, brown, dark-brown or black discoloured;
- (b) Kernels of which the germ has dark-brown to black discoloration;

**"insect"** in relation to maize, means any live insect which is injurious to stored grain, irrespective of the stage of development of the insect;

**"maize"** means the threshed kernels or pieces of kernels of the plants of *Zea mays indurata* and *Zea mays indentata* or one or more crossings of the two types;

**"mouldy"** means kernels or pieces of kernels that-

- (a) are visibly infected by fungi and are characterised by black, blue, green, yellow or white fungi growth anywhere on the kernel, or are characterised by fungi growth underneath the bran layer of the kernel;
- (b) are infected by ear-rot and are characterised by red, pink or brown discolorations. The kernels are partially to completely infected;

**"other colour maize kernels"** in relation to -

- (a) white maize, means maize kernels or pieces of maize kernels of which the endosperm as a result of genetic (characteristics) composition have another colour than white, excluding pinked maize kernels;
- (b) yellow maize, means maize kernels or pieces of maize kernels of which the endosperm as a result of genetic (characteristics) composition have another colour than yellow;

"oxidation stained maize kernels" means maize kernels with a shiny light brown colour that are discoloured from the crown and not from the tip cap;

"pinked maize kernels" means kernels and pieces of kernels of white maize of which the pericarp or part thereof is shaded red or pink in colour;

"poisonous seeds" means seeds or part of seeds of plant species that may in terms of the Foodstuffs, Cosmetics and Disinfectants Act, 1972 (Act No. 54 of 1972) represent a hazard to human or animal health when consumed, including seeds of *Argemone mexicana*, *Convolvulus* spp., *Crotalaria* spp., *Datura* spp., *Ipomoea* spp., *Lolium temulentum*, *Ficus communis* or *Xanthium* spp;

"shrivelled or obviously immature maize kernels" means maize kernels with a thin and shrunken appearance;

"sprouted maize kernels" means maize kernels which have sprouted so far that developing roots and/or sprouts are clearly visible, or the shoot (plumule) in the germ is visibly discoloured;

"the Act" means the Agricultural Product Standards Act, 1990 (Act No. 119 of 1990);

"the 6,35 mm round-hole sieve" means a sieve-

- (a) with a flat metal sheet bottom of 1,0 mm thickness perforated with round holes of 6,35 mm ( $\pm 0,05$  mm), in diameter that are arranged with the centres of the holes at the points of intersection of an equilateral triangular grid with a pitch of 8 mm;
- (b) of which the upper surface of the bottom is smooth;
- (c) the frame of which is at least 40 mm high;
- (d) with the inner width of at least 200 mm and the inner length of at least 300 mm, or, in the case of a circular sieve, the inner diameter of at least 278 mm;
- (e) with a minimum area of 600cm<sup>2</sup> and a maximum of 750cm<sup>2</sup>; and
- (f) that fits onto a tray with a solid bottom and must be at least 20mm above the bottom of the tray; and

"water damaged maize kernels" means maize kernels with a light yellow shine from the tip cap in a band around the maize kernel.

#### **Scope of regulations**

2. These regulations are the minimum standards applicable to maize that are destined for sale in the Republic of South Africa but does not include -

- (a) maize in retail quantities; and
- (b) maize for seed production purposes.

#### **Restrictions on sale of maize**

3. (1) No person shall sell maize in the Republic of South Africa -
  - (a) unless the maize is sold according to the classes set out in regulation 4;

- (b) unless the maize complies with the standards for the class concerned set out in regulation 5;
  - (c) unless the maize complies with the grades of maize and the standards for grades, where applicable, set out in regulations 6 and 7 respectively;
  - (d) unless the maize is packed in accordance with the packing requirements set out in regulation 8;
  - (e) unless the containers or sale documents, as the case may be, are marked in accordance with the marking requirements set out in regulation 9; and
  - (f) if such maize contains a substance that renders it unfit for human consumption or for processing into or utilisation thereof as food or feed.
- (2) The Executive Officer may grant written exemption, entirely or partially to any person on such conditions as he or she may deem necessary, from the provisions of subregulation 1: Provided that such exemption is done in terms of section 3 (1) (c) of the Act.

#### PART I QUALITY STANDARDS

##### *Classes of maize*

4. The classes of Maize shall be -
- (a) Class White Maize;
  - (b) Class Yellow Maize; and
  - (c) Class Other Maize.

##### *Standards for classes of maize*

5. (1) A consignment of maize shall be classified as Class White Maize if -
- (a) subject to the allowable deviation in respect of other colour maize kernels that apply to the different grades of white maize, it consists of maize the endosperm of which is by nature white in colour; and
  - (b) it complies with the standards for one of the grades of white maize set out in regulation 7.
- (2) A consignment of maize shall be classified as Class Yellow Maize if -
- (a) subject to the allowable deviation in respect of other colour maize kernels that apply to the different grades of yellow maize, it consists of maize the endosperm of which is by nature yellow in colour; and
  - (b) it complies with the standards for one of the grades of yellow maize set out in regulation 7.
- (3) A consignment of maize shall be classified as Class Other Maize if the consignment does not comply with the standards for Class White Maize or Class Yellow Maize.



**Grades of maize**

6. (1) Maize of the Class White Maize shall be graded as WM1, WM2 or WM3.
- (2) Maize of the Class Yellow Maize shall be graded as YM1, YM2 or YM3.
- (3) No grades are determined for Class Other Maize.

**Standards for grades of Class White Maize and Class Yellow Maize**

7. All grades of maize -
  - (a) shall be free from a musty, sour or other undesired odour;
  - (b) shall be free from glass, metal, coal or dung;
  - (c) shall be free from a substance which renders it unfit for human consumption or for processing into or utilisation thereof as food or feed;
  - (d) shall be free from insects;
  - (e) shall be free from stones which cannot pass through the 6,35 mm round-hole sieve;
  - (f) shall contain not more than one gram of stones, which can pass through the 6,35 mm round-hole sieve, per 10 kg;
  - (g) shall contain not more poisonous seeds than permitted in terms of the Foodstuffs, Cosmetics and Disinfectants Act, 1972 (Act No. 54 of 1972);
  - (h) shall have a moisture content of not more than 14 per cent; and
  - (i) shall not exceed the maximum percentage of permissible deviation as determined in the table in the Annexure for each grade.

**PART II  
PACKING AND MARKING REQUIREMENTS****Packing requirements**

8. Maize of different classes and grades shall be packed in different containers.

**Marking requirements**

9. Each container or the accompanying sales document of a consignment of maize shall be marked or endorsed with -
  - (a) the class of the maize;
  - (b) the grade, in the case of Class White Maize or Class Yellow Maize; and

**PART III  
SAMPLING****Obtaining sample**

10. (1) A sample of a consignment of maize shall -

- (a) in the case of maize delivered in bags and subject to regulation 11, be obtained by sampling at least ten per cent of the bags, chosen from that consignment at random, with a bag probe: Provided that at least 25 bags in a consignment shall be sampled and where a consignment consists of less than 25 bags, all the bags in that consignment shall be sampled; and
  - (b) in the case of maize delivered in bulk and subject to regulation 10, be obtained by sampling that consignment throughout the whole depth of the layer, in at least six different places, chosen at random in that bulk quantity, with a bulk sampling apparatus.
- (2) The collective sample obtained in subregulation (1) (a) or (b) shall -
- (a) have a total mass of at least 10 kg; and
  - (b) be thoroughly mixed by means of dividing before further examination.
- (3) If it is suspected that the sample referred to in subregulation (1)(a) is not representative of that consignment, an additional five per cent of the remaining bags, chosen from that consignment at random, shall be emptied into a suitable bulk container and sampled in the manner contemplated in subregulation (1)(b).
- (4) A sample taken in terms of these regulations shall be deemed representative of the consignment from which it was taken.

***Sampling if contents differ***

11. (1) If, after an examination of the maize taken from different bags in a consignment in terms of regulation 10(1), it appears that the contents of those bags differ substantially -
- (a) the bags concerned shall be placed separately;
  - (b) all the bags in the consignment concerned shall be sampled in order to do such separation; and
  - (c) each group of bags with similar contents in that consignment shall for the purposes of these regulations be deemed to be a separate consignment.
- (2) If, after the discharge of a consignment of maize in bulk has commenced, it is suspected that the consignment could be of a class or grade other than that determined by means of the initial sampling, the discharge shall immediately be stopped and the part of the consignment remaining in the bulk container, as well as the grain that is already in the collecting tray, shall be sampled anew with a bulk sampling apparatus or by catching at least 20 samples at regular intervals throughout the whole offloading period with a suitable container from the stream of grain that is flowing in bulk.

***Working sample***

12. A working sample shall be obtained by dividing the representative sample of the consignment according to the ICC 101/1 method.

**PART IV  
DETERMINATION OF OTHER SUBSTANCES**

***Determination of undesirable odours and harmful substances***

13. A sample of a consignment of maize shall be sensorial assessed or chemically analysed in order to determine -

- (a) whether it has a musty, sour or other undesirable odour: Provided that a working sample of unscreened maize that is ground in a grain mill to a fine meal may be used for the determination concerned; and
- (b) whether it contains a substance that renders the maize unfit for human consumption or for processing into or for utilisation as food or feed.

***Determination of glass, metal, coal, dung, stone, poisonous seed and insect content***

14. A consignment of maize shall be sensorial assessed and a sample of that consignment shall be sensorial assessed and sorted by hand in order to determine whether the sample contains glass, metal, coal, dung, insects, stones and poisonous seeds.

***Determination of percentage of foreign matter***

15. The percentage of foreign matter in a consignment of maize shall be determined as follows:
- (a) Obtain a working sample with a mass of at least 150g from the sample of the consignment.
  - (b) Remove all foreign matter from the working sample and determine the mass thereof.
  - (c) Express the mass thus determined as a percentage of the total mass of the working sample.
  - (d) Such percentage shall represent the percentage of foreign matter in the consignment concerned.

**PART V  
MAIZE KERNELS**

***Determination of percentage of defective maize kernels***

16. The percentage of defective maize kernels in a consignment of maize shall be determined as follows:
- (a) Obtain a working sample with a mass of at least 150g from the sample of the consignment.
  - (b) Place the working sample on the 6, 35 mm round-hole sieve and screen the sample by moving the sieve 20 strokes to and fro, alternately away from and towards the operator of the sieve. Move the sieve, which rests on a table or other suitable smooth surface, 250 mm to 460 mm away from and towards the operator with each stroke. The prescribed 20 strokes must be completed within 20 to 30 seconds.
  - (c) Determine the mass of the matter that has passed through the sieve and express it as a percentage of the mass of the working sample.
  - (d) Remove all defective maize kernels from that part of the working sample remaining on the sieve and determine the mass thereof.
  - (e) Express the mass as a percentage of the mass of the working sample.
  - (f) Calculate the sum of the masses determined in terms of paragraphs (c) and (d).
  - (g) Express the combined mass calculated in terms of paragraph (f) as a percentage of the mass of the working sample.
  - (h) In the case of yellow maize the percentage obtained -

- (i) in terms of paragraph (c), represents the percentage of defective maize kernels in the consignment concerned, which can pass through the 6,35 mm round-hole sieve; and
  - (ii) in terms of paragraph (e), represents the percentage of defective maize kernels in the consignment concerned, which can not pass through the 6,35 mm round-hole sieve.
- (i) In the case of white maize, the percentage obtained in terms of paragraph (g) represents the percentage of defective maize kernels in the consignment concerned.

***Determination of percentage of other colour maize kernels***

17. The percentage of other colour maize kernels in a consignment of maize shall be determined as follows:

- (a) Obtain a working sample with a mass of at least 150g from the sample of the consignment.
- (b) Remove all other colour maize kernels from the working sample and determine the mass thereof.
- (c) Express the mass thus determined as a percentage of the mass of the working sample.
- (d) Such percentage shall represent the percentage of other colour maize kernels in the consignment concerned.

***Determination of percentage of pinked maize kernels***

18. The percentage of pinked maize kernels in a consignment of maize shall be determined as follows:

- (a) Obtain a working sample with a mass of at least 150g from the sample of the consignment.
- (b) Remove all pinked maize kernels from the working sample and determines the mass thereof.
- (c) Express the mass thus determined as a percentage of the mass of the working sample.
- (d) Such percentage shall represent the percentage of pinked maize kernels in the consignment concerned.

**PART VI  
MOISTURE CONTENT**

***Determination of moisture content***

19. The moisture content of a consignment of maize may be determined according to any suitable method: Provided that the results thus obtained are in accordance with the maximum permissible deviation for a class 1 moisture meter as detailed in ISO 7700/1 based on the results of the 72 hour, 103°C oven dried method (AACC Method 44-15A).

**OFFENCE AND PENALTIES**

20. Any person who contravenes or fails to comply with any provision of these regulations shall be guilty of an offence and upon conviction be liable to a fine or imprisonment in terms of section 11 of the Act.

**ANNEXURE/AANHANGSEL  
TABLE/TABEL**

**STANDARDS FOR GRADES OF CLASS WHITE MAIZE AND CLASS YELLOW MAIZE/  
STANDAARDE VIR GRADE VAN KLAS WITMIELIES EN KLAS GEELMIELIES**

Deviation/Afwyking	Maximum permissible deviation/ Maksimum toelaatbare afwyking					
	White maize/ Witmielies			Yellow maize/ Geelmielies		
	WM1	WM 2	WM 3	YM1	YM2	YM3
1	2	3	4	5	6	7
1. Foreign matter [regulation 15]/ Vreemde voorwerpe [regulasie 15]	0,3%	0,5%	0,75 %	0,3%	0,5%	0,75%
2. Defective maize kernels, above and below the 6,35 mm round-hole sieve [regulations 16]/ <i>Gebrekkige mieliepitte, bo en onder die 6,35 mm-rondegatsif [regulasies 16]</i>	7%	13%	30%	*	*	*
3. Defective maize kernels that can pass through the 6,35 mm round-hole sieve [regulation 16(c)]/ <i>Gebrekkige mieliepitte wat deur die 6,35 mm rondegatsif kan gaan [regulasie 16(c)]</i>	*	*	*	4%	10%	30%
4. Defective maize kernels that can not pass through the 6,35 mm round-hole sieve [regulation 16(e)]/ <i>Gebrekkige mieliepitte wat nie deur die 6,35 mm-rondegatsif kan gaan nie [regulasie 16(e)]</i>	*	*	*	9%	20%	30%
5. Other colour maize kernels [regulation 17]/ <i>Mieliepitte van 'n ander kleur [regulasie 17]</i>	3%	6%	10%	2%	5%	5%
6. Deviations referred to in items 1, 3, 4 and 5 individually within the specified limits/ <i>Afwykinge in items 1, 3, 4 en 5 bedoel, gesamentlik: met dien verstande dat die afwykinge individueel binne die gespesifiseerde perke is</i>	8%	16%	30%	9%	20%	30%
7. Pinked maize kernels [regulation 18]/ <i>Verrooide mieliepitte [regulasie 18]</i>	12%	12%	12%	*	*	*

\* Not specified/Nie gespesifiseer nie.



agriculture,  
forestry & fisheries

Department:  
Agriculture, Forestry and Fisheries  
REPUBLIC OF SOUTH AFRICA

Directorate Food Safety and Quality Assurance Private Bag X258, PRETORIA, 0001 • Tel: +27 12 319 6291

**FAX COVER SHEET**

TO: Grain Silo Industry

ATTENTION: Mr. Dirk Kok

FAX: 012 348 2980

FROM: Mrs. Caroline Makobe

E-MAIL: CarolineL@nda.agric.za

TEL: (012) 319 6291

REF NO: 20/4/14/1 / Dispensation

FAX: (012) 319 6055

SERIAL NO:

NO. PAGES: 1

DATE: 15 April 2010

Subject

**INDUSTRY-WIDE DISPENSATION: AMENDMENT OF THE REGULATION RELATING TO THE GRADING, PACKING AND MARKING OF MAIZE INTENDED FOR SALE IN THE REPUBLIC OF SOUTH AFRICA.**

Please refer to the e-mail dated 29<sup>th</sup> March 2010 from Grain Silo Industry

Permission is hereby granted by the Executive Officer: Agricultural Product Standards, in terms of Section (3) of the Agricultural Products Standards Act, 1990 (Act No. 119 of 1990), to all producers, wholesalers, traders, retailers and importers of Maize to sell and import maize whereby the definition of "Foreign matter" is amended in the English version in order to align it to the Afrikaans one of the above mentioned Regulation to read as follows : **"Foreign matter" means all matter other than maize, glass, stones above the sieve, coal, dung or metal.**

This dispensation is extended further to apply to item 6 of the Annexure in the Table relating to Standards for grades of Class White and Yellow maize which is amended and replaced with the following item: Provided that all provisions of the regulations shall be complied with:

ANNEXURE/AANHANGSEL  
TABLE/TABEL


STANDARDS FOR GRADES OF CLASS WHITE MAIZE AND CLASS YELLOW MAIZE/  
STANDAARDE VIR GRADE VAN KLAS WITMIELIES EN KLAS GEELMIELIES

Deviation/Afwyking	Maximum permissible deviation/ Maksimum toelaatbare afwyking					
	White maize/ Witmielies			Yellow maize/ Geelmielies		
	WM 1	WM2	WM3	YM1	YM2	YM3
1	2	3	4	5	6	7
6. Deviations referred to in items 1, 2, 3, 4 and 5 collectively: Provided that the deviations are individually within the specified limits/Afwykinge in items 1, 2, 3,4 en 5 bedoel, gesamentlik: Met dien verstande dat die afwykinge individueel binne die gespesifiseerde perke is	8%	16%	30%	9%	20%	30%

**INDUSTRY-WIDE DISPENSATION: AMENDMENT OF THE REGULATION RELATING TO THE GRADING, PACKING AND MARKING OF MAIZE INTENDED FOR SALE IN THE REPUBLIC OF SOUTH AFRICA.**

This permission is subject to the following conditions:

- (a) All other conditions of the regulations shall be complied with.
- (b) It may be withdrawn at any time should a valid complaint be received
- (c) All producers, wholesalers, traders, retailers and importers of Maize, Indemnifies this Directorate and the Department from any detrimental effect, financially or otherwise, which may emanate as a result of this permission.
- (d) Termination date: until the regulation is reviewed and gazetted.



**EXECUTIVE OFFICER:**

**AGRICULTURAL PRODUCT STANDARDS ACT, NO. 119 OF 1990**

Copies: APIS : NPPIS North – Attention: Jimmy Mogodi

