



SOUTH AFRICAN  
Wheat Crop  
Quality Report  
2010/2011 Season

# INDEX

	Page
Introduction	1
Production and protein distribution graphs	2
Quality 2010/2011 season and wheat grades	3
Protein distribution per production area (graphs)	4
Wheat class and grade per production area (pie charts)	4
Cultivars and mycotoxins	5
Bread Wheat Grading table	6
Mycotoxin results	6
Weighted average protein, falling number, hectolitre mass and mixogram peak times per region compared over the last three seasons	7
Protein, falling number, hectolitre mass and mixogram peak time over ten seasons (graphs)	8
Regional quality	9 - 12
Regional quality weighted averages (table)	13
RSA chart and seed figures	14
South African Winter Cereal Production	15
Average yield per province (graph)	15
Comparison of Flour Quality over last four seasons	16
Correlation graphs for protein content, loaf volume and gluten content	17
SAGIS Wheat Imports / Exports 2009/2010 Season	18
SAGIS Wheat Imports / Exports 2010/2011 Season	19
SAGIS Wheat Exports and Imports per country 2009/2010 Season	20
SAGIS Wheat Exports and Imports per country 2010/2011 Season	21
SAGIS Wheat Stock Figures	22
Quality data plus rheological graphs per production region	23 - 54
Summary of RSA wheat quality of 2008/2009 and 2010/2011	55 - 56
Summary of RSA wheat quality of 2009/2010 and 2010/2011	57 - 58
Methods	59 - 61
Imported wheat quantity (2009/2010 season)	62
Average quality data of imported wheat during the 2009/2010 season table	62
Imported wheat quality compared to RSA crop quality during the 2009/2010 production season	63 - 72
Accuracy Awards	73
SANAS Certificate and Schedule of Accreditation	74 - 76
Grading Regulations for Bread Wheat, Regulation No. R. 1186 of 17 December 2010	77 - 88

Compiled and issued by the:

Southern African Grain Laboratory  
 Grain Building  
 477 Witherite Road  
 The Willows  
 Pretoria  
 SOUTH AFRICA

PostNet Suite # 391  
 Private Bag X 1  
 The Willows  
 0041

Tel: +27 (12) 807 4019  
 Fax: +27(12) 807 4160



E-mail: info@sagl.co.za  
 Website: www.sagl.co.za

# SOUTH AFRICAN

## COMMERCIAL WHEAT QUALITY FOR THE 2010/2011 SEASON

### Acknowledgements

With gratitude to:

- The Winter Cereal Trust for its financial support in conducting this survey.
- The Grain Silo Industry and its members for their cooperation in providing the samples to make this survey possible.
- The National Chamber of Milling and its members for providing samples of wheat delivered directly to the mills.

### Introduction

The final wheat production of 1 430 000 tons for the 2010/2011 season, was 27 % lower than the previous season's 1 958 000 tons. This is 26,6 % lower than the 10 year average of 1 948 591 tons (2001/2002 to 2010/2011 seasons). A total area of 558 100 hectares was utilized for wheat production. (Figures obtained from the Crop Estimates Committee).

The whole wheat protein average was 12.1 % compared to the 11.7 % of the previous season and the ten year average of 12.0 %. The average hectolitre mass was 80.3 kg/hl. The hectolitre mass of this season's samples was determined according to ISO 7971-3 by means of the Kern 222 instrument. This method was accredited by SANAS during 2010. The average mixogram peak time of 3.0 minutes was similar to the previous two season's 2.9 minutes.

The percentage of samples in this survey graded as B1 increased from 33 % the previous season to 44 % this season. The main contributing factors being the change in instrument for the hectolitre mass determination and higher protein levels. The low rainfall experienced during the planting and growing periods in parts of the production regions like the Southern Cape and Free State resulted in lower yields and higher protein levels. Preharvest sprouting and low falling numbers observed in parts of the Free State was due to late rains towards the end of October 2010.

Differences in the flour and dough qualities between the winter rainfall, summer rainfall and irrigation areas were observed as in previous seasons. The overall flour and dough quality were good.

During the harvesting season, a representative sample of each delivery of wheat is taken

A sub-sample of each of these grading samples is collected in a bin according to grade and class per silo bin at each silo. This composite bin sample is then divided and a 3 kg sample is sent to the Southern African Grain Laboratory (SAGL) for the annual wheat crop quality survey. SAGL analysed all 372 samples received, to represent the production of wheat in all the different production regions. The number of samples analysed was significantly less than in previous seasons (480 samples) due to the decrease in wheat production.

The samples were fully graded and thousand kernel mass was done. Small samples were milled on the Quadromat mill, followed by a mixograph analysis.

Cultivar identification was done on these samples and sales figures of seed sold by the commercial grain silo owners were obtained.


Composite samples were made up per class and grade for each production region and milled on the Bühler mill. Rheological tests, namely gluten, mixogram, farinogram, alveogram, extensogram and 100-gram baking tests, were then performed.

The results (as averages per region) are made available weekly on the SAGL website ([www.sagl.co.za](http://www.sagl.co.za)) as soon as the first samples are received. The hard copy reports are distributed to all interested parties and can also be downloaded from the website.

Summaries comparing the quality of the local wheat for the 2008/2009 and 2010/2011 as well as the 2009/2010 and 2010/2011 seasons are provided.


## WHEAT PRODUCTION IN THE RSA OVER THE LAST 10 SEASONS

### Wheat Production RSA



### DIFFERENCES IN THE DISTRIBUTION OF PROTEIN CONTENT OVER THE LAST 3 SEASONS

#### Distribution of protein content



## Crop quality of the 2010/2011 season

The protein graph of the wheat produced in the 2010/2011 season showed a normal distribution around the 12.0 - 12.9 % protein level (see page 2). The highest percentage of samples (31.5 %) had protein contents ranging from 12.0 - 12.9 %. The second highest percentage of 27.7 % was for protein contents 11.0 - 11.9 % and thirdly 21.0 % for 13.0 - 13.9 % protein content. The Summer rainfall area (Free State) had the highest average protein content of 12.6 % and the Winter rainfall and other Summer rainfall and Irrigation areas the lowest with 11.8 %.

The Winter (Western Cape) and Summer (Free State) rainfall areas had similar average hectolitre masses of 79.8 and 79.7 kg/hl respectively. The Irrigation (81.3 kg/hl) and Other Summer rainfall and Irrigation areas (81.6 kg/hl) also had similar average hectolitre masses but tested approximately 1.7 kg/hl higher than the Winter and Summer rainfall areas.

The weighted average thousand kernel mass of 39.5 g was 0.4 g higher than the previous season. The weighted average screenings (1.8 mm sieve) of 1.68 % was similar to the 1.63 % in the 2009/2010 season.

The weighted average falling number was 372 seconds. Twenty five samples gave falling number values of less than 250 seconds and of these, twenty two had falling number values lower than 220 seconds. These samples were mainly from the Free State (96 %).

The weighted mixogram peak time on flour from the Quadromat mill averaged 3.0 minutes, similar to the ten year average (2.9 minutes). The weighted mixogram peak time of the flour from the Bühler mill averaged 2.8 minutes.

The weighted average Bühler extraction was 74.9 %, with a weighted average Kent Jones colour of -1.8 KJ.

The farinogram had a weighted average water absorption of 63.2 % (61.0 % the previous season) and a weighted average development time of 5.5 minutes (3.5 minutes previous season). The weighted average alveogram strength was 36.2 cm<sup>2</sup> and the weighted average P/L value 1.29 (35.5 cm<sup>2</sup> and 1.17 the previous season). The weighted average extensogram strength was 97 cm<sup>2</sup> (83 cm<sup>2</sup> previous season).





The loaves baked using the 100 g straight-dough optimized bread making method, which refers to the relationship between the protein content and the bread volume, was evaluated and scored from "Excellent" to "Extremely Poor" (one sample). The average scores were "Very Good" to "Good".

## Wheat grades

The 372 representative crop samples were graded as follows: 44 % was graded B1, 25 % was graded B2, 13 % was graded B3, 6 % was graded B4, 8 % UT (Utility Grade) and 4 % COW (Class Other Wheat). The determination of the hectolitre mass according to ISO 7971-3 by means of the Kern 222 instrument (please see Methods page 59), contributed to the increase in the percentage samples graded B1, compared to the 2008/2009 and earlier seasons.


Grade B1 wheat in the Free State province amounted to 51 % (45 % the previous season) and grade B1 in other summer rainfall and irrigation areas amounted to 45 % (36 % in the previous season). In the irrigation areas 48 % (43 % in the previous season) of the wheat graded as B1 and in the Western Cape Province 33 % graded as B1 (15 % in the previous season).

# Protein distribution graphs per production area




# Wheat class and grades per production area


Number of samples per class and grade in the irrigation areas




Number of samples per class and grade in the summer rainfall area



Number of samples per class and grade in the winter rainfall area



Number of samples per class and grade in the other summer rainfall and irrigation areas



## Cultivars

In the Western Cape, SST 88 dominated the market (35 %). SST 027 (29 %) and SST 015 (26 %) were also popular cultivars. The Western Cape produced 38 % of all wheat grown in South Africa during the 2010/2011 season.

Regions 21 to 24 of the Free State were dominated by PAN 3120 (26 %) and SST 835 (23 %). Elands dominated regions 25 to 28 (40 %). Matlabas and SST 356 were also popular cultivars with 16 % and 13 % respectively.

Farmers in the Vaal and Orange River areas preferred SST 835 (34 %), Duzi (30 %) and SST 843 (15 %).

The most preferred cultivar in the North West was SST 835 (36 %), followed by SST 843 (27 %) and Duzi (8 %).

In Limpopo, Gauteng, Mpumalanga and KwaZulu-Natal, SST 835 (42 %) was the dominant cultivar followed by SST 843 (22 %), Kariega (19 %) and Duzi (10 %).

The above information was calculated from the cultivar identification done on all 372 crop samples.

## Mycotoxins

Mycotoxins are secondary metabolites produced by fungi on agricultural commodities intended for human and animal consumption. These mycotoxins are potentially dangerous to humans and animals since they are, amongst other also carcinogens. Aside from health risks, mycotoxin contamination can also reduce the value of the crops. Environmental factors such as temperature, humidity, soil and storage conditions influence toxin production.

During 2010 SAGL implemented a multi-mycotoxin screening method using UPLC-MS/MS. With this technique simultaneous quantification and confirmation of Aflatoxin G<sub>1</sub>; B<sub>1</sub>; G<sub>2</sub>; B<sub>2</sub>, Fumonisin B<sub>1</sub> and B<sub>2</sub>, Deoxynivalenol, T2-toxin, Zearalenone and Ochratoxin A are possible in one run.

Thirty samples (representing different regions as well as different classes and grades) were selected randomly for mycotoxin analyses.

All ten mycotoxins were not detected on all thirty samples tested.

# BREAD WHEAT GRADING TABLE

2010/2011

Grade	Minimum			Maximum percentage permissible deviation (m/m)									
				A	B	C	D	E	F	G	H	I	J
	Hectolitre mass, kg/hl	Falling number, seconds	Protein content, %	Heavily frost damaged kernels	Field fungi	Storage fungi	Screenings	Other grain and unthreshed ears	Gravel, stones, turf and glass	Foreign matter plus F	Heat damaged kernels	Damaged kernels plus H	Combined deviations (D+E+G+I)
Grade 1	77	220	12	5	2	0.5	3	1	0.5	1	0.5	2	5
Grade 2	76	220	11	5	2	0.5	3	1	0.5	1	0.5	2	5
Grade 3	74	220	10	5	2	0.5	3	1	0.5	1	0.5	2	5
Grade 4	72	200	9	5	2	0.5	3	1	0.5	1	0.5	2	5
Utility grade	70	150	8	10	2	0.5	10	4	0.5	3	0.5	5	10
Other Wheat	<70	<150	<8	>10	>2	>0.5	>10	>4	>0.5	>3	>0.5	>5	>10
Minimum size of working samples	1 kg	300 g clean	Apparatus instructions	25 g sifted	25 g sifted	25 g sifted	500 g unsifted	50 g sifted	100 g sifted	100 g sifted	100 g sifted	25 g sifted	-

## Mycotoxin results for the 2010/2011 season

Region	Grade	Aflatoxin µg/kg				Fumonisin µg/kg		DON µg/kg	Ochratoxin A µg/kg	Zearalenone µg/kg	T2 - Toxin µg/kg
		G <sub>1</sub>	B <sub>1</sub>	G <sub>2</sub>	B <sub>2</sub>	B <sub>1</sub>	B <sub>2</sub>				
		LOD									
		4 µg/kg	2 µg/kg	4 µg/kg	4 µg/kg	100 µg/kg	100 µg/kg	100 µg/kg	20 µg/kg	20 µg/kg	20 µg/kg
1	B1	0	0	0	0	0	0	0	0	0	0
2	UT	0	0	0	0	0	0	0	0	0	0
3	B2	0	0	0	0	0	0	0	0	0	0
4	B3	0	0	0	0	0	0	0	0	0	0
5	B2	0	0	0	0	0	0	0	0	0	0
6	B1	0	0	0	0	0	0	0	0	0	0
10	B1	0	0	0	0	0	0	0	0	0	0
11	B1	0	0	0	0	0	0	0	0	0	0
12	B1	0	0	0	0	0	0	0	0	0	0
14	B4	0	0	0	0	0	0	0	0	0	0
15	B2	0	0	0	0	0	0	0	0	0	0
16	B2	0	0	0	0	0	0	0	0	0	0
17	B1	0	0	0	0	0	0	0	0	0	0
18	B1	0	0	0	0	0	0	0	0	0	0
19	B1	0	0	0	0	0	0	0	0	0	0
20	B2	0	0	0	0	0	0	0	0	0	0
21	B2	0	0	0	0	0	0	0	0	0	0
22	B1	0	0	0	0	0	0	0	0	0	0
23	B3	0	0	0	0	0	0	0	0	0	0
24	B1	0	0	0	0	0	0	0	0	0	0
25	B1	0	0	0	0	0	0	0	0	0	0
26	B1	0	0	0	0	0	0	0	0	0	0
27	B1	0	0	0	0	0	0	0	0	0	0
28	B1	0	0	0	0	0	0	0	0	0	0
29	B1	0	0	0	0	0	0	0	0	0	0
30	COW	0	0	0	0	0	0	0	0	0	0
32	B1	0	0	0	0	0	0	0	0	0	0
34	B2	0	0	0	0	0	0	0	0	0	0
35	B1	0	0	0	0	0	0	0	0	0	0
36	B1	0	0	0	0	0	0	0	0	0	0
<b>Average</b>		<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>
<b>Number of samples</b>		<b>30</b>	<b>30</b>	<b>30</b>	<b>30</b>	<b>30</b>	<b>30</b>	<b>30</b>	<b>30</b>	<b>30</b>	<b>30</b>

Note: All results <LOD and non detected are reported as 0





LOD: Limit of detection, see table

µg/kg = ppb (parts per billion)

## WEIGHTED AVERAGE RESULTS FOR THE LAST THREE SEASONS

Region	2010/2011					2009/2010					2008/2009				
	Protein (12% mb), %	FN, sec	Hlm, kg/hl	Mixo PT, min	n	Protein (12% mb), %	FN, sec	Hlm, kg/hl	Mixo PT, min	n	Protein (12% mb), %	FN, sec	Hlm, kg/hl	Mixo PT, min	n
1	12.2	393	79.7	2.9	3	11.2	299	78.7	2.9	4	11.2	435	79.7	2.7	4
2	12.3	422	77.7	2.7	12	11.4	314	77.8	2.6	30	10.3	390	76.7	3.0	24
3	11.7	410	79.7	2.7	44	10.9	351	79.5	2.5	63	10.5	395	77.7	2.7	71
4	11.2	387	81.2	2.7	25	10.6	382	80.1	2.5	23	10.3	377	79.4	2.7	14
5	11.8	355	79.6	2.7	20	11.1	379	80.0	2.5	30	11.7	304	77.6	2.4	19
6	12.5	355	80.2	2.6	11	11.8	355	79.3	2.4	24	11.8	339	78.0	2.5	34
7	12.9	401	81.7	2.4	1	12.2	375	79.5	2.7	2	-	-	-	-	-
8	-	-	-	-	-	-	-	-	-	-	13.1	397	76.4	2.9	4
9	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10	12.4	413	82.4	2.5	32	12.1	421	82.6	2.2	27	11.6	430	79.0	2.6	23
11	11.9	395	80.9	2.7	14	12.5	454	80.6	2.8	26	11.8	446	77.5	2.8	24
12	12.8	405	81.4	3.1	5	13.1	414	79.2	2.6	7	13.5	446	75.8	3.0	7
13	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
14	11.6	415	79.7	2.9	4	12.4	369	78.6	2.5	7	-	-	-	-	-
15	11.9	375	81.2	3.0	9	12.1	400	81.2	2.9	6	12.9	467	76.0	2.6	9
16	11.8	434	78.9	2.6	3	12.3	418	76.4	2.4	3	-	-	-	-	-
17	11.9	416	80.6	2.9	8	11.7	361	79.3	2.5	8	13.8	344	74.1	2.8	6
18	12.7	386	79.6	4.2	2	12.3	361	80.7	2.6	2	12.3	329	77.6	2.7	3
19	11.9	389	81.2	3.4	8	11.4	361	79.8	2.6	10	12.2	396	76.2	2.6	13
20	11.6	411	80.3	3.1	15	11.3	393	79.4	2.9	10	11.6	411	76.4	3.3	25
21	11.8	397	81.7	3.1	5	12.0	384	81.4	2.5	5	11.5	426	77.8	2.8	2
22	11.7	357	81.4	3.4	6	12.1	367	82.6	3.0	8	14.1	320	76.8	2.9	10
23	11.4	362	81.4	3.1	22	11.6	325	81.5	3.1	15	12.9	379	77.2	2.9	23
24	12.0	356	80.1	3.4	16	11.8	322	80.8	3.0	29	13.4	347	76.1	3.3	17
25	13.1	240	78.2	3.8	25	11.7	361	78.8	3.5	35	13.5	314	77.6	3.4	31
26	13.3	305	80.2	3.7	13	12.2	348	79.9	3.5	22	14.0	320	77.2	3.6	25
27	13.6	328	79.1	3.7	8	12.0	411	79.8	3.3	7	12.8	370	77.9	3.4	3
28	13.1	298	78.8	3.4	31	12.2	372	79.0	3.7	34	13.2	376	78.5	3.2	29
29	12.6	421	85.0	2.8	1	-	-	-	-	-	-	-	-	-	-
30	13.0	407	76.9	4.2	1	-	-	-	-	-	12.6	438	80.0	2.3	3
31	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
32	12.2	396	82.6	3.5	1	11.5	278	81.6	2.9	5	11.9	321	78.9	3.0	7
33	-	-	-	-	-	11.8	372	78.6	3.5	9	11.1	372	78.4	2.9	10
34	11.5	436	82.3	2.2	11	12.3	416	80.3	2.6	5	11.9	346	77.2	3.0	18
35	11.9	415	80.6	2.9	8	11.8	386	79.7	3.2	14	11.3	468	79.3	3.1	17
36	12.7	432	82.2	2.9	8	12.8	348	81.2	3.1	10	12.5	367	79.2	2.9	5
Ave.	12.1	372	80.3	3.0	372	11.7	367	79.9	2.9	480	12.0	378	77.6	2.9	480

## WEIGHTED AVERAGE QUALITY OVER 10 SEASONS



\* Includes addition of 2 kg/hl according to Hectolitre mass Dispensation.

\*\* Hectolitre mass determined using Kern 222 instrument.


## REGIONAL QUALITY

### WINTER RAINFALL AREA (Western Cape)

Production regions 1 to 6 fall within the winter rainfall area (Western Cape Province). Region 1 is Namaqualand, regions 2 - 4 are the Swartland area and regions 5 and 6 the Rûens area. The Western Cape had the highest production of all the provinces this season, namely 556 500 tons (38 %) (CEC).

The hectolitre mass averaged 79.8 kg/hl. The thousand kernel mass averaged 39.9 gram, which is equal to the previous season. The average falling number was 391 seconds. The average protein content of 11.75 % (12 % mb) was the lowest of the different production areas.

Fairly favourable climatic conditions were experienced in the Swartland. Growing conditions in the Rûens were the worst experienced over a period of four years.



The screenings of 1.82 % was lower than the previous season's 2.02 %. The Bühler extraction averaged 74.7 % (average of wheat grades B1 to B4, UT and COW) and the average colour of the flour was -2.2 KJ units. This colour indicates a very white flour that is preferred by millers and bakers.

The mixogram peak time (Quadromat mill) averaged 2.7 minutes. The average farinogram absorption was 61.9 %. The average alveogram strength was 30.3 cm<sup>2</sup> and the average strength on the extensogram was 85 cm<sup>2</sup>. The alveogram strength in the Free State was 41.2 cm<sup>2</sup> and in the irrigation areas 36.0 cm<sup>2</sup>.

The 100-gram baking test showed a very good relationship between protein content and bread volume.

## SUMMER RAINFALL AREA

(Free State)

Production regions 21 to 28, which fall within the Free State Province, had the second highest production, namely 377 400 tons (26 %) (CEC).

The 1.9 tons/ha average yield in the Free State was lower than the 2.7 tons/ha of the previous season.

The average hectolitre mass was 79.7 kg/hl. The physical characteristic thousand kernel mass (38.1 g) was lower than the previous season's 38.4 g. The average screenings was 1.51 %. The average protein content increased from 11.93 % the previous season to 12.59 % (12 % mb) this season. Although the average falling number was 314 seconds, twenty four samples gave falling numbers lower than 250 seconds and twenty one samples were below 220 seconds. These low falling numbers were caused by preharvest sprouting due to late rains.


The mixogram (Quadromat) peak time of 3.5 minutes was similar to the previous season, giving the Free State the longest average mixogram peak time of the different production areas.

The average Bühler extraction percentage in the Free State was 73.7 % (75.6 % previous season). The Kent Jones flour colour was -0.9 KJ units (-2.3 KJ units in the previous season).

The average farinogram water absorption was 64.3 %, the highest of all four areas. The average alveogram strength of 41.1 cm<sup>2</sup> and extensogram strength of 107 cm<sup>2</sup> were also the highest reported for all four areas.


The 100-gram baking test showed that the relationship between protein content and bread volume was good on average between the different grades.

## FREE STATE



## IRRIGATION AREAS

(Northern Cape, North West (plus other irrigation areas))



Production regions 7, 10 - 12, 14 - 20 and 36 falls within the irrigation areas. These areas produced 428 420 tons of wheat this season (29 % of total production) with an average yield of 5.4 tons/hectare.

Due to the significant decrease in wheat produced under dryland conditions over the past ten years, the stable production situation of wheat under irrigation has proven to be an increasingly important stabilising factor in the total production of wheat in the RSA.

The average hectolitre mass was 81.3 kg/hl and the thousand kernel mass was 40.3 g (38.6 g the previous season). The average falling number was 407 seconds. The average screenings was 1.69 % and the protein averaged 12.11 % (12 % mb).

The average mixogram (Quadromat) peak time was 2.8 minutes which was more or less the same as the previous season.

The average Bühler extraction percentage was 76.0 (73.9 % during the previous season), with an average flour colour of -2.4 KJ units.

The average farinogram water absorption was 62.8 % (61.6 % during previous season), with an average farinogram development time of 5.7 minutes.

The average alveogram strength was 36.0 cm<sup>2</sup> and the average P/L 1.01 (40.0 cm<sup>2</sup> and 1.68 respectively the previous season).

The average extensogram strength was 100 cm<sup>2</sup>. The relationship between protein content and bread volume (with the 100-gram baking test) was shown to be very good.

## OTHER SUMMER RAINFALL AND IRRIGATION AREAS (Mpumalanga, Limpopo and Gauteng)

Other summer rainfall regions, excluding the Free State, are mainly regions 29, 30, 32, 33 (Mpumalanga), 34 (Gauteng) and 35 (Limpopo). They produced in total 102 650 tons during this season (7 % of the total production).

The average hectolitre mass was 81.6 kg/hl, the highest of the four production areas. The average thousand kernel mass was 41.4 g (40.6 g the previous season), also the highest of the four areas.


The average falling number was 425 seconds, with the average percentage screenings 1.89 %. The average protein content was 11.78 % (12 % mb), which is similar to the previous year.

The average mixogram (Quadromat) peak time was 2.6 minutes (3.1 minutes the previous season).

The average Bühler extraction was 76.0 %, with an average colour of -2.0 KJ units (76.2 % and -2.1 KJ units the previous season). The farinogram average water absorption was 63.3 % (60.1 % the previous season) and had an average development time of 5.0 minutes.

The average alveogram strength was 31.9 cm<sup>2</sup> and the average extensogram strength 78 cm<sup>2</sup>.

The 100-gram baking test showed a good relationship between protein content and bread volume.



Regional maps with gratitude to ARC - Small Grain Institute

# REGIONAL QUALITY WEIGHTED AVERAGES

	<i>Winter rainfall area (Western Cape)</i>			<i>Summer rainfall area (Free State)</i>			<i>Irrigation areas</i>			<i>Other Summer rainfall and Irrigation areas</i>			<i>RSA average</i>
<i>Individual samples n</i>	115			126			109			22			372
<b>Regions</b>	1 - 6			21 - 28			7, 10 - 12, 14 - 20, 36			29 - 35			All
<b>Hectolitre mass dirty, kg/hl</b>	79.8			79.7			81.3			81.6			80.3
<b>1000 kernel mass (13% mb), g</b>	39.9			38.1			40.3			41.4			39.5
<b>Falling number, sec</b>	391			314			407			425			372
<b>Screenings (1,8 mm), %</b>	1.82			1.51			1.69			1.89			1.68
<b>Protein (12% mb), % (ww)</b>	11.75			12.59			12.11			11.78			12.14
<b>Mixogram peak time, min (Quadromat)</b>	2.7			3.5			2.8			2.6			3.0
<i>Individual samples per class and grade, n</i>	38	35	21	64	21	13	52	32	10	10	5	4	164 93 48
	7	13	1	7	8	13	7	7	1	2	0	1	23 28 16
<i>Composite samples per class and grade, n = 100</i>	B1	B2	B3	B1	B2	B3	B1	B2	B3	B1	B2	B3	B1 B2 B3
	B4	UT	COW	B4	UT	COW	B4	UT	COW	B4	UT	COW	B4 UT COW
<i>Composite samples, n</i>	6	5	3	8	7	6	11	10	6	4	2	2	29 24 17
	3	3	1	4	4	5	3	3	1	1	0	1	11 10 8
<b>Bühler extraction, %</b>	74.5	74.6	75.1	74.1	74.1	73.7	75.8	76.0	75.8	75.9	75.6	76.5	75.1 75.1 75.0
	75.1	74.9	72.7	73.2	72.8	73.3	76.2	76.2	77.0	75.3	-	77.0	74.7 74.5 74.2
<b>Flour colour, KJ</b>	-2.2	-2.3	-2.4	-1.5	-1.7	-1.3	-2.2	-2.4	-2.5	-2.1	-2.1	-2.5	-2.0 -2.2 -2.1
	-2.4	-2.1	-0.7	-0.9	-0.2	1.1	-2.6	-2.6	-1.8	-2.5	-	0.3	-1.9 -1.5 0.4
<b>Flour protein (12% mb), %</b>	11.8	10.4	9.4	12.3	10.7	10.3	12.1	10.5	9.5	11.7	10.2	9.5	12.0 10.5 9.8
	9.0	10.6	11.9	9.9	12.7	12.5	10.7	11.1	10.2	9.5	-	11.9	9.8 11.6 12.1
<b>Wet gluten (14% mb), %</b>	32.6	27.9	24.3	32.8	28.4	26.6	32.8	28.7	25.3	32.1	28.3	26.2	32.7 28.4 25.7
	25.9	28.4	33.4	27.6	33.7	32.5	30.4	31.2	28.0	26.5	-	25.1	27.8 31.4 31.1
<b>Dry gluten (14% mb), %</b>	11.9	10.0	8.7	11.8	9.9	9.5	11.5	9.9	8.8	11.2	9.6	8.7	11.6 9.9 9.0
	9.2	10.2	11.8	9.4	12.0	11.7	10.2	10.7	9.5	8.9	-	8.7	9.5 11.1 11.1
<b>Farinogram:</b>	63.2	62.3	60.7	65.1	63.5	63.1	64.0	62.4	60.9	64.5	63.1	61.5	64.2 62.8 61.7
<b>Water absorption (14% mb), %</b>	59.6	61.9	63.1	63.3	66.0	64.9	63.6	62.8	62.0	63.0	-	62.4	62.3 63.8 64.0
<b>Farinogram:</b>	5.8	5.1	4.6	7.6	6.0	5.6	7.3	5.1	4.8	6.6	4.0	4.4	7.0 5.3 5.0
<b>Development time, min</b>	3.1	5.2	4.7	4.0	5.0	5.3	4.8	5.5	4.2	4.0	-	2.5	4.0 5.2 4.7
<b>Alveogram:</b>	34.7	30.8	27.1	46.5	35.6	35.8	45.7	32.6	29.2	41.4	27.0	24.0	43.1 32.6 30.5
<b>Strength (S), cm<sup>2</sup></b>	24.0	29.5	32.1	33.4	51.1	45.6	31.6	34.0	24.6	20.9	-	30.6	29.2 39.5 39.4
<b>Alveogram:</b>	0.86	1.15	1.10	1.44	1.53	2.02	0.96	1.06	1.20	1.03	0.97	0.98	1.08 1.21 1.45
P/L	1.07	1.16	0.85	2.39	1.76	1.35	0.96	0.75	0.73	1.42	-	4.21	1.55 1.28 1.57
<b>Extensogram:</b>	93	86	75	114	101	100	121	94	85	104	57	60	111 91 86
<b>Strength, cm<sup>2</sup></b>	67	89	102	97	112	117	84	98	74	56	-	-	82 101 95
<b>Mixogram peak time, min</b>	2.5	2.5	2.3	3.1	2.8	3.1	2.8	2.7	2.8	2.8	2.3	2.3	2.8 2.7 2.8
	2.4	2.7	2.3	3.3	3.3	3.5	2.3	2.3	2.3	2.0	-	3.5	2.7 2.8 3.2
<b>Relationship between protein and bread volume</b>	VG	EX	EX	Q	G	Q	VG	EX	VG	VG	EX	EX	G VG VG
	EX = Excellent	VG = Very Good	G = Good	Q = Questionable	VG = Very Good	EX = Excellent	VG = Very Good	EX = Excellent	VG = Very Good	VG = Very Good	VG = Very Good	VG = Very Good	VG = Good G

Q = Questionable

EP = Extremely Poor

## RSA WHEAT PRODUCTION AREAS



**WHEAT SEED SOLD BY COMMERCIAL GRAIN SILO OWNERS TO  
WHEAT PRODUCERS FOR THE 2010 PLANTING SEASON**

<u>Cultivar</u>	<u>%</u>	<u>Cultivar</u>	<u>%</u>
SST 027	16.20	SST 347	0.25
SST 835	13.14	PAN 3355	0.23
SST 88	12.77	PAN 3349	0.22
SST 843	9.30	Buffels	0.214
SST 015	8.72	SST 334	0.205
Duzi	8.43	SST 966	0.146
Elands	6.17	PAN 3377	0.106
SST 047	5.28	Gariep	0.105
SST 356	4.72	Olifants	0.103
CRN 826	4.17	SST 399	0.090
SST 822	2.15	Kariega	0.084
Krokodil	1.15	PAN 3368	0.082
SST 877	0.94	Baviaans	0.067
SST 806	0.73	Komati	0.038
SST 056	0.68	PAN 3144	0.030
Matlabas	0.66	Betta DN	0.027
SST 876	0.64	SST 322	0.012
SST 363	0.62	SST 825	0.007
PAN 3120	0.43	PAN 3364	0.006
Tugela	0.39	Inia	0.002
PAN 3118	0.34	SST 935	0.0019
PAN 3434	0.33	SST 367	<u>0.0004</u>
			100

Note: These figures are not absolute, but the best and only figures available.

## South African Winter Cereal Production

Wheat is by far the biggest winter cereal crop planted in South Africa. Other winter crops are barley for malting purposes and canola. Summer field crops are better suited for the South African climatic conditions. Maize has the largest crop size of the different crops, followed by wheat, then sunflower seed, soya-beans, sorghum, barley, groundnuts, dry beans and canola.

South Africa (made up of nine provinces) is divided into 36 crop production regions with wheat planted in about 32 of these regions. These production regions are described on pages 23 to 54 (in the header of the left page) giving the specific intake silo names for each region.


The three main wheat producing provinces are Western Cape (winter rainfall), Free State (summer rainfall) and the Northern Cape (irrigation). A fourth province worth mentioning is the North-West (mainly irrigation). The local production is not sufficient for domestic requirements and South Africa has to import wheat to meet its domestic consumption.

The Western Cape province produced 556 500 tons and the Free State province followed with 377 400 tons. (Seventh estimate by the Crop Estimates Committee, CEC). These two provinces were responsible for 64 % of the total wheat produced. The Northern Cape produced 254 600 tons and North West 123 200 tons.

The yield in the main production areas ranged from 6.7 tons per hectare in the Northern Cape (irrigation area), 1.9 tons per hectare in the Free State and 2.1 tons per hectare for the Western Cape. Gauteng gave a yield of 6.4 tons per hectare, followed by North West with 5.6 tons per hectare, Limpopo 5.5 tons per hectare and Mpumalanga with 5.4 tons per hectare. KwaZulu-Natal and the Eastern Cape yielded 5.2 and 4.2 tons per hectare respectively. See graph on page 13.

South Africa has three major wheat-breeding programs. The South African breeders can only release a new cultivar or an introduction cultivar if it has better agronomical as well as better flour quality characteristics than the cultivars planted commercially in a specific area. Producers continuously strive to improve the wheat yield and quality by selecting the best cultivars for commercial production in a specific area. Grading standards are also set high to ensure adequate quality control.

**Average yield per province  
(Irrigation versus summer and winter rainfall areas)**



(Based on figures obtained from the CEC)

## Comparison of Flour Quality over the last four seasons



Flour Quality 2010/2011 season			
Flour protein (12% mb) (%)	11.0	Farinogram abs. (14% mb) (%)	63.2
Bread volume 100g (cm <sup>3</sup> )	832	Farinogram dev. time (min.)	5.5
Mixogram (Bühler) peak time (min)	2.8	Alveogram strength (cm <sup>2</sup> )	36.2
Wet gluten (14% mb) (%)	29.7	Alveogram P/L	1.29
Dry gluten (14% mb) (%)	10.4	Extensogram strength (cm <sup>2</sup> )	97

Flour Quality 2009/2010 season			
Flour protein (12% mb) (%)	10.5	Farinogram abs. (14% mb) (%)	61.0
Bread volume 100g (cm <sup>3</sup> )	843	Farinogram dev. time (min.)	3.5
Mixogram (Bühler) peak time (min)	2.6	Alveogram strength (cm <sup>2</sup> )	35.5
Wet gluten (14% mb) (%)	28.6	Alveogram P/L	1.17
Dry gluten (14% mb) (%)	10.0	Extensogram strength (cm <sup>2</sup> )	83



Flour Quality 2008/2009 season			
Flour protein (12% mb) (%)	11.1	Farinogram abs. (14% mb) (%)	61.1
Bread volume 100g (cm <sup>3</sup> )	902	Farinogram dev. time (min.)	4.0
Mixogram (Bühler) peak time (min)	2.6	Alveogram strength (cm <sup>2</sup> )	38.0
Wet gluten (14% mb) (%)	-	Alveogram P/L	0.90
Dry gluten (14% mb) (%)	-	Extensogram strength (cm <sup>2</sup> )	90

Flour Quality 2007/2008 season			
Flour protein (12% mb) (%)	10.8	Farinogram abs. (14% mb) (%)	60.8
Bread volume 100g (cm <sup>3</sup> )	827	Farinogram dev. time (min.)	3.6
Mixogram (Bühler) peak time (min)	2.8	Alveogram strength (cm <sup>2</sup> )	41.9
Wet gluten (14% mb) (%)	-	Alveogram P/L	0.94
Dry gluten (14% mb) (%)	-	Extensogram strength (cm <sup>2</sup> )	97



## Comparison of protein content vs loaf volume



## Comparison of wet gluten content vs loaf volume



## Comparison of protein content vs dry gluten content





**South African Grain Information Service**  
**Suid-Afrikaanse Graan Inligtingsdiens**

Association incorporated under section 21 / Vereniging ingelyf kragtens artikel 21 (Reg No. 1997/019186/08)

## WHEAT IMPORTS / EXPORTS

2009/10 Season

Progressive: 2009/10/03 - 2010/10/01

Updated: 2010/11/26

Imported from:	Total Tons	Destined for:		
		RSA	Africa	Overseas
Overseas	1387813	1277972	109841	0
RSA	0	0	0	0
Africa	0	0	0	0
	1387813	1277972	109841	0
Less: RSA for RSA		0	0	n/a
	<b>1387813</b>	<b>1277972</b>	<b>109841</b>	<b>0</b>

Exported from:	Total Tons	Destined for:		
		RSA	Africa	Overseas
Overseas	116392	0	116392	0
RSA	204127	0	204127	0
Africa	0	0	0	0
	320519	0	320519	0
Less: RSA for RSA	0	0	n/a	n/a
	<b>320519</b>	<b>0</b>	<b>320519</b>	<b>0</b>

Information updated due to:

- i) Revised information received from importers/exporters.
- ii) Grain originally allocated for RSA but finally exported to Africa or vice versa.



**South African Grain Information Service**  
**Suid-Afrikaanse Graan Inligtingsdiens**

Association incorporated under section 21 / Vereniging ingelyf kragtens artikel 21 (Reg No. 1997/019186/08)

---

**WHEAT IMPORTS / EXPORTS**

**2010/11 Season**

Progressive: 2010/10/02 - 2011/05/13

Published: 2011/05/17

Imported from:	Total Tons	Destined for:		
		RSA	Africa	Overseas
Overseas	1005172	964340	34429	6403
RSA	0	0	0	0
Africa	0	0	0	0
	1005172	964340	34429	6403
Less: RSA for RSA		0	0	n/a
	<b>1005172</b>	<b>964340</b>	<b>34429</b>	<b>6403</b>

Exported from:	Total Tons	Destined for:		
			Africa	Overseas
Overseas	36899	0	36899	0
RSA	100393	0	100393	0
Africa	0	0	0	0
	137292	0	137292	0
Less: RSA for RSA	0	0	n/a	n/a
	<b>137292</b>	<b>0</b>	<b>137292</b>	<b>0</b>

**2009/10 Season**  
**Progressive: 2009/10/03 - 2010/10/01**  
**Updated: 2010/11/26**

### **WHEAT EXPORTS**

COUNTRY	AFRICA TONS	RSA TONS	OVERSEAS TONS
Botswana	95164		
Mozambique	1991		
Lesotho	61699		
Namibia	21630		
Swaziland	38538		
Zimbabwe	101497		
	<b>320519</b>		

### **WHEAT IMPORTS**

FROM COUNTRY	FOR AFRICA TONS	FOR RSA TONS	FOR OVERSEAS TONS
Germany	80487	809934	
Ukraine		41230	
USA	2454	173030	
Australia	3080	55312	
Canada		72911	
Brazil	12691	123944	
Lithuania	11129	1611	
	<b>109841</b>	<b>1277972</b>	

Information updated due to:

- i) Revised information received from importers/exporters.
- ii) Grain originally allocated for RSA but finally exported to Africa or vice versa.

**2010/11 Season**

Progressive: 2010/10/02 - 2011/05/13

Published: 2011/05/17

**WHEAT EXPORTS**

COUNTRY	AFRICA TONS	COUNTRY	OVERSEAS TONS
Botswana	52723	Madagascar	6403
Lesotho	50004		
Mozambique	6		
Namibia	8147		
Swaziland	13944		
Zambia	4124		
Zimbabwe	1941		
	130889		6403

**WHEAT IMPORTS**

FROM COUNTRY	FOR AFRICA TONS	FOR RSA TONS	FOR OVERSEAS TONS
Argentina	12430	510115	0
Australia	5116	53198	0
Brazil	5808	28600	0
Canada	0	47415	0
Germany	0	62186	0
Uruguay	0	25249	0
USA	11075	237577	6403
	34429	964340	6403

**WHEAT**      '000 t

**Updated:**

		Oct - Sep	1997/98	1998/99	1999/00	2000/01	2001/02	2002/03	2003/04	2004/05	2005/06	2006/07	2007/08	2008/09	2009/10	Projection	Actual
																Months:	Oct-Apr
																12	7
Opening stock		578.0	1241.0	771.0	507.0	551.0	580.0	897.0	598.0	574.0	582.0	376.0	509.0	694.0	579.0	579.0	
Prod deliveries (a)		2449.0	1644.0	1725.0	2353.0	2415.0	2387.0	1512.0	1670.0	1893.0	2045.0	1876.0	2130.0	1910.0		1371.0	
Crop estimate (b)		2283.5	1531.0	1725.0	2349.0	2492.9	2320.7	1540.0	1680.0	1905.0	2105.0	1905.0	2130.0	1958.0		1430.0	
Imports		469.0	484.0	624.0	308.0	407.0	747.0	1042.0	1227.0	1055.0	777.0	1396.0	1192.0	1285.0	936.0	936.0	
Surplus		0.0	0.0	0.0	0.0	0.0	0.0	6.0	6.0	9.0	32.0	0.0	13.0	0.0	9.0	9.0	
<b>Available</b>		<b>3496.0</b>	<b>3369.0</b>	<b>3120.0</b>	<b>3168.0</b>	<b>3373.0</b>	<b>3714.0</b>	<b>3457.0</b>	<b>3501.0</b>	<b>3531.0</b>	<b>3436.0</b>	<b>3648.0</b>	<b>3844.0</b>	<b>3889.0</b>	<b>2954.0</b>	<b>2895.0</b>	
Processed		2181.0	2400.0	2371.0	2427.0	2541.0	2577.0	2653.0	2736.0	2793.0	2820.0	2845.0	2857.0	3017.0	2849.1	1662.0	
-human		2138.0	2348.0	2345.0	2424.0	2519.0	2575.0	2652.0	2734.0	2781.0	2818.0	2844.0	2849.0	2991.0	2847.4	1661.0	
-animal		43.0	52.0	24.0	2.0	22.0	2.0	1.0	2.0	12.0	2.0	1.0	8.0	26.0	1.7	1.0	
-gristing		0.0	0.0	2.0	1.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
-bio-fuel		0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Producers		0.0	0.0	43.0	33.0	31.0	24.0	13.0	7.0	10.0	7.0	12.0	12.0	14.0	5.0	5.0	
End consumers		2.0	5.0	12.0	4.0	7.0	5.0	2.0	2.0	4.0	4.0	2.0	5.0	3.0	2.0	2.0	
Seed		0.0	16.0	26.0	24.0	27.0	20.0	21.0	16.0	26.0	17.0	22.0	26.0	17.0	10.0	10.0	
Net receipts(-)/dispatch(+)		-7.0	76.0	37.0	9.0	15.0	11.0	12.0	6.0	5.0	1.0	26.0	19.0	15.0	-1.0	-1.0	
Deficit		0.0	60.0	52.0	17.0	23.0	1.0	0.0	0.0	0.0	0.0	9.0	0.0	4.0	0.0	0.0	
Exports		79.0	75.0	72.0	103.0	149.0	179.0	158.0	158.0	111.0	211.0	223.0	231.0	240.0	108.0	108.0	
Utilised:		<b>2255.0</b>	<b>2632.0</b>	<b>2613.0</b>	<b>2617.0</b>	<b>2793.0</b>	<b>2817.0</b>	<b>2859.0</b>	<b>2925.0</b>	<b>2949.0</b>	<b>3060.0</b>	<b>3139.0</b>	<b>3150.0</b>	<b>3310.0</b>	<b>2973.1</b>	<b>1786.0</b>	
<b>Stock</b>		<b>1241.0</b>	<b>737.0</b>	<b>507.0</b>	<b>551.0</b>	<b>580.0</b>	<b>897.0</b>	<b>598.0</b>	<b>576.0</b>	<b>582.0</b>	<b>376.0</b>	<b>509.0</b>	<b>694.0</b>	<b>579.0</b>	<b>-19.1</b>	<b>1109.0</b>	
- processed p/month		181.8	200.0	197.6	202.3	211.8	214.8	221.1	228.0	232.8	235.0	237.1	238.1	251.4	237.4	237.4	
- months' stock		6.8	3.7	2.6	2.7	2.7	4.2	2.7	2.5	2.5	1.6	2.1	2.9	2.3	-0.1	4.7	

**SOUTH AFRICAN**  
**WINTER RAINFALL WHEAT**  
**Western Cape Province**

PRODUCTION REGION	(1) Namaqualand				(2) Swartland Western Region							
	Intake silos	Bitterfontein	Graafwater	Landplaas	Vanrhynsdorp	Vredendal	Bergvvier	Darling	Koperfontein	Vredenburg		
<b>WHEAT</b>												
Protein (12% mb), %	ave 12.2	min 11.3	max 13.3	stdev 1.01	ave 12.3	min 11.6	max 13.0	stdev 0.44				
Falling number, sec	393	377	407	15.10	422	395	458	19.96				
1000 Kernel mass (13% mb), g	43.2	40.2	48.9	4.91	37.9	34.4	38.9	1.25				
Hectolitre mass (dirty), kg hl	79.7	79.4	80.0	0.31	77.7	74.5	81.2	1.95				
Screenings (<1.8mm), %	2.40	2.20	2.54	0.18	2.67	1.78	4.72	0.81				
Total damaged kernels, %	1.15	1.08	1.28	0.12	0.63	0.18	0.84	0.21				
<i>Number of samples</i>	3				12							
<b>CULTIVARS</b>												
cultivars with highest % occurrence	SST 88 36.0				SST 027 36.9							
	SST 015 31.7				SST 015 28.8							
	SST 027 26.7				SST 88 24.7							
	SST 047 3.0				SST 047 3.5							
	PAN 3492 2.7				SST 056 2.3							
<i>Number of samples</i>	3				12							
<b>MIXOGRAM (Quadromat)</b>												
Peak time, min	ave 2.9	min 2.6	max 3.3	stdev 0.38	ave 2.7	min 2.5	max 2.8	stdev 0.11				
Tail height (6min), mm	50	49	51	1.00	48	45	50	1.73				
<i>Number of samples</i>	3				12							
<b>BÜHLER EXTRACTION, %</b>	B1 74.4	B2 74.0	B3 -	B4 -	UT -	COW -	B1 74.5	B2 -	B3 -			
										73.7	-	
<b>FLOUR</b>												
Protein (12% mb), %	11.4	9.7	-	-	-	-	11.7	-	-	-	11.2	-
Colour, KJ	-2.3	-2.2	-	-	-	-	-2.4	-	-	-	-2.0	-
<b>GLUTEN</b>												
Wet gluten (14% mb), %	31.5	24.5	-	-	-	-	31.8	-	-	-	30.5	-
Dry gluten (14% mb), %	11.5	8.3	-	-	-	-	11.9	-	-	-	11.2	-
<b>FARINOGRAM</b>												
Water absorption (14% mb), %	63.6	63.0	-	-	-	-	62.6	-	-	-	61.1	-
Development time, min	6.2	2.7	-	-	-	-	5.7	-	-	-	5.3	-
Stability, min	7.3	9.2	-	-	-	-	7.7	-	-	-	6.8	-
Mixing tolerance index, BU	45	13	-	-	-	-	34	-	-	-	38	-
<b>EXTENSOGRAM (45 min pull)</b>												
Area, cm <sup>2</sup>	97	88	-	-	-	-	87	-	-	-	86	-
Maximum height, BU	335	400	-	-	-	-	330	-	-	-	325	-
Extensibility, mm	202	156	-	-	-	-	186	-	-	-	184	-
<b>ALVEOGRAM</b>												
Strength (S), cm <sup>2</sup>	37.0	34.1	-	-	-	-	32.3	-	-	-	29.4	-
Stability (P), mm	91	107	-	-	-	-	82	-	-	-	81	-
Distensibility (L), mm	93	63	-	-	-	-	92	-	-	-	84	-
Configuration ratio (P/L)	0.98	1.70	-	-	-	-	0.89	-	-	-	0.96	-
<b>MIXOGRAM</b>												
Peak time, min	2.7	2.7	-	-	-	-	2.5	-	-	-	2.8	-
<b>100g BAKING TEST</b>												
Loaf volume, cm <sup>3</sup>	870	775	-	-	-	-	880	-	-	-	890	-
Evaluation	1	0	-	-	-	-	1	-	-	-	0	-

# RHEOLOGICAL GRAPHS PER PRODUCTION REGION

## MIXOGRAM

1

2

## FARINOGRAM

1

2

## EXTENSOGRAM

1

2

## ALVEOGRAM

1

2

**SOUTH AFRICAN  
WINTER RAINFALL WHEAT  
Western Cape Province**

PRODUCTION REGION	(3) Swartland Central Region				(4) Swartland Eastern Region									
	Eendekuil	Klipheuwel	Koringberg	Malmesbury	Moorreesburg	Moravia	Piketberg	Pools	Ruststasie	Ceres	Gouda	Halfmanshof	Leliedam	Porterville
Intake silos														
<b>WHEAT</b>														
Protein (12% mb), %	<b>ave</b> 11.7	<b>min</b> 9.7	<b>max</b> 13.5	<b>stdev</b> 1.00	<b>ave</b> 11.2	<b>min</b> 9.3	<b>max</b> 13.2	<b>stdev</b> 0.93						
Falling number, sec	410	350	475	28.72	387	331	451	30.13						
1000 Kernel mass (13% mb), g	40.6	34.0	45.6	2.41	39.9	35.8	44.0	2.10						
Hectolitre mass (dirty), kg/hl	79.7	76.2	83.6	1.47	81.2	77.5	84.1	1.31						
Screenings (<1.8mm), %	2.02	0.30	6.64	0.99	1.09	0.04	5.45	1.37						
Total damaged kernels, %	0.32	0.00	0.72	0.17	0.43	0.08	1.36	0.32						
<b>Number of samples</b>	<b>44</b>				<b>25</b>									
<b>CULTIVARS</b>														
cultivars with highest % occurrence	SST 015				SST 88				29.3					
	SST 027				SST 027				26.2					
	SST 88				SST 015				22.9					
	SST 835				SST 57				11.8					
	SST 047				SST 047				7.5					
<b>Number of samples</b>	<b>44</b>				<b>25</b>									
<b>MIXOGRAM (Quadromat)</b>														
Peak time, min	<b>ave</b> 2.7	<b>min</b> 2.2	<b>max</b> 3.8	<b>stdev</b> 0.27	<b>ave</b> 2.7	<b>min</b> 2.1	<b>max</b> 3.3	<b>stdev</b> 0.25						
Tail height (6min), mm	48	42	55	3.06	47	42	51	2.78						
<b>Number of samples</b>	<b>44</b>				<b>25</b>									
<b>BÜHLER EXTRACTION, %</b>														
<b>FLOUR</b>	<b>B1</b> 74.5	<b>B2</b> 74.7	<b>B3</b> 75.1	<b>B4</b> 75.7	<b>UT</b> 74.9	<b>COW</b> -	<b>B1</b> 74.3	<b>B2</b> 74.4	<b>B3</b> 74.6	<b>B4</b> 73.9	<b>UT</b> 76.1	<b>COW</b> -		
Protein (12% mb), %	12.0	10.5	9.3	9.5	11.5	-	11.8	10.7	9.3	9.3	9.0	-		
Colour, KJ	-2.4	-2.5	-2.6	-2.4	-2.3	-	-2.6	-2.7	-2.6	-2.6	-2.1	-		
<b>GLUTEN</b>														
Wet gluten (14% mb), %	33.0	28.4	24.4	25.4	31.5	-	33.6	29.2	24.2	26.3	23.3	-		
Dry gluten (14% mb), %	12.1	10.6	8.9	9.0	11.5	-	12.2	10.3	8.5	9.3	7.8	-		
<b>FARINOGRAM</b>														
Water absorption (14% mb), %	63.8	62.1	60.8	61.7	62.7	-	62.8	61.6	60.4	60.7	61.9	-		
Development time, min	6.2	5.4	4.8	3.3	5.7	-	5.9	6.0	4.8	3.9	4.5	-		
Stability, min	7.4	7.0	6.1	6.2	6.8	-	7.4	7.4	6.1	5.4	5.5	-		
Mixing tolerance index, BU	35	43	50	35	39	-	35	38	43	45	47	-		
<b>EXTENSOGRAM (45 min pull)</b>														
Area, cm <sup>2</sup>	96	94	79	62	92	-	95	85	73	73	-	-		
Maximum height, BU	315	350	310	290	330	-	320	325	320	310	-	-		
Extensibility, mm	210	189	175	147	191	-	209	183	161	157	-	-		
<b>ALVEOGRAM</b>														
Strength (S), cm <sup>2</sup>	37.3	32.3	27.7	26.3	32.6	-	32.3	29.4	26.8	25.7	26.6	-		
Stability (P), mm	85	86	83	87	84	-	77	79	77	80	94	-		
Distensibility (L), mm	107	86	76	66	90	-	100	82	80	73	59	-		
Configuration ratio (P/L)	0.79	1.00	1.09	1.32	0.93	-	0.77	0.96	0.96	1.10	1.59	-		
<b>MIXOGRAM</b>														
Peak time, min	2.5	2.4	2.3	2.5	2.5	-	2.5	2.4	2.2	2.2	2.8	-		
<b>100g BAKING TEST</b>														
Loaf volume, cm <sup>3</sup>	885	830	760	725	900	-	880	790	820	775	700	-		
Evaluation	1	0	0	2	0	-	1	2	0	0	2	-		

# RHEOLOGICAL GRAPHS PER PRODUCTION REGION

## MIXOGRAM

3

4

## FARINOGRAM

3

4

## EXTENSOGRAM

3

4

## ALVEOGRAM

3

4

**SOUTH AFRICAN  
WINTER RAINFALL WHEAT  
Western Cape Province**

PRODUCTION REGION	(5) Rüens Western Region				(6) Rüens Eastern Region							
	Bredasdorp	Caledon	Klipdale	Krike	Albertinia	Ashton	Camfer	Heidelberg				
Intake silos	Napier	Protem	Rietpoel	Villiersdorp	Karringmelksrivier	Kleinberg	Protom	Riversdal				
								Swellendam				
<b>WHEAT</b>	<b>ave</b>	<b>min</b>	<b>max</b>	<b>stdev</b>	<b>ave</b>	<b>min</b>	<b>max</b>	<b>stdev</b>				
Protein (12% mb), %	11.8	9.8	13.1	0.88	12.5	10.9	15.1	1.08				
Falling number, sec	355	81	417	73.63	355	321	407	26.00				
1000 Kernel mass (13% mb), g	39.8	36.6	43.5	2.14	38.7	35.7	42.4	2.30				
Hectolitre mass (dirty), kg/hl	79.6	76.2	80.8	1.41	80.2	78.2	82.0	1.05				
Screenings (<1.8mm), %	1.79	0.60	2.54	0.49	1.64	0.48	2.80	0.79				
Total damaged kernels, %	0.68	0.00	2.02	0.59	0.53	0.12	1.28	0.39				
<b>Number of samples</b>	<b>20</b>				<b>11</b>							
<b>CULTIVARS</b>	SST 88      56.7				SST 88      41.7							
cultivars	SST 027      25.1				SST 027      34.0							
with highest % occurrence	SST 015      15.5				SST 015      21.5							
	SST 047      1.8				SST 57      1.1							
	PAN 3492      0.6				PAN 3492      0.9							
<b>Number of samples</b>	<b>20</b>				<b>11</b>							
<b>MIXOGRAM (Quadromat)</b>	<b>ave</b>	<b>min</b>	<b>max</b>	<b>stdev</b>	<b>ave</b>	<b>min</b>	<b>max</b>	<b>stdev</b>				
Peak time, min	2.7	2.3	3.0	0.18	2.6	2.3	3.3	0.29				
Tail height (6min), mm	48	41	50	2.20	48	46	54	2.16				
<b>Number of samples</b>	<b>20</b>				<b>11</b>							
<b>BÜHLER EXTRACTION, %</b>	<b>B1</b>	<b>B2</b>	<b>B3</b>	<b>B4</b>	<b>UT</b>	<b>COW</b>	<b>B1</b>	<b>B2</b>	<b>B3</b>	<b>B4</b>	<b>UT</b>	<b>COW</b>
	74.2	74.8	75.7	75.8	-	72.7	75.3	75.1	-	-	-	-
<b>FLOUR</b>												
Protein (12% mb), %	11.9	10.6	9.5	8.2	-	11.9	11.9	10.6	-	-	-	-
Colour, KJ	-2.2	-2.1	-1.9	-2.3	-	-0.7	-1.5	-1.8	-	-	-	-
<b>GLUTEN</b>												
Wet gluten (14% mb), %	32.6	27.8	24.4	-	-	33.4	32.8	29.6	-	-	-	-
Dry gluten (14% mb), %	11.8	10.0	8.8	-	-	11.8	11.8	10.9	-	-	-	-
<b>FARINOGRAM</b>												
Water absorption (14% mb), %	62.4	61.5	60.8	56.4	-	63.1	63.9	63.1	-	-	-	-
Development time, min	6.2	6.0	4.3	2.2	-	4.7	4.8	5.4	-	-	-	-
Stability, min	7.4	7.0	6.7	5.5	-	5.2	6.3	6.3	-	-	-	-
Mixing tolerance index, BU	39	41	38	40	-	64	38	38	-	-	-	-
<b>EXTENSOGRAM (45 min pull)</b>												
Area, cm <sup>2</sup>	93	78	74	67	-	102	87	86	-	-	-	-
Maximum height, BU	325	295	320	315	-	360	325	295	-	-	-	-
Extensibility, mm	201	172	160	137	-	201	195	190	-	-	-	-
<b>ALVEOGRAM</b>												
Strength (S), cm <sup>2</sup>	34.4	29.4	26.9	20.0	-	32.1	34.6	28.9	-	-	-	-
Stability (P), mm	81	84	86	63	-	79	89	85	-	-	-	-
Distensibility (L), mm	105	80	69	79	-	93	94	81	-	-	-	-
Configuration ratio (P/L)	0.77	1.05	1.25	0.80	-	0.85	0.95	1.05	-	-	-	-
<b>MIXOGRAM</b>												
Peak time, min	2.5	2.4	2.5	2.5	-	2.3	2.3	2.4	-	-	-	-
<b>100g BAKING TEST</b>												
Loaf volume, cm <sup>3</sup>	860	835	800	760	-	880	875	845	-	-	-	-
Evaluation	2	0	0	0	-	1	1	0	-	-	-	-

# RHEOLOGICAL GRAPHS PER PRODUCTION REGION

## MIXOGRAM

5

6

## FARINOGRAM

5

5

## EXTENSOGRAM

5

6

## ALVEOGRAM

5

6

**SOUTH AFRICAN  
IRRIGATION WHEAT  
Eastern Cape and Vaal and Orange river area**

PRODUCTION REGION	(7) Eastern Cape Southern Region					(10) Griqualand-West						
	Avontuur Humansdorp Paterson Uitenhage					Britstown Douglas Havenga Brug Marydale Modderrivier Oranjerivierstasie Prieska Rietrivier Upington						
<b>WHEAT</b>												
Protein (12% mb), %	ave 12.9	min -	max -	stdev -		ave 12.4	min 10.5	max 14.6	stdev 0.92			
Falling number, sec	401	-	-	-		413	372	514	24.92			
1000 Kernel mass (13% mb), g	39.4	-	-	-		42.1	37.5	48.6	3.09			
Hectolitre mass (dirty), kg/hl	81.7	-	-	-		82.4	78.4	85.6	1.95			
Screenings (<1.8mm), %	1.46	-	-	-		1.30	0.28	4.12	0.99			
Total damaged kernels, %	0.34	-	-	-		0.14	0.00	0.40	0.09			
<b>Number of samples</b>	<b>1</b>				<b>32</b>							
<b>CULTIVARS</b>												
cultivars	SST 835				100.0				SST 835			
with highest % occurrence									Duzi 40.0			
									SST 843 24.9			
									Baviaans 13.4			
									PAN 3434 7.5			
<b>Number of samples</b>	<b>1</b>				<b>32</b>							
<b>MIXOGRAM (Quadromat)</b>												
Peak time, min	ave 2.4	min -	max -	stdev -		ave 2.5	min 1.8	max 4.6	stdev 0.53			
Tail height (6min), mm	51	-	-	-		49	43	54	2.71			
<b>Number of samples</b>	<b>1</b>				<b>32</b>							
<b>BÜHLER EXTRACTION, %</b>	<b>B1</b> 76.1	<b>B2</b> -	<b>B3</b> -	<b>B4</b> -	<b>UT</b> -	<b>COW</b> -	<b>B1</b> 76.1	<b>B2</b> 75.6	<b>B3</b> 76.2	<b>B4</b> 76.6	<b>UT</b> 78.3	<b>COW</b> -
<b>FLOUR</b>												
Protein (12% mb), %	11.7	-	-	-	-	-	11.9	10.7	9.4	11.3	11.5	-
Colour, KJ	-1.8	-	-	-	-	-	-2.4	-2.3	-2.6	-2.8	-2.7	-
<b>GLUTEN</b>												
Wet gluten (14% mb), %	32.9	-	-	-	-	-	34.2	30.4	25.6	33.2	34.6	-
Dry gluten (14% mb), %	11.5	-	-	-	-	-	12.1	10.6	8.9	11.0	11.9	-
<b>FARINOGRAM</b>												
Water absorption (14% mb), %	64.9	-	-	-	-	-	64.8	64.3	61.1	64.4	64.6	-
Development time, min	4.8	-	-	-	-	-	5.8	4.7	4.2	5.3	5.3	-
Stability, min	5.7	-	-	-	-	-	6.5	6.5	6.3	4.7	5.0	-
Mixing tolerance index, BU	60	-	-	-	-	-	41	43	45	55	53	-
<b>EXTENSOGRAM (45 min pull)</b>												
Area, cm <sup>2</sup>	-	-	-	-	-	-	101	87	75	92	95	-
Maximum height, BU	-	-	-	-	-	-	320	345	325	330	350	-
Extensibility, mm	-	-	-	-	-	-	215	184	163	198	192	-
<b>ALVEOGRAM</b>												
Strength (S), cm <sup>2</sup>	39.0	-	-	-	-	-	37.5	35.6	29.1	33.5	33.6	-
Stability (P), mm	100	-	-	-	-	-	90	95	78	85	82	-
Distensibility (L), mm	85	-	-	-	-	-	92	80	87	92	103	-
Configuration ratio (P/L)	1.18	-	-	-	-	-	0.98	1.19	0.90	0.92	0.80	-
<b>MIXOGRAM</b>												
Peak time, min	2.3	-	-	-	-	-	2.2	2.7	2.4	2.0	2.0	-
<b>100g BAKING TEST</b>												
Loaf volume, cm <sup>3</sup>	925	-	-	-	-	-	935	935	815	910	965	-
Evaluation	0	-	-	-	-	-	0	0	0	0	0	-

# RHEOLOGICAL GRAPHS PER PRODUCTION REGION

## MIXOGRAM

7

10

## FARINOGRAM

7

10

## EXTENSOGRAM

7

10

## ALVEOGRAM

7

10

**SOUTH AFRICAN  
IRRIGATION WHEAT  
Vaal and Orange river area**

**MAINLY IRRIGATION WHEAT  
North-West Province**

PRODUCTION REGION	(11) Vaalharts	(12) North-West Western Region										
Intake silos	Barkly-West Hartswater Jan Kemp Magogong Taung	Bloubank	Vryburg	Buhrmannsdrif	Vryhof	Kameel						
<b>WHEAT</b>												
Protein (12% mb), %	ave 11.9	min 10.9	max 12.6	stdev 0.53	ave 12.8	min 11.4	max 14.8					
Falling number, sec	395	381	423	12.7	405	375	440					
1000 Kernel mass (13% mb), g	41.9	31.8	46.2	3.34	38.8	34.1	40.6					
Hectolitre mass (dirty), kg/hl	80.9	79.2	82.4	0.99	81.4	80.3	81.9					
Screenings (<1.8mm), %	2.30	1.46	3.32	0.55	1.32	0.87	1.99					
Total damaged kernels, %	0.30	0.06	0.66	0.16	0.37	0.22	0.64					
<b>Number of samples</b>	14				5							
<b>CULTIVARS</b>												
cultivars with highest % occurrence	Duzi	36.0		SST 835	39.0							
	SST 835	28.3		SST 843	36.0							
	SST 843	16.6		Duzi	15.4							
	PAN 3434	6.5		CRN 826	8.4							
	Baviaans	6.0		SST 825	1.2							
<b>Number of samples</b>	14				5							
<b>MIXOGRAM (Quadromat)</b>												
Peak time, min	ave 2.7	min 2.5	max 2.8	stdev 0.13	ave 3.1	min 2.4	max 4.3					
Tail height (6min), mm	49	47	52	1.44	55	49	67					
<b>Number of samples</b>	14				5							
<b>BÜHLER EXTRACTION, %</b>												
<b>FLOUR</b>	B1 76.8	B2 76.5	B3 76.3	B4 75.9	UT -	COW -	B1 76.1	B2 76.0	B3 -	B4 -	UT -	COW -
<b>GLUTEN</b>												
Wet gluten (14% mb), %	11.5	10.5	9.7	10.9	-	-	13.1	10.7	-	-	-	-
Colour, KJ	-2.3	-2.5	-2.6	-2.3	-	-	-2.5	-2.6	-	-	-	-
<b>FARINOGRAM</b>												
Water absorption (14% mb), %	63.2	62.6	62.9	64.2	-	-	64.8	62.6	-	-	-	-
Development time, min	5.5	4.7	4.8	4.5	-	-	10.2	4.8	-	-	-	-
Stability, min	6.2	5.5	6.3	5.1	-	-	12.4	6.1	-	-	-	-
Mixing tolerance index, BU	38	52	43	48	-	-	26	43	-	-	-	-
<b>EXTENSOGRAM (45 min pull)</b>												
Area, cm <sup>2</sup>	89	79	81	76	-	-	145	96	-	-	-	-
Maximum height, BU	290	300	310	285	-	-	400	335	-	-	-	-
Extensibility, mm	218	179	170	187	-	-	252	199	-	-	-	-
<b>ALVEOGRAM</b>												
Strength (S), cm <sup>2</sup>	35.6	31.6	28.0	32.0	-	-	57.6	31.2	-	-	-	-
Stability (P), mm	78	81	92	87	-	-	103	80	-	-	-	-
Distensibility (L), mm	114	92	64	86	-	-	113	88	-	-	-	-
Configuration ratio (P/L)	0.68	0.88	1.44	1.01	-	-	0.91	0.91	-	-	-	-
<b>MIXOGRAM</b>												
Peak time, min	2.3	2.3	2.3	2.2	-	-	3.0	2.5	-	-	-	-
<b>100g BAKING TEST</b>												
Loaf volume, cm <sup>3</sup>	900	820	810	825	-	-	900	875	-	-	-	-
Evaluation	0	0	0	1	-	-	3	0	-	-	-	-

# RHEOLOGICAL GRAPHS PER PRODUCTION REGION

## MIXOGRAM

11

12

## FARINOGRAM

11

12

## EXTENSOGRAM

11

12

## ALVEOGRAM

11

12

**SOUTH AFRICAN**  
**MAINLY IRRIGATION WHEAT**  
**North-West Province**

PRODUCTION REGION	(14) North-West Southern Region				(15) North-West South-Eastern Region							
Intake silos	Amalia				Bloemhof							
	Barberspan				Christiana							
	Delareyville				Hertzogville							
	Excelsior				Hoopstad							
	Geysdorp				Kingwood							
	Hallat's Hope											
	Migdal											
	Nooitgedacht											
	Schweizer-Reneke											
	Taaibospan											
<b>WHEAT</b>												
Protein (12% mb), %	ave 11.6	min 10.8	max 13.7	stdev 1.43	ave 11.9	min 10.5	max 13.0	stdev 0.75				
Falling number, sec	415	410	420	4.20	375	267	460	74.42				
1000 Kernel mass (13% mb), g	39.1	26.7	47.5	9.22	38.1	33.3	46.8	3.98				
Hectolitre mass (dirty), kg hl	79.7	74.1	84.3	4.20	81.2	79.7	82.8	0.98				
Screenings (<1.8mm), %	4.28	3.18	6.54	1.57	1.19	0.44	2.37	0.78				
Total damaged kernels, %	0.14	0.00	0.24	0.10	0.08	0.00	0.20	0.08				
<b>Number of samples</b>	<b>4</b>				<b>9</b>							
<b>CULTIVARS</b>												
	SST 835	39.3			SST 835	23.4						
cultivars	Baviaans	26.3			PAN 3120	15.6						
with highest % occurrence	PAN 3434	11.0			Gariep	15.1						
	SST 843	9.3			Duzi	12.4						
	CRN 826	7.8			SST 843	9.7						
<b>Number of samples</b>	<b>4</b>				<b>9</b>							
<b>MIXOGRAM (Quadromat)</b>												
Peak time, min	ave 2.9	min 2.3	max 3.3	stdev 0.48	ave 3.0	min 2.2	max 3.7	stdev 0.46				
Tail height (6min), mm	50	46	54	3.32	49	47	52	2.07				
<b>Number of samples</b>	<b>4</b>				<b>9</b>							
<b>BÜHLER EXTRACTION, %</b>	B1	B2	B3	B4	UT	COW	B1	B2				
	-	-	-	76.0	74.3	-	74.4	74.9				
							74.1	-				
							-	-				
<b>FLOUR</b>												
Protein (12% mb), %	-	-	-	9.8	11.3	-	11.9	10.4				
Colour, KJ	-	-	-	-2.7	-2.5	-	-2.2	-2.4				
							-2.6	-				
							-	-				
<b>GLUTEN</b>												
Wet gluten (14% mb), %	-	-	-	27.6	30.2	-	33.3	28.7				
Dry gluten (14% mb), %	-	-	-	9.5	10.6	-	11.7	9.8				
							9.2	-				
							-	-				
<b>FARINOGRAM</b>												
Water absorption (14% mb), %	-	-	-	62.2	61.7	-	65.7	63.9				
Development time, min	-	-	-	4.5	6.0	-	5.7	4.9				
Stability, min	-	-	-	5.9	7.8	-	7.2	6.2				
Mixing tolerance index, BU	-	-	-	47	41	-	34	43				
							21	-				
							-	-				
<b>EXTENSOGRAM (45 min pull)</b>												
Area, cm <sup>2</sup>	-	-	-	84	123	-	97	84				
Maximum height, BU	-	-	-	325	400	-	340	320				
Extensibility, mm	-	-	-	178	213	-	201	186				
							177	-				
							-	-				
<b>ALVEOGRAM</b>												
Strength (S), cm <sup>2</sup>	-	-	-	29.4	39.3	-	39.6	33.3				
Stability (P), mm	-	-	-	80	81	-	103	94				
Distensibility (L), mm	-	-	-	83	116	-	79	73				
Configuration ratio (P/L)	-	-	-	0.96	0.70	-	1.30	1.29				
							1.18	-				
							-	-				
<b>MIXOGRAM</b>												
Peak time, min	-	-	-	2.6	2.7	-	2.5	2.5				
							3.5	-				
							-	-				
<b>100g BAKING TEST</b>												
Loaf volume, cm <sup>3</sup>	-	-	-	810	875	-	895	830				
Evaluation	-	-	-	0	0	-	1	0				
							3	-				
							-	-				

# RHEOLOGICAL GRAPHS PER PRODUCTION REGION

## MIXOGRAM

14

15

## FARINOGRAM

14

15

## EXTENSOGRAM

14

15

## ALVEOGRAM

14

15

**SOUTH AFRICAN  
MAINLY IRRIGATION WHEAT  
North-West Province**

PRODUCTION REGION	(16) North-West Central Eastern Region				(17) North-West Central Northern Region (Ottosdal)							
Intake silos	Bamboesspruit Klerksdorp Leeudoringstad Makwassie Regina Strydpoort Wolmaranstad				Bospoort Lethabong (Hartbeesfontein) Kleincharts Melliodora Ottosdal Rostrataville Vermaas Werda							
<b>WHEAT</b>												
Protein (12% mb), %	<b>ave</b> 11.8	<b>min</b> 11.2	<b>max</b> 12.5	<b>stdev</b> 0.67	<b>ave</b> 11.9	<b>min</b> 10.2	<b>max</b> 13.0	<b>stdev</b> 0.94				
Falling number, sec	434	428	444	8.96	416	371	455	32.73				
1000 Kernel mass (13% mb), g	39.8	37.5	41.4	2.03	40.8	34.3	50.7	5.85				
Hectolitre mass (dirty), kg/hl	78.9	76.9	80.1	1.74	80.6	79.2	82.2	1.12				
Screenings (<1.8mm), %	2.17	1.88	2.54	0.34	1.12	0.52	1.40	0.29				
Total damaged kernels, %	0.17	0.12	0.20	0.04	0.19	0.06	0.32	0.11				
<b>Number of samples</b>	<b>3</b>				<b>8</b>							
<b>CULTIVARS</b>												
cultivars with highest % occurrence	SST 835 SST 843 Duzi CRN 826	42.0 33.7 15.3 9.0			SST 835 Baviaans SST 843 Duzi CRN 826	48.6 20.8 15.4 6.8 5.1						
<b>Number of samples</b>	<b>3</b>				<b>8</b>							
<b>MIXOGRAM (Quadromat)</b>												
Peak time, min	<b>ave</b> 2.6	<b>min</b> 2.3	<b>max</b> 3.2	<b>stdev</b> 0.52	<b>ave</b> 2.9	<b>min</b> 2.3	<b>max</b> 3.3	<b>stdev</b> 0.36				
Tail height (6min), mm	46	44	48	2.08	50	42	54	3.81				
<b>Number of samples</b>	<b>3</b>				<b>8</b>							
<b>BÜHLER EXTRACTION, %</b>	<b>B1</b> 75.2	<b>B2</b> 75.7	<b>B3</b> -	<b>B4</b> -	<b>UT</b> -	<b>COW</b> -	<b>B1</b> 75.4	<b>B2</b> 77.4	<b>B3</b> 76.3	<b>B4</b> -	<b>UT</b> -	<b>COW</b> -
<b>FLOWER</b>												
Protein (12% mb), %	11.8	10.1	-	-	-	-	11.7	10.2	8.7	-	-	-
Colour, KJ	-2.1	-2.3	-	-	-	-	-2.2	-2.6	-2.7	-	-	-
<b>GLUTEN</b>												
Wet gluten (14% mb), %	33.2	26.9	-	-	-	-	31.8	28.1	21.7	-	-	-
Dry gluten (14% mb), %	11.3	9.2	-	-	-	-	11.2	9.5	7.8	-	-	-
<b>FARINOGRAM</b>												
Water absorption (14% mb), %	62.5	60.5	-	-	-	-	63.3	61.9	57.5	-	-	-
Development time, min	5.0	4.8	-	-	-	-	5.7	4.5	3.0	-	-	-
Stability, min	5.4	5.5	-	-	-	-	8.1	6.4	7.4	-	-	-
Mixing tolerance index, BU	48	47	-	-	-	-	37	49	26	-	-	-
<b>EXTENOGRAM (45 min pull)</b>												
Area, cm <sup>2</sup>	-	87	-	-	-	-	119	86	76	-	-	-
Maximum height, BU	-	335	-	-	-	-	375	345	405	-	-	-
Extensibility, mm	-	177	-	-	-	-	219	175	135	-	-	-
<b>ALVEOGRAM</b>												
Strength (S), cm <sup>2</sup>	34.1	26.6	-	-	-	-	41.9	30.9	22.2	-	-	-
Stability (P), mm	74	73	-	-	-	-	87	90	70	-	-	-
Distensibility (L), mm	119	86	-	-	-	-	111	73	63	-	-	-
Configuration ratio (P/L)	0.62	0.85	-	-	-	-	0.78	1.23	1.11	-	-	-
<b>MIXOGRAM</b>												
Peak time, min	2.3	2.5	-	-	-	-	2.7	2.7	2.9	-	-	-
<b>100g BAKING TEST</b>												
Loaf volume, cm <sup>3</sup>	905	805	-	-	-	-	825	800	775	-	-	-
Evaluation	0	0	-	-	-	-	2	0	0	-	-	-

# RHEOLOGICAL GRAPHS PER PRODUCTION REGION

## MIXOGRAM

16

17

## FARINOGRAM

16

17

## EXTENSOGRAM

16

17

## ALVEOGRAM

16

17

**SOUTH AFRICAN**  
**MAINLY IRRIGATION WHEAT**  
**North-West Province**

PRODUCTION REGION	(18) North-West Central Region (Ventersdorp)				(19) North-West Central Region (Lichtenburg)									
	Intake silos	Bodenstein	Grootpan		Buckingham	Halfpad		Coligny	Hibernia	Ense spruit	Lichtenburg	Makokskraal	Lottiehalte	Potchefstroom
<b>WHEAT</b>														
Protein (12% mb), %	<b>ave</b> 12.7	<b>min</b> 11.2	<b>max</b> 14.2	<b>stdev</b> 2.12	<b>ave</b> 11.9	<b>min</b> 10.3	<b>max</b> 12.8	<b>stdev</b> 0.79						
Falling number, sec	386	360	411	36.06	389	333	413	26.17						
1000 Kernel mass (13% mb), g	38.7	38.6	38.7	0.07	37.5	36.0	39.7	1.15						
Hectolitre mass (dirty), kg/hl	79.6	78.5	80.6	1.48	81.2	79.6	82.1	0.92						
Screenings (<1.8mm), %	1.31	1.22	1.40	0.13	1.88	1.48	2.15	0.27						
Total damaged kernels, %	0.35	0.30	0.40	0.07	0.26	0.12	0.60	0.15						
<b>Number of samples</b>	<b>2</b>				<b>8</b>									
<b>CULTIVARS</b>														
cultivars with highest % occurrence	SST 843				SST 835				52.5					
	SST 88				SST 843				37.0					
					CRN 826				6.4					
					Olifants				3.4					
					SST 876				0.8					
<b>Number of samples</b>	<b>2</b>				<b>8</b>									
<b>MIXOGRAM (Quadromat)</b>														
Peak time, min	<b>ave</b> 4.2	<b>min</b> 3.2	<b>max</b> 5.2	<b>stdev</b> 1.41	<b>ave</b> 3.4	<b>min</b> 2.8	<b>max</b> 5.0	<b>stdev</b> 0.81						
Tail height (6min), mm	59	48	69	14.85	52	43	60	5.01						
<b>Number of samples</b>	<b>2</b>				<b>8</b>									
<b>BÜHLER EXTRACTION, %</b>	<b>B1</b> 75.4	<b>B2</b> 75.4	<b>B3</b> -	<b>B4</b> -	<b>UT</b> -	<b>COW</b> -	<b>B1</b> 75.6	<b>B2</b> 75.7	<b>B3</b> 75.6	<b>B4</b> -	<b>UT</b> -	<b>COW</b> -		
<b>FLOUR</b>														
Protein (12% mb), %	13.7	10.1	-	-	-	-	11.9	10.9	9.5	-	-	-		
Colour, KJ	-2.3	-2.2	-	-	-	-	-2.3	-2.4	-2.1	-	-	-		
<b>GLUTEN</b>														
Wet gluten (14% mb), %	33.1	26.0	-	-	-	-	31.7	29.0	25.9	-	-	-		
Dry gluten (14% mb), %	12.4	9.6	-	-	-	-	11.0	10.1	9.0	-	-	-		
<b>FARINOGRAM</b>														
Water absorption (14% mb), %	64.9	60.5	-	-	-	-	63.8	63.0	62.4	-	-	-		
Development time, min	16.4	4.7	-	-	-	-	7.8	6.5	5.5	-	-	-		
Stability, min	17.6	9.0	-	-	-	-	12.7	8.2	7.2	-	-	-		
Mixing tolerance index, BU	-	25	-	-	-	-	20	35	34	-	-	-		
<b>EXTENSOGRAM (45 min pull)</b>														
Area, cm <sup>2</sup>	139	108	-	-	-	-	148	119	-	-	-	-		
Maximum height, BU	475	435	-	-	-	-	430	390	-	-	-	-		
Extensibility, mm	204	184	-	-	-	-	239	217	-	-	-	-		
<b>ALVEOGRAM</b>														
Strength (S), cm <sup>2</sup>	81.0	33.5	-	-	-	-	51.5	37.6	26.9	-	-	-		
Stability (P), mm	136	85	-	-	-	-	96	93	94	-	-	-		
Distensibility (L), mm	102	83	-	-	-	-	112	84	55	-	-	-		
Configuration ratio (P/L)	1.33	1.02	-	-	-	-	0.86	1.11	1.71	-	-	-		
<b>MIXOGRAM</b>														
Peak time, min	4.8	3.2	-	-	-	-	3.2	2.7	2.7	-	-	-		
<b>100g BAKING TEST</b>														
Loaf volume, cm <sup>3</sup>	980	850	-	-	-	-	760	820	720	-	-	-		
Evaluation	2	0	-	-	-	-	5	1	2	-	-	-		

# RHEOLOGICAL GRAPHS PER PRODUCTION REGION

## MIXOGRAM

18

19

## FARINOGRAM

18

19

## EXTENSOGRAM

18

19

## ALVEOGRAM

18

19

**SOUTH AFRICAN  
MAINLY IRRIGATION WHEAT  
North-West Province**

PRODUCTION REGION	(20) North-West Eastern Region						(21) Free State North-Western Region (Viljoenskroon)					
Intake silos	Battery	Boons	Brits	Derby	Koster	Rustenburg	Swartruggens	Syferbult	Attie	Groenebloem	Heuningspruit	Koppies
									Rooiwal	Vierfontein	Viljoenskroon	Vrededorf
									Weiveld			
<b>WHEAT</b>	<b>ave</b>	<b>min</b>	<b>max</b>	<b>stdev</b>	<b>ave</b>	<b>min</b>	<b>max</b>	<b>stdev</b>				
Protein (12% mb), %	11.6	10.6	13.2	0.81	11.8	10.6	13.5	1.07				
Falling number, sec	411	357	521	40.75	397	351	437	30.58				
1000 Kernel mass (13% mb), g	39.7	33.9	46.1	2.75	37.8	32.7	42.7	3.76				
Hectolitre mass (dirty), kg/hl	80.3	73.0	84.0	2.94	81.7	79.8	82.6	1.19				
Screenings (<1.8mm), %	2.42	0.27	5.84	1.86	1.91	1.49	2.52	0.42				
Total damaged kernels, %	0.31	0.00	0.88	0.25	0.18	0.08	0.32	0.09				
<b>Number of samples</b>	<b>15</b>				<b>5</b>							
<b>CULTIVARS</b>												
cultivars with highest % occurrence	SST 835			40.3			SST 835			50.8		
	Duzi			17.3			SST 843			25.6		
	SST 843			16.5			Elands			16.4		
	Kariega			12.8			CRN 826			4.0		
	CRN 826			4.7			SST 347, Baviaans			1.6		
<b>Number of samples</b>	<b>15</b>				<b>5</b>							
<b>MIXOGRAM (Quadromat)</b>	<b>ave</b>	<b>min</b>	<b>max</b>	<b>stdev</b>	<b>ave</b>	<b>min</b>	<b>max</b>	<b>stdev</b>				
Peak time, min	3.1	2.3	4.8	0.72	3.1	2.3	4.5	0.87				
Tail height (6min), mm	50	43	65	5.07	52	47	59	5.55				
<b>Number of samples</b>	<b>15</b>				<b>5</b>							
<b>BÜHLER EXTRACTION, %</b>	<b>B1</b>	<b>B2</b>	<b>B3</b>	<b>B4</b>	<b>UT</b>	<b>COW</b>	<b>B1</b>	<b>B2</b>	<b>B3</b>	<b>B4</b>	<b>UT</b>	<b>COW</b>
	75.8	76.5	76.4	-	76.0	77.0	74.8	76.0	75.2	-	-	-
<b>FLOUR</b>												
Protein (12% mb), %	11.6	10.6	9.3	-	10.4	10.2	13.1	10.9	9.6	-	-	-
Colour, KJ	-2.3	-2.3	-2.2	-	-2.5	-1.8	-2.2	-1.6	-2.0	-	-	-
<b>GLUTEN</b>												
Wet gluten (14% mb), %	30.8	29.5	24.6	-	28.8	28.0	32.5	30.3	25.2	-	-	-
Dry gluten (14% mb), %	11.2	10.3	8.5	-	9.5	9.5	12.4	10.5	8.9	-	-	-
<b>FARINOGRAM</b>												
Water absorption (14% mb), %	61.5	62.3	59.7	-	62.2	62.0	63.4	64.3	61.4	-	-	-
Development time, min	7.0	6.0	5.5	-	5.2	4.2	10.5	5.5	5.7	-	-	-
Stability, min	10.5	7.7	7.2	-	5.2	4.3	16.5	5.9	6.8	-	-	-
Mixing tolerance index, BU	30	44	42	-	50	56	15	46	47	-	-	-
<b>EXTENSOGRAM (45 min pull)</b>												
Area, cm <sup>2</sup>	148	103	88	-	75	74	126	99	101	-	-	-
Maximum height, BU	475	400	370	-	285	260	465	360	435	-	-	-
Extensibility, mm	218	185	176	-	188	193	197	194	160	-	-	-
<b>ALVEOGRAM</b>												
Strength (S), cm <sup>2</sup>	45.1	35.2	28.9	-	29.1	24.6	57.6	36.7	28.3	-	-	-
Stability (P), mm	89	89	75	-	74	69	104	95	92	-	-	-
Distensibility (L), mm	106	85	89	-	99	95	104	84	60	-	-	-
Configuration ratio (P/L)	0.84	1.05	0.84	-	0.75	0.73	1.00	1.13	1.53	-	-	-
<b>MIXOGRAM</b>												
Peak time, min	3.3	2.8	2.8	-	2.3	2.3	3.8	2.5	2.8	-	-	-
<b>100g BAKING TEST</b>												
Loaf volume, cm <sup>3</sup>	810	805	730	-	740	795	775	790	770	-	-	-
Evaluation	3	1	1	-	3	1	6	2	0	-	-	-

# RHEOLOGICAL GRAPHS PER PRODUCTION REGION

## MIXOGRAM

20

21

## FARINOGRAM

20

21

## EXTENSOGRAM

20

21

## ALVEOGRAM

20

21

## **SOUTH AFRICAN SUMMER RAINFALL WHEAT (AND IRRIGATION) Free State Province (Central)**

PRODUCTION REGION	(26) Free State South-Eastern Region (Senekal)				(27) Free State Northern Region									
Intake silos	Arlington	Kaallaagte	Libertas	Marquard	Meets	Monte Video	Senekal	Steynsrus	Gottenburg	Heilbron	Hooge	Mooigeleë	Petrus Steyn	Wolwehoek
<b>WHEAT</b>														
Protein (12% mb), %	ave 13.2	min 12.2	max 14.6	stdev 0.63	ave 13.6	min 11.7	max 14.7	stdev 1.19						
Falling number, sec	305	104	466	104.21	328	114	498	112.80						
1000 Kernel mass (13% mb), g	39.1	35.4	42.2	1.98	37.8	35.6	40.1	1.86						
Hectolitre mass (dirty), kg/hl	80.2	74.3	82.2	2.40	79.1	75.7	81.0	1.60						
Screenings (<1.8mm), %	1.22	0.47	2.49	0.69	1.10	0.24	1.81	0.57						
Total damaged kernels, %	0.88	0.00	6.48	1.71	0.85	0.50	1.24	0.22						
<b>Number of samples</b>	13				8									
<b>CULTIVARS</b>														
cultivars with highest % occurrence	Elands				Elands				50.4					
	Matlabas				SST 347				15.7					
	SST 356				Matlabas				6.3					
	PAN 3120				SST 835				5.7					
	Gariep				PAN 3120				5.6					
<b>Number of samples</b>	13				8									
<b>MIXOGRAM (Quadromat)</b>														
Peak time, min	ave 3.7	min 2.8	max 4.7	stdev 0.62	ave 3.7	min 3.2	max 4.5	stdev 0.40						
Tail height (6min), mm	55	51	62	3.08	56	51	62	3.98						
<b>Number of samples</b>	13				8									
<b>BÜHLER EXTRACTION, %</b>	B1 74.8	B2 -	B3 -	B4 -	UT 72.8	COW 74.7	B1 74.7	B2 74.7	B3 -	B4 -	UT -	COW 73.5		
<b>FLOUR</b>														
Protein (12% mb), %	12.3	-	-	-	12.7	12.3	13.3	10.7	-	-	-	14.1		
Colour, KJ	-0.9	-	-	-	1.7	2.7	0.3	-1.4	-	-	-	2.8		
<b>GLUTEN</b>														
Wet gluten (14% mb), %	32.8	-	-	-	33.0	29.9	35.6	26.9	-	-	-	37.2		
Dry gluten (14% mb), %	11.5	-	-	-	11.6	11.0	13.2	9.3	-	-	-	13.3		
<b>FARINOGRAM</b>														
Water absorption (14% mb), %	65.6	-	-	-	64.5	63.9	66.5	62.3	-	-	-	67.2		
Development time, min	8.0	-	-	-	3.0	2.0	7.2	6.7	-	-	-	8.8		
Stability, min	11.3	-	-	-	10.6	7.6	11.8	10.4	-	-	-	10.9		
Mixing tolerance index, BU	28	-	-	-	24	33	20	25	-	-	-	32		
<b>EXTENSOGRAM (45 min pull)</b>														
Area, cm <sup>2</sup>	110	-	-	-	90	110	140	106	-	-	-	119		
Maximum height, BU	390	-	-	-	480	440	445	450	-	-	-	365		
Extensibility, mm	193	-	-	-	143	183	215	163	-	-	-	225		
<b>ALVEOGRAM</b>														
Strength (S), cm <sup>2</sup>	42.7	-	-	-	39.6	45.0	50.3	35.2	-	-	-	48.6		
Stability (P), mm	122	-	-	-	134	120	128	107	-	-	-	106		
Distensibility (L), mm	61	-	-	-	46	63	69	62	-	-	-	85		
Configuration ratio (P/L)	2.00	-	-	-	2.91	1.90	1.86	1.73	-	-	-	1.25		
<b>MIXOGRAM</b>														
Peak time, min	3.1	-	-	-	3.5	4.4	3.3	3.3	-	-	-	3.4		
<b>100g BAKING TEST</b>														
Loaf volume, cm <sup>3</sup>	860	-	-	-	720	870	940	740	-	-	-	1035		
Evaluation	2	-	-	-	6	2	2	3	-	-	-	1		

# RHEOLOGICAL GRAPHS PER PRODUCTION REGION

## MIXOGRAM

26

27

## FARINOGRAM

26

27

## EXTENSOGRAM

26

27

## ALVEOGRAM

26

27

**SOUTH AFRICAN**  
**SUMMER RAINFALL WHEAT (AND IRRIGATION)**  
**Free State Province (North-Western)**

PRODUCTION REGION	(22) Free-State North-Western Region (Bothaville)				(23) Free-State North-Western Region (Bultfontein)							
Intake silos	Allanridge Bothaville Mirage Odendaalsrus Schoonspruit Schuttesdraai	Bultfontein Losdoorns Protespan Tierfontein Wesselsbron Willemsrust										
<b>WHEAT</b>												
Protein (12% mb), %	<b>ave</b> 11.7	<b>min</b> 9.5	<b>max</b> 13.6	<b>stdev</b> 1.61	<b>ave</b> 11.4	<b>min</b> 9.7	<b>max</b> 13.6	<b>stdev</b> 1.13				
Falling number, sec	357	316	404	39.78	362	285	510	57.39				
1000 Kernel mass (13% mb), g	36.3	34.8	38.3	1.47	36.5	30.9	41.8	3.39				
Hectolitre mass (dirty), kg/hl	81.4	80.0	82.4	0.87	81.4	77.7	84.1	1.49				
Screenings (<1.8mm), %	1.51	0.85	2.03	0.40	1.75	0.23	3.17	0.77				
Total damaged kernels, %	0.16	0.00	0.26	0.10	0.27	0.08	0.80	0.15				
<b>Number of samples</b>	<b>6</b>				<b>22</b>							
<b>CULTIVARS</b>												
	PAN 3120	34.3			PAN 3120	29.1						
cultivars	SST 835	16.8			SST 835	20.4						
with highest % occurrence	SST 347	10.0			CRN 826	11.0						
	PAN 3355	9.2			PAN 3118	10.2						
	Gariep	8.3			Gariep	7.4						
<b>Number of samples</b>	<b>6</b>				<b>22</b>							
<b>MIXOGRAM (Quadromat)</b>												
Peak time, min	<b>ave</b> 3.4	<b>min</b> 2.5	<b>max</b> 4.7	<b>stdev</b> 0.88	<b>ave</b> 3.1	<b>min</b> 2.4	<b>max</b> 3.8	<b>stdev</b> 0.42				
Tail height (6min), mm	51	42	57	5.50	50	43	57	4.15				
<b>Number of samples</b>	<b>6</b>				<b>22</b>							
<b>BÜHLER EXTRACTION, %</b>	<b>B1</b> 74.1	<b>B2</b> 72.6	<b>B3</b> 73.3	<b>B4</b> 72.0	<b>UT</b> -	<b>COW</b> -	<b>B1</b> 73.8	<b>B2</b> 73.2	<b>B3</b> 73.8	<b>B4</b> 73.7	<b>UT</b> -	<b>COW</b> -
<b>FLOUR</b>												
Protein (12% mb), %	12.2	9.9	9.1	7.8	-	-	11.8	10.4	9.2	8.8	-	-
Colour, KJ	-2.0	-2.3	-1.6	-2.2	-	-	-2.3	-2.4	-2.3	-2.1	-	-
<b>GLUTEN</b>												
Wet gluten (14% mb), %	34.2	26.8	21.3	-	-	-	31.0	27.7	24.4	21.9	-	-
Dry gluten (14% mb), %	12.0	9.0	7.6	-	-	-	11.2	10.0	8.7	7.6	-	-
<b>FARINOGRAM</b>												
Water absorption (14% mb), %	67.1	62.8	63.4	62.6	-	-	63.0	63.4	62.0	61.5	-	-
Development time, min	7.2	5.5	1.9	1.7	-	-	7.2	6.0	5.5	3.7	-	-
Stability, min	8.4	7.7	1.6	1.3	-	-	10.1	8.0	8.6	7.4	-	-
Mixing tolerance index, BU	35	42	62	73	-	-	32	38	33	33	-	-
<b>EXTENSOGRAM (45 min pull)</b>												
Area, cm <sup>2</sup>	108	-	-	-	-	-	119	95	80	80	-	-
Maximum height, BU	330	-	-	-	-	-	410	390	355	380	-	-
Extensibility, mm	223	-	-	-	-	-	202	172	162	143	-	-
<b>ALVEOGRAM</b>												
Strength (S), cm <sup>2</sup>	43.9	32.6	28.7	21.3	-	-	40.8	34.9	29.7	26.3	-	-
Stability (P), mm	110	102	129	125	-	-	96	107	104	108	-	-
Distensibility (L), mm	83	64	35	28	-	-	85	63	54	41	-	-
Configuration ratio (P/L)	1.33	1.59	3.69	4.46	-	-	1.13	1.70	1.93	2.63	-	-
<b>MIXOGRAM</b>												
Peak time, min	2.7	2.5	3.0	3.7	-	-	3.0	2.8	2.8	2.8	-	-
<b>100g BAKING TEST</b>												
Loaf volume, cm <sup>3</sup>	815	740	640	570	-	-	795	735	655	685	-	-
Evaluation	4	2	4	5	-	-	4	3	4	2	-	-

# RHEOLOGICAL GRAPHS PER PRODUCTION REGION

## MIXOGRAM

22

23

## FARINOGRAM

22

23

## EXTENSOGRAM

22

23

## ALVEOGRAM

22

23

**SOUTH AFRICAN**  
**SUMMER RAINFALL WHEAT (AND IRRIGATION)**  
**Free State Province (Eastern)**

PRODUCTION REGION	(25) Free State South-Western Region (Bethlehem)				(28) Free State Eastern Region							
	Bethlehem	Clocolan	De Wetshoek	Ficksburg	Afrikaskop	Tweeling	Villiers	Vrede				
Intake silos	Fouriesburg	Marseilles	Modderpoort	Slabberts	Cornelia	Warden	Windfield					
	Zastron				Daniëlsrus							
					Eeram							
					Frankfort							
					Harrismith							
					Jim Fouché							
					Kransfontein							
					Memel							
					Reitz							
<b>WHEAT</b>												
Protein (12% mb), %	ave 13.1	min 11.2	max 15.3	stdev 0.79	ave 13.1	min 11.5	max 15.3	stdev 0.85				
Falling number, sec	240	89	379	84.88	298	63	439	92.58				
1000 Kernel mass (13% mb), g	39.4	31.8	43.4	3.04	39.5	35.0	45.3	2.41				
Hectolitre mass (dirty), kg/hl	78.2	72.4	82.0	2.42	78.8	72.9	83.3	2.46				
Screenings (<1.8mm), %	1.72	0.22	4.74	0.98	1.24	0.26	3.72	0.77				
Total damaged kernels, %	0.74	0.08	3.64	0.76	0.58	0.08	5.68	0.99				
<i>Number of samples</i>	<b>25</b>				<b>31</b>							
<b>CULTIVARS</b>												
cultivars	Elands SST 356				Elands Matlabas							
with highest % occurrence	28.7 25.5				37.5 18.2							
	Matlabas SST 835				SST 835 SST 356							
	7.6 5.4				13.0 Duzi							
<i>Number of samples</i>	<b>25</b>				<b>31</b>							
<b>MIXOGRAM (Quadromat)</b>												
Peak time, min	ave 3.8	min 2.6	max 6.8	stdev 0.99	ave 3.4	min 2.4	max 4.5	stdev 0.48				
Tail height (6min), mm	54	42	65	4.60	54	46	61	3.90				
<i>Number of samples</i>	<b>25</b>				<b>31</b>							
<b>BÜHLER EXTRACTION, %</b>	<b>B1</b> 73.7	<b>B2</b> 73.8	<b>B3</b> 73.1	<b>B4</b> 74.5	<b>UT</b> 74.0	<b>COW</b> 73.1	<b>B1</b> 74.0	<b>B2</b> 74.5	<b>B3</b> 73.4	<b>B4</b> 72.7	<b>UT</b> 73.5	<b>COW</b> 72.9
<b>FLOUR</b>												
Protein (12% mb), %	12.3	11.7	11.9	12.0	11.2	11.9	11.8	11.1	12.9	11.1	14.3	12.5
Colour, KJ	-1.6	-1.1	-0.3	-1.2	-0.1	0.7	-1.4	-0.5	1.6	1.8	-0.6	0.7
<b>GLUTEN</b>												
Wet gluten (14% mb), %	33.1	31.3	30.7	32.3	28.6	31.4	31.6	29.3	35.5	28.6	39.2	33.6
Dry gluten (14% mb), %	11.7	10.8	11.2	11.3	10.0	11.2	10.9	10.0	12.5	9.4	14.2	11.9
<b>FARINOGRAM</b>												
Water absorption (14% mb), %	65.1	66.2	63.7	65.1	64.4	63.4	65.2	63.4	67.8	64.0	67.2	65.3
Development time, min	7.5	6.2	6.2	8.3	2.4	3.2	7.0	6.2	7.0	2.2	6.7	4.2
Stability, min	13.4	9.2	11.2	11.1	7.8	6.4	9.4	9.0	11.6	9.0	14.7	7.9
Mixing tolerance index, BU	18	22	21	29	29	39	32	36	22	16	21	33
<b>EXTENSOGRAM (45 min pull)</b>												
Area, cm <sup>2</sup>	117	98	105	122	126	124	92	98	110	89	121	103
Maximum height, BU	420	350	405	410	435	375	350	410	410	350	365	400
Extensibility, mm	198	188	185	210	195	230	188	172	195	185	236	184
<b>ALVEOGRAM</b>												
Strength (S), cm <sup>2</sup>	50.3	37.3	42.8	50.6	48.0	42.8	42.5	37.3	49.2	35.5	67.9	43.9
Stability (P), mm	113	107	105	111	124	89	113	101	126	94	110	105
Distensibility (L), mm	86	67	74	92	70	93	72	74	74	74	121	78
Configuration ratio (P/L)	1.31	1.60	1.42	1.21	1.77	0.96	1.57	1.36	1.70	1.27	0.91	1.35
<b>MIXOGRAM</b>												
Peak time, min	3.3	2.4	3.5	3.3	3.8	3.5	2.7	3.2	3.0	3.3	3.2	3.2
<b>100g BAKING TEST</b>												
Loaf volume, cm <sup>3</sup>	880	850	800	900	800	925	835	825	935	935	1040	890
Evaluation	2	2	4	1	2	0	2	1	1	0	1	2

# RHEOLOGICAL GRAPHS PER PRODUCTION REGION

## MIXOGRAM

25

28

## FARINOGRAM

25

28

## EXTENSOGRAM

25

28

## ALVEOGRAM

25

28

**SOUTH AFRICAN**  
**SUMMER RAINFALL WHEAT (AND IRRIGATION)**  
**Free State Province (South-Western)**

**OTHER SUMMER RAINFALL AND  
IRRIGATION WHEAT**  
**Mpumalanga**

PRODUCTION REGION	(24) Free State Central Region				(29) Mpumalanga Southern Region									
	Bloemfontein	Welgeleë			Balfour									
Intake silos	Brandfort	Winburg			Greylingstad									
	De Brug				Grootvlei									
	Geneva				Harvard									
	Hennenman				Holmdene									
	Koffiefontein				Leeuspruit									
	Kroonstad				Platrand									
	Petrusburg				Standerton									
	Theunissen				Val									
	Van Tonder													
<b>WHEAT</b>														
Protein (12% mb), %	ave 12.0	min 10.0	max 13.6	stdev 1.08	ave 12.6	min -	max -	stdev -						
Falling number, sec	356	276	408	40.76	421	-	-	-						
1000 Kernel mass (13% mb), g	35.5	29.4	46.3	3.92	39.7	-	-	-						
Hectolitre mass (dirty), kg/hl	80.1	76.6	81.6	1.46	85.0	-	-	-						
Screenings (<1.8mm), %	1.66	0.53	4.04	0.91	0.96	-	-	-						
Total damaged kernels, %	0.45	0.08	1.46	0.40	0.16	-	-	-						
<b>Number of samples</b>	<b>16</b>				<b>1</b>									
<b>CULTIVARS</b>														
cultivars with highest % occurrence	PAN 3120	38.7			SST 843	55.0								
	PAN 3118	16.8			SST 835	45.0								
	Gariep	11.6												
	SST 835	5.8												
	PAN 3355	5.6												
<b>Number of samples</b>	<b>16</b>				<b>1</b>									
<b>MIXOGRAM (Quadromat)</b>														
Peak time, min	ave 3.4	min 2.8	max 4.2	stdev 0.48	ave 2.8	min -	max -	stdev -						
Tail height (6min), mm	52	49	56	2.19	48	-	-	-						
<b>Number of samples</b>	<b>16</b>				<b>1</b>									
<b>BÜHLER EXTRACTION, %</b>	<b>B1</b> 73.2	<b>B2</b> 73.9	<b>B3</b> 73.4	<b>B4</b> -	<b>UT</b> 70.8	<b>COW</b> 72.4	<b>B1</b> 76.8	<b>B2</b> -						
<b>FLOUR</b>														
Protein (12% mb), %	11.6	10.2	9.1	-	12.4	11.7	11.6	-						
Colour, KJ	-2.1	-2.3	-2.9	-	-1.7	-1.3	-2.3	-						
<b>GLUTEN</b>														
Wet gluten (14% mb), %	31.8	26.5	22.5	-	34.1	30.3	32.3	-						
Dry gluten (14% mb), %	11.8	9.6	8.2	-	12.2	11.2	11.4	-						
<b>FARINOGRAM</b>														
Water absorption (14% mb), %	64.9	62.1	60.3	-	67.8	64.8	65.4	-						
Development time, min	6.4	6.2	7.5	-	7.8	8.3	6.7	-						
Stability, min	10.0	9.1	12.5	-	10.7	9.7	6.6	-						
Mixing tolerance index, BU	24	34	23	-	31	39	43	-						
<b>EXTENSOGRAM (45 min pull)</b>														
Area, cm <sup>2</sup>	100	112	103	-	-	131	101	-						
Maximum height, BU	375	430	470	-	-	445	365	-						
Extensibility, mm	195	179	160	-	-	204	199	-						
<b>ALVEOGRAM</b>														
Strength (S), cm <sup>2</sup>	43.9	35.0	35.8	-	48.8	47.9	41.9	-						
Stability (P), mm	109	102	110	-	120	110	101	-						
Distensibility (L), mm	82	65	59	-	82	85	89	-						
Configuration ratio (P/L)	1.33	1.57	1.86	-	1.46	1.29	1.13	-						
<b>MIXOGRAM</b>														
Peak time, min	2.8	3.2	3.6	-	2.5	2.8	2.7	-						
<b>100g BAKING TEST</b>														
Loaf volume, cm <sup>3</sup>	930	855	665	-	825	825	920	-						
Evaluation	0	0	3	-	4	2	0	-						

# RHEOLOGICAL GRAPHS PER PRODUCTION REGION

## MIXOGRAM

24

29

## FARINOGRAM

24

29

## EXTENSOGRAM

24

29

## ALVEOGRAM

24

29

**SOUTH AFRICAN**  
**OTHER SUMMER RAINFALL WHEAT AND IRRIGATION**  
**Mpumalanga**

PRODUCTION REGION	(30) Mpumalanga Eastern Region	(32) Mpumalanga Western Region
Intake silos	Amersfoort Badplaas Carolina Davel Ermelo Estancia Lothair Maizefield Mkondo Morgenzon Overvaal Panbuilt	Argent Dryden Endicott Elof Hawerklip Kendal Ogies
<b>WHEAT</b>		
Protein (12% mb), %	ave 13.0 min - max - stdev -	ave 12.2 min - max - stdev -
Falling number, sec	407	396
1000 Kernel mass (13% mb), g	42.7	41.1
Hectolitre mass (dirty), kg/hl	76.9	82.6
Screenings (<1.8mm), %	0.32	1.31
Total damaged kernels, %	0.32	0.18
<i>Number of samples</i>	1	1
<b>CULTIVARS</b>	Kariega	100.0
cultivars with highest % occurrence		SST 835 90.0
		SST 843 10.0
<i>Number of samples</i>	1	1
<b>MIXOGRAM (Quadromat)</b>		
Peak time, min	ave 4.2 min - max - stdev -	ave 3.5 min - max - stdev -
Tail height (6min), mm	45	51
<i>Number of samples</i>	1	1
<b>BÜHLER EXTRACTION, %</b>	B1 - B2 - B3 - B4 - UT - COW 77.0	B1 76.8 B2 - B3 - B4 - UT - COW -
<b>FLOUR</b>		
Protein (12% mb), %	-	11.9
Colour, KJ	-	11.2
<b>GLUTEN</b>		
Wet gluten (14% mb), %	-	25.1
Dry gluten (14% mb), %	-	29.8
<b>FARINOGRAM</b>		
Water absorption (14% mb), %	-	62.4
Development time, min	-	62.5
Stability, min	-	7.7
Mixing tolerance index, BU	-	8.4
	-	9.8
<b>EXTENSOGRAM (45 min pull)</b>		
Area, cm <sup>2</sup>	-	124
Maximum height, BU	-	460
Extensibility, mm	-	200
<b>ALVEOGRAM</b>		
Strength (S), cm <sup>2</sup>	-	30.6
Stability (P), mm	-	45.6
Distensibility (L), mm	-	139
Configuration ratio (P/L)	-	97
	-	33
<b>MIXOGRAM</b>		
Peak time, min	-	102
<b>100g BAKING TEST</b>		
Loaf volume, cm <sup>3</sup>	-	4.21
Evaluation	-	0.95
	-	6
	-	0

# RHEOLOGICAL GRAPHS PER PRODUCTION REGION

## MIXOGRAM

30

32

## FARINOGRAM

30

32

## EXTENSOGRAM

30

32

## ALVEOGRAM

30

32

**SOUTH AFRICAN**  
**OTHER SUMMER RAINFALL WHEAT AND IRRIGATION**  
**Gauteng and Limpopo Provinces**

PRODUCTION REGION	(34) Gauteng				(35) Limpopo							
	Intake silos	Bloekomspruit	Bronhorstspruit	Glenroy	Alma	Bela-Bela (Warmbad)	Crecy	Immerpan				
		Goeie Hoek	Kaalfontein	Middelvlei	Lehau	Modimolle (Nylstroom)	Mokopane (Potgietersrus)	Naboomspruit				
		Nigel	Oberholzer	Raathsvlei		Northam	Nutfield	Pienaarsrivier				
						Polokwane (Pietersburg)	Roedtan	Settlers				
						Tzaneen	Vaalwater					
<b>WHEAT</b>												
Protein (12% mb), %	ave 11.5	min 10.6	max 13.0	stdev 0.93	ave 11.9	min 10.1	max 13.1	stdev 1.05				
Falling number, sec	436	401	474	23.05	415	371	469	33.54				
1000 Kernel mass (13% mb), g	41.2	36.3	45.2	30.4	41.7	36.8	46.5	3.99				
Hectolitre mass (dirty), kg/hl	82.3	80.9	83.3	0.70	80.6	77.7	82.5	1.95				
Screenings (<1.8mm), %	2.41	1.72	3.22	0.53	1.56	0.38	2.82	0.87				
Total damaged kernels, %	0.32	0.12	0.80	0.18	0.51	0.12	0.90	0.25				
<b>Number of samples</b>	<b>11</b>				<b>8</b>							
<b>CULTIVARS</b>												
cultivars	SST 835	34.4			Duzi	32.5						
with highest % occurrence	Duzi	28.6			SST 835	28.1						
	SST 876	12.6			SST 843	21.1						
	SST 843	6.7			Kariega	11.0						
	CRN 826	6.4			SST 876	7.3						
<b>Number of samples</b>	<b>11</b>				<b>8</b>							
<b>MIXOGRAM (Quadromat)</b>												
Peak time, min	ave 2.2	min 1.8	max 2.3	stdev 0.16	ave 2.9	min 2.4	max 4.3	stdev 0.60				
Tail height (6min), mm	46	41	51	2.73	50	46	59	3.92				
<b>Number of samples</b>	<b>11</b>				<b>8</b>							
<b>BÜHLER EXTRACTION, %</b>	B1 74.6	B2 76.0	B3 75.3	B4 75.3	UT -	COW -	B1 75.3	B2 75.1	B3 77.7	B4 -	UT -	COW -
<b>FLOUR</b>												
Protein (12% mb), %	12.2	10.1	9.8	9.5	-	-	11.8	10.2	9.2	-	-	-
Colour, KJ	-1.8	-2.1	-2.5	-2.5	-	-	-2.3	-2.0	-2.4	-	-	-
<b>GLUTEN</b>												
Wet gluten (14% mb), %	34.7	27.9	27.4	26.5	-	-	31.4	28.6	25.0	-	-	-
Dry gluten (14% mb), %	11.8	9.2	8.9	8.9	-	-	10.8	9.9	8.5	-	-	-
<b>FARINOGRAM</b>												
Water absorption (14% mb), %	65.8	63.4	61.5	63.0	-	-	64.1	62.8	61.4	-	-	-
Development time, min	5.2	3.3	4.0	4.0	-	-	6.7	4.7	4.8	-	-	-
Stability, min	6.5	5.0	4.6	4.7	-	-	8.1	5.5	5.9	-	-	-
Mixing tolerance index, BU	29	43	48	47	-	-	35	44	48	-	-	-
<b>EXTENSOGRAM (45 min pull)</b>												
Area, cm <sup>2</sup>	74	57	60	56	-	-	115	-	59	-	-	-
Maximum height, BU	290	240	250	235	-	-	425	-	265	-	-	-
Extensibility, mm	176	158	159	153	-	-	192	-	149	-	-	-
<b>ALVEOGRAM</b>												
Strength (S), cm <sup>2</sup>	34.6	26.6	23.1	20.9	-	-	43.4	27.4	24.8	-	-	-
Stability (P), mm	92	84	72	81	-	-	96	75	76	-	-	-
Distensibility (L), mm	89	75	78	57	-	-	97	91	74	-	-	-
Configuration ratio (P/L)	1.03	1.12	0.92	1.42	-	-	0.99	0.82	1.03	-	-	-
<b>MIXOGRAM</b>												
Peak time, min	2.3	2.2	2.3	2.0	-	-	2.8	2.3	2.3	-	-	-
<b>100g BAKING TEST</b>												
Loaf volume, cm <sup>3</sup>	880	815	800	780	-	-	965	915	820	-	-	-
Evaluation	2	0	0	0	-	-	0	0	0	-	-	-

# RHEOLOGICAL GRAPHS PER PRODUCTION REGION

## MIXOGRAM

34

35

## FARINOGRAM

34

35

## EXTENSOGRAM

34

35

## ALVEOGRAM

34

35

**SOUTH AFRICAN  
OTHER SUMMER RAINFALL WHEAT AND IRRIGATION  
KwaZulu-Natal Province**

PRODUCTION REGION	(36) KwaZulu-Natal				
Intake silos	Bergville Bloedrivier Dannhauser Dundee Mizpah New Amalfi Paulpietersburg Vryheid Winterton				
<b>WHEAT</b>					
Protein (12% mb), %	<b>ave</b>	<b>min</b>	<b>max</b>	<b>stdev</b>	
	12.7	11.1	13.9	0.93	
Falling number, sec	432	373	475	37.31	
1000 Kernel mass (13% mb), g	39.1	33.0	42.7	3.11	
Hectolitre mass (dirty), kg/hl	82.2	80.6	83.9	1.02	
Screenings (<1.8mm), %	0.64	0.49	0.81	0.11	
Total damaged kernels, %	0.16	0.00	0.76	0.25	
<b>Number of samples</b>	<b>8</b>				
<b>CULTIVARS</b>					
cultivars	SST 835                    53.6				
with highest % occurrence	SST 843                    42.0				
	SST 877                    2.3				
	Kariega                    2.1				
<b>Number of samples</b>	<b>8</b>				
<b>MIXOGRAM (Quadromat)</b>					
Peak time, min	<b>ave</b>	<b>min</b>	<b>max</b>	<b>stdev</b>	
	2.9	2.0	3.3	0.42	
Tail height (6min), mm	49	44	52	3.04	
<b>Number of samples</b>	<b>8</b>				
<b>BÜHLER EXTRACTION, %</b>	<b>B1</b>	<b>B2</b>	<b>B3</b>	<b>B4</b>	<b>UT</b>
76.4	76.4	-	-	-	-
<b>FLOUR</b>					
Protein (12% mb), %	12.0	10.7	-	-	-
Colour, KJ	-2.0	-1.9	-	-	-
<b>GLUTEN</b>					
Wet gluten (14% mb), %	33.3	30.0	-	-	-
Dry gluten (14% mb), %	11.2	10.4	-	-	-
<b>FARINOGRAM</b>					
Water absorption (14% mb), %	64.8	62.2	-	-	-
Development time, min	6.3	5.8	-	-	-
Stability, min	6.7	6.6	-	-	-
Mixing tolerance index, BU	37	41	-	-	-
<b>EXTENSOGRAM (45 min pull)</b>					
Area, cm <sup>2</sup>	102	95	-	-	-
Maximum height, BU	375	350	-	-	-
Extensibility, mm	191	182	-	-	-
<b>ALVEOGRAM</b>					
Strength (S), cm <sup>2</sup>	40.2	30.7	-	-	-
Stability (P), mm	97	86	-	-	-
Distensibility (L), mm	91	80	-	-	-
Configuration ratio (P/L)	1.07	1.08	-	-	-
<b>MIXOGRAM</b>					
Peak time, min	2.7	2.8	-	-	-
<b>100g BAKING TEST</b>					
Loaf volume, cm <sup>3</sup>	950	885	-	-	-
Evaluation	0	0	-	-	-

# RHEOLOGICAL GRAPHS PER PRODUCTION REGION

## MIXOGRAM

36

## FARINOGRAM

36

## EXTENSOGRAM

36

## ALVEOGRAM

36

# RSA WHEAT CROP QUALITY SUMMARY

RSA Crop Quality 2008/2009 and 2010/2011 Seasons

Country of origin		RSA Crop Average 2008/2009							RSA Crop Average 2010/2011						
Class and Grade bread wheat		B1	B2	B3	B4	UT	COW	Average	B1	B2	B3	B4	UT	COW	Average
No. of samples		126	121	101	49	16	67	480	164	93	48	23	16	28	372
<b>WHEAT GRADING</b>															
Protein (12% mb), %		13.27	12.00	11.25	10.69	12.69	11.54	12.00	12.91	11.53	10.69	11.20	13.31	12.28	12.14
Moisture, %		11.5	11.1	11.0	10.9	11.6	11.0	11.2	11.8	11.7	11.6	11.6	12.2	11.8	11.8
Falling number, sec		364	395	382	346	364	393	378	385	389	371	359	180	358	372
1000 Kernel mass (13% mb), g		37.5	38.5	39.1	40.6	36.6	37.4	38.3	39.1	40.0	39.9	40.8	39.3	38.5	39.5
Hl/m (dirty), kg/hl		78.6	77.8	77.2	77.4	76.0	76.4	77.6	80.9	80.4	80.6	80.8	76.6	78.1	80.3
Screenings (<1,8mm), %		1.36	1.52	1.74	1.33	2.76	2.77	1.72	1.34	1.61	1.51	2.66	1.49	3.48	1.68
Gravel, stones, turf and glass, %		0.00	0.00	0.00	0.01	0.13	0.00	0.01	0.00	0.00	0.00	0.00	0.07	0.00	0.01
Foreign matter, %		0.09	0.09	0.08	0.10	0.29	0.17	0.11	0.07	0.08	0.07	0.10	0.42	0.18	0.10
Other grain & unthreshed ears, %		0.20	0.24	0.29	0.26	0.32	0.34	0.26	0.28	0.34	0.33	0.43	0.26	0.85	0.35
Heat damaged kernels, %		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.02	0.00	0.00
Immature kernels, %		0.12	0.07	0.04	0.05	0.04	0.06	0.07	0.06	0.02	0.02	0.02	0.10	0.06	0.04
Insect damaged kernels, %		0.27	0.42	0.46	0.54	4.02	1.13	0.62	0.24	0.28	0.29	0.27	0.28	0.35	0.27
Heavily frost damaged kernels, %		0.00	0.00	0.00	0.02	0.00	0.01	0.00	0.02	0.00	0.00	0.00	0.00	0.00	0.01
Sprouted kernels, %		0.10	0.09	0.11	0.10	1.01	0.14	0.13	0.04	0.03	0.03	0.06	1.43	0.15	0.11
Total damaged kernels, %		0.49	0.58	0.62	0.75	5.07	1.40	0.85	0.34	0.33	0.35	0.35	1.83	0.56	0.42
Combined deviations, %		2.14	2.42	2.71	2.27	8.44	4.49	2.88	2.03	2.36	2.26	3.55	3.99	5.07	2.55
Field fungi, %		0.08	0.12	0.09	0.04	0.23	0.12	0.10	0.18	0.12	0.10	0.14	1.12	0.29	0.20
Storage fungi, %		0.01	0.01	0.01	0.00	0.04	0.01	0.01	0.02	0.01	0.01	0.02	0.14	0.03	0.02
Ergot, %		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Noxious seeds ( <i>Crotalaria spp.</i> , <i>Datura spp.</i> )		0	0	0	0	0	0	0	0	0	0	0	0	0	0
Noxious seeds ( <i>Argemone mexicana</i> )		0	0	0	0	0	0	0	0	0	0	0	0	0	0
Live insects		No	No	No	No	No	No	No	No	No	No	No	No	No	No
Undesirable odour		No	No	No	No	No	No	No	No	No	No	No	No	No	No
		B1	B2	B3	B4	UT	COW	Average	B1	B2	B3	B4	UT	COW	Average
<b>No. of samples</b>		22	23	22	12	5	16	100	29	24	17	11	8	10	99
<b>BÜHLER EXTRACTION, %</b>		76.0	76.1	75.7	75.2	75.1	75.2	75.7	75.1	75.1	75.0	74.7	74.2	74.4	74.9
<b>FLOUR</b>															
Colour, KJ		-1.3	-1.7	-1.7	-1.6	-0.9	-1.5	-1.5	-2.0	-2.1	-2.0	-1.9	0.4	-1.5	-1.8
Protein (12% mb), %		12.1	11.1	10.6	10.5	11.3	11.0	11.1	12.0	10.5	9.7	9.8	12.1	11.5	11.0
Wet Gluten (14% mb), %		-	-	-	-	-	-	-	32.7	28.4	25.7	28.0	31.1	31.4	29.7
Dry Gluten (14% mb), %		-	-	-	-	-	-	-	11.6	9.9	9.0	9.6	11.1	11.1	10.4
<b>100g BAKING TEST</b>															
Baking water absorption, %		61.9	60.8	60.0	60.0	60.1	60.3	60.7	62.3	60.5	59.6	59.9	62.4	61.8	61.1
Loaf volume, cm <sup>3</sup>		956	909	870	849	926	892	902	882	824	766	789	853	846	832
Evaluation		1	0	0	1	0	0	0	2	1	1	1	2	2	1
<b>FARINOGRAM</b>															
Water absorption, %		62.2	61.1	60.4	60.9	60.6	60.7	61.1	64.2	62.7	61.7	62.3	64.0	63.8	63.2
Development time, min		5.0	4.1	3.6	3.1	3.7	3.8	4.0	7.0	5.3	5.0	4.0	4.7	5.2	5.5
Stability, mm		8.6	7.7	7.3	6.8	7.4	7.4	7.6	9.1	7.3	7.5	6.0	7.6	8.1	7.8
Mixing tolerance index, BU		42	42	44	46	43	45	43	33	39	38	43	40	37	37

## RSA Crop Quality of 2008/2009 and 2010/2011 Seasons

Country of origin	RSA Crop Average 2008/2009							RSA Crop Average 2010/2011						
Class and Grade bread wheat	B1	B2	B3	B4	UT	COW	Average	B1	B2	B3	B4	UT	COW	Average
No. of samples	22	23	22	12	5	16	100	164	93	48	23	16	28	372
<b>ALVEOGRAM</b>														
Strength (S) , cm <sup>2</sup>	42.4	38.0	35.6	35.8	36.6	37.2	38.0	43.1	32.6	30.5	29.2	39.4	39.5	36.2
Stability (P), mm	85	83	82	87	78	82	83	98	91	92	91	102	98	95
Distensibility (L), mm	111	103	95	88	114	101	101	94	78	69	70	78	87	81
P/L	0.80	0.86	0.93	1.10	0.91	0.94	0.90	1.08	1.21	1.44	1.55	1.57	1.28	1.29
<b>EXTENSOGRAM</b>														
Strength, cm <sup>2</sup>	101	89	83	85	89	90	90	110	93	85	80	109	101	97
Max. height, BU	348	330	330	337	325	340	336	381	357	358	323	378	371	362
Extensibility, mm	201	184	172	174	187	179	183	204	181	166	170	203	193	187
<b>MIXOGRAM</b>														
Peak time, min	2.5	2.6	2.7	2.7	2.8	2.7	2.6	2.8	2.6	2.8	2.6	3.2	2.8	2.8
Absorption, %	62.3	61.0	60.6	60.6	61.4	61.1	61.2	62.2	60.4	59.6	59.8	62.3	61.7	61.0
<b>MYCOTOXINS</b>														
Aflatoxin, ppb [max.value]	1.23 [3.00]							0 [0]						
Deoxynivalenol, ppm [max. value]	0.47 [3.00]							0 [0]						
Ochratoxin A, ppb [max. value]	0.03 [1.00]							0 [0]						
<b>No. of samples</b>	<b>30</b>							<b>30</b>						

# RSA WHEAT CROP QUALITY SUMMARY

RSA Crop Quality 2009/2010 and 2010/2011 Seasons

Country of origin		RSA Crop Average 2009/2010							RSA Crop Average 2010/2011						
Class and Grade bread wheat		B1	B2	B3	B4	UT	COW	Average	B1	B2	B3	B4	UT	COW	Average
No. of samples		159	138	79	30	16	58	480	164	93	48	23	16	28	372
<b>WHEAT</b>															
<b>GRADING</b>															
Protein (12% mb), %		12.84	11.50	10.62	9.64	11.29	11.51	11.68	12.91	11.53	10.69	11.20	13.31	12.28	12.14
Moisture, %		11.5	11.3	11.3	10.8	11.0	11.0	11.3	11.8	11.7	11.6	11.6	12.2	11.8	11.8
Falling number, sec		382	375	359	339	240	365	367	385	389	371	359	180	358	372
1000 Kernel mass (13% mb), g		38.7	39.5	40.4	40.2	36.6	38.1	39.2	39.1	40.0	39.9	40.8	39.3	38.5	39.5
Hl <sub>m</sub> (dirty), kg/hl		80.8	80.0	79.5	79.7	75.2	79.0	79.9	80.9	80.4	80.6	80.8	76.6	78.1	80.3
Screenings (<1.8mm), %		1.36	1.30	1.47	1.44	3.36	3.03	1.63	1.34	1.61	1.51	2.66	1.49	3.48	1.68
Gravel, stones, turf and glass, %		0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.07	0.00	0.01
Foreign matter, %		0.05	0.05	0.09	0.09	0.12	0.11	0.07	0.07	0.08	0.07	0.10	0.42	0.18	0.10
Other grain & unthreshed ears, %		0.25	0.26	0.30	0.35	0.64	0.65	0.33	0.28	0.34	0.33	0.43	0.26	0.85	0.35
Heat damaged kernels, %		0.00	0.00	0.00	0.00	0.05	0.00	0.00	0.00	0.00	0.00	0.00	0.02	0.00	0.00
Immature kernels, %		0.07	0.05	0.04	0.03	0.01	0.06	0.05	0.06	0.02	0.02	0.02	0.10	0.06	0.04
Insect damaged kernels, %		0.19	0.26	0.29	0.27	1.10	0.52	0.30	0.24	0.28	0.29	0.27	0.28	0.35	0.27
Heavily frost damaged kernels, %		0.00	0.00	0.00	0.00	0.13	0.00	0.01	0.02	0.00	0.00	0.00	0.00	0.00	0.01
Sprouted kernels, %		0.09	0.11	0.16	0.14	2.73	0.36	0.23	0.04	0.03	0.03	0.06	1.43	0.15	0.11
Total damaged kernels, %		0.36	0.43	0.50	0.44	3.96	0.96	0.60	0.34	0.33	0.35	0.35	1.83	0.56	0.42
Combined deviations, %		2.02	2.04	2.34	2.32	8.07	4.76	2.63	2.03	2.36	2.26	3.55	3.99	5.07	2.55
Field fungi, %		0.19	0.19	0.21	0.15	0.54	0.16	0.20	0.18	0.12	0.10	0.14	1.12	0.29	0.20
Storage fungi, %		0.01	0.01	0.01	0.00	0.03	0.01	0.01	0.02	0.01	0.01	0.02	0.14	0.03	0.02
Ergot, %		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Noxious seeds ( <i>Crotalaria spp.</i> , <i>Datura spp.</i> )		0	0	0	0	0	0	0	0	0	0	0	0	0	0
Noxious seeds ( <i>Argemone mexicana</i> )		0	0	0	0	0	0	0	0	0	0	0	0	0	0
Live insects		No	No	No	No	No	No	No	No	No	No	No	No	No	No
Undesirable odour		No	No	No	No	No	No	No	No	No	No	No	No	No	No
		B1	B2	B3	B4	UT	COW	Average	B1	B2	B3	B4	UT	COW	Average
<b>No. of samples</b>		26	28	20	10	5	11	100	29	24	17	11	8	10	99
<b>BÜHLER EXTRACTION, %</b>		74.8	75.1	74.8	73.8	74.1	74.9	74.7	75.1	75.1	75.0	74.7	74.2	74.4	74.9
<b>FLOUR</b>															
Colour, KJ		-2.2	-2.3	-2.3	-2.3	-1.7	-2.2	-2.2	-2.0	-2.1	-2.0	-1.9	0.4	-1.5	-1.8
Protein (12% mb), %		11.7	10.6	9.6	8.6	10.4	10.6	10.5	12.0	10.5	9.7	9.8	12.1	11.5	11.0
Wet Gluten (14% mb), %		32.1	29.0	25.9	22.2	28.3	29.1	28.6	32.7	28.4	25.7	28.0	31.1	31.4	29.7
Dry Gluten (14% mb), %		11.3	10.1	9.0	7.9	9.6	10.2	10.0	11.6	9.9	9.0	9.6	11.1	11.1	10.4
<b>100g BAKING TEST</b>															
Baking water absorption, %		61.5	60.3	59.3	58.2	59.5	60.4	60.2	62.3	60.5	59.6	59.9	62.4	61.8	61.1
Loaf volume, cm <sup>3</sup>		902	853	803	727	868	847	843	882	824	766	789	853	846	832
Evaluation		1	0	0	0	0	0	0	2	1	1	1	2	2	1
<b>FARINOGRAM</b>															
Water absorption, %		62.3	61.3	60.5	59.3	58.3	60.6	61.0	64.2	62.7	61.7	62.3	64.0	63.8	63.2
Development time, min		4.8	3.4	2.9	2.0	2.7	3.7	3.5	7.0	5.3	5.0	4.0	4.7	5.2	5.5
Stability, mm		8.4	6.8	5.7	3.8	4.9	6.9	6.6	9.1	7.3	7.5	6.0	7.6	8.1	7.8
Mixing tolerance index, BU		39	44	52	71	71	48	49	33	39	38	43	40	37	37

## RSA Crop Quality of 2009/2010 and 2010/2011 Seasons

Country of origin	RSA Crop Average 2009/2010							RSA Crop Average 2010/2011						
Class and Grade bread wheat	B1	B2	B3	B4	UT	COW	Average	B1	B2	B3	B4	UT	COW	Average
No. of samples	26	28	20	10	5	11	100	164	93	48	23	16	28	372
<b>ALVEOGRAM</b>														
Strength (S) , cm <sup>2</sup>	42.4	36.5	32.7	25.0	26.8	35.4	35.5	43.1	32.6	30.5	29.2	39.4	39.5	36.2
Stability (P), mm	93	90	90	87	61	83	89	98	91	92	91	102	98	95
Distensibility (L), mm	97	86	75	58	101	90	85	94	78	69	70	78	87	81
P/L	1.03	1.13	1.31	1.80	0.61	0.97	1.17	1.08	1.21	1.44	1.55	1.57	1.28	1.29
<b>EXTENSOGRAM</b>														
Strength, cm <sup>2</sup>	96	85	74	61	71	87	83	110	93	85	80	109	101	97
Max. height, BU	355	344	327	304	295	345	337	381	357	358	323	378	371	362
Extensibility, mm	187	174	158	139	171	173	170	204	181	166	170	203	193	187
<b>MIXOGRAM</b>														
Peak time, min	2.6	2.6	2.7	2.8	2.8	2.5	2.6	2.8	2.6	2.8	2.6	3.2	2.8	2.8
Absorption, %	61.8	60.4	59.4	58.5	60.3	60.6	60.4	62.2	60.4	59.6	59.8	62.3	61.7	61.0
<b>MYCOTOXINS</b>														
Aflatoxin, ppb [max.value]	1.00 [4.00]							0 [0]						
Deoxynivalenol, ppm [max. value]	0.05 [0.48]							0 [0]						
Ochratoxin A, ppb [max. value]	0.17 [1.00]							0 [0]						
No. of samples	30							30						

# METHODS

## GRADING:

Full grading was done in accordance with the Regulations relating to the grading, packing and marking of wheat intended for sale in the Republic of South Africa (No. R. 1186 of 17 December 2010).

Hectolitre mass, screenings, protein and falling number were determined. The determination of deviations relating to wheat kernels comprised foreign matter including gravel, stones, turf and glass; other grain and unthreshed ears; damaged kernels including heat-damaged kernels, immature kernels, insect-damaged kernels and sprouted kernels; heavily frost-damaged kernels; field fungi; storage fungi; ergot; noxious seeds; possible presence of undesirable odours and live insects.

Hectolitre mass means the mass in kilogram per hectolitre and was determined according to ISO 7971-3 by means of the Kern 222 instrument.

During previous seasons the hectolitre mass was determined by means of the Two-level funnel method. In the 2009/2010 season the hectolitre mass value was adjusted by the addition of  $2 \text{ kg}/\text{hl}$  to all hectolitre mass values as per an Industry-wide Hectolitre Mass Dispensation published by the National Department of Agriculture.

Hectolitre mass provides a measure of the bulk density of grain and is also useful as a guide to grain soundness and potential milling extraction.

Screenings means all material that passes through a standard sieve. For the definition of a standard sieve please refer to the definitions of Regulation No. R. 1186 on pages 78 - 79 of this report.

Damaged wheat means wheat -

- (a) which have been damaged by insects;
- (b) which have been distinctly discoloured (orange-brown, dark brown or black) by external heat or as a result of heating caused by internal fermentation in wheat with an excessive moisture content, excluding wheat kernels in respect of which the discolouration is confined to the germ end;
- (c) which are immature and have a distinctly green colour; and
- (d) in which germination has proceeded to such an extent that the skin covering the embryo has been broken or the developing sprouts and/or rootlets are clearly visible.

## THOUSAND KERNEL MASS:

This is the weight in grams of one thousand kernels of grain and provides a measure of grain size and density. This determination does not include kernels that are broken or chipped.

## FALLING NUMBER MILLING:

At least 300 g of wheat is cleaned by using the standard 1,8 mm sieve and by removing coarser impurities by hand. The sample is then milled on the falling number hammer mill fitted with a 0,8 mm screen.

## MOISTURE:

ICC Standard No. 110/1, latest edition is used to determine the moisture content of wheat flour. This method determines moisture content as a loss in weight of a sample when dried in an oven at 130°C for 90 minutes or 2 hours for flour and whole wheat flour respectively.

## PROTEIN:

The Dumas combustion analysis technique is used, according to AACC method 46-30.01, latest edition.

This method prescribes a generic combustion method for the determination of crude protein. Combustion at high temperature in pure oxygen sets nitrogen free, which is measured by thermal conductivity detection. The total nitrogen content of the flour sample is determined and converted to equivalent protein by multiplication with a factor of 5.7 to obtain the protein content.

## FALLING NUMBER:

This method is based upon the rapid gelatinization of an aqueous suspension of meal or flour in a boiling water bath and subsequent measurement of the liquefaction of the starch paste by the alpha-amylase in the sample. The method measures the alpha-amylase activity.

ICC Standard No.107/1, latest edition is used to determine the falling number. Only the altitude-corrected value is reported.

## QUADROMAT MILLING:

Cleaned wheat samples are conditioned by adding 3 ml water per 100 g wheat, 18 hours prior to milling.

The samples are then milled on the Quadromat junior laboratory mill.

#### MIXOGRAPH:

A 35 g mixograph is used. The amount of flour weighed is adjusted according to the flour moisture content and the amount of water added to the flour is adjusted according to the flour protein content. Industry Accepted Method 020 based on AACC method 54-40.02, latest edition is followed.

Mixogram peak time is the time measured in minutes that a dough takes to reach its maximum consistency or first indication of dough weakening. The peak time is a measure of optimum dough development and thus a measure of protein quality.

Mixogram tail height at 6 minutes is the distance in millimetres measured from the base line of the paper at 6 minutes to the graph centre point at 6 minutes. This figure is an indication of the weakening effect of the dough. Higher values indicate flours that are more tolerant to mixing.

#### BÜHLER MILLING:

Cleaned wheat samples are damped to between 15,0 % and 16,0 % moisture according to the wheat moisture and kernel hardness and allowed to stand for 20 hours. Samples are then milled on a standard Bühler MLU 202 mill and passed through a bran finisher.

#### BÜHLER EXTRACTION:

The extraction represents the flour yield after milling plus flour obtained from bran that passed through a bran finisher. Flour extraction is calculated from the mass of the total products. Bühler MLU 202 mill set for South African wheat, mill settings and sieve sizes deviate from AACC method 26-21.02, latest edition.

#### COLOUR:

The Kent Jones colour is determined by following FTP Method No. 0007/3, 7/1991. This method determines the influence of the branny material present in flour by measuring reflectance with a light source in the green band of the light spectrum. The lower the Kent Jones colour, the lighter the flour.

#### GLUTEN:

Wheat gluten is the water-insoluble complex protein fraction separated from wheat flours. The ability of wheat flour to produce dough with good gas retaining properties is attributed to gluten.

The gluten content of wheat flour is determined by means of AACC Method 38-12.02, latest edition. Wet gluten is washed from meal or flour by an automatic washing apparatus (Glutomatic). Wet gluten is a plastic elastic substance composed principally of two protein fractions. Glutenin, the higher molecular weight fraction, contributes elasticity and Gliadin, the lower molecular weight fraction, provides extensibility.

The wet gluten is dried under standardized conditions in a Glutork to obtain the dry gluten. The total wet and total dry gluten contents are expressed as percentages of the sample on a 14% moisture basis.

#### FARINOGRAPH:

AACC method 54-21.01, latest edition constant flour weight procedure is followed, using 300 g of flour on a 14 % moisture basis.

The farinograph measures and records the resistance of a dough to mixing, as it is formed from flour and water, developed and broken down. The dough is subjected to a prolonged, relatively gentle mixing action at a constant temperature.

The water absorption is the amount of water required for a dough to reach a definite consistency (500 Brabender units). The amount of water added to the flour is expressed as a percentage of the flour mass and reported on a 14 % moisture basis.

The development time is the time from the beginning of water addition until the dough reaches its optimum consistency and the point immediately before the first indication of weakening. A long mixing time can be associated with flours with a high percentage of gluten-forming proteins.

The stability is the time during which the top of the curve intercepts a horizontal line through the centre of the curve. This gives an indication of the dough's tolerance to mixing: the longer the stability, the longer the mixing time that the dough can withstand. A dough with a longer stability can also withstand a longer fermentation period. The mixing tolerance index value is the difference,

in Brabender units, between the top of the curve at the peak and the top of the curve measured 5 minutes after the peak is reached. The value gives an indication of the extent to which breakdown of the dough occurs. The higher the value, the more and the quicker the breakdown of the dough occurs. This value is similar to the mixogram tail height.

#### EXTENSOGRAPH:

ICC Standard No. 114/1, latest edition is followed.

The strength gives an indication of the total force (work) needed to stretch the dough and is represented by the area under the curve.

The maximum height gives an indication of the dough's resistance to stretching and is measured as the mean of the maximum heights of the curves of the two test pieces.

The extensibility is the mean length at the base of the 2 curves and indicates the stretchability of the dough.

#### ALVEOGRAPH:

ICC Standard No.121, latest edition is followed.

The alveograph measures the resistance of the dough to stretching and also how extensible the dough is. The alveograph stretches the dough in more than one direction (as is happening during proofing), whereas the extensograph stretches the dough in only one direction.

Strength (S): The area under the curve gives an indication of the dough strength.

Stability (P): Obtained by multiplying the maximum height of the curve with a constant factor of 1.1. This value is an indication of the resistance of the dough to extension.

Distensibility (L): The length of the curve, measured along the base line, gives an indication of the extensibility of the dough and also predicts the handling characteristics of the dough.

P/L-value: This ratio is obtained by dividing the P-value by the L-value, thus providing an approximate indication of the shape of the curve that combines stability and extensibility.

#### 100 g BAKING TEST:

This procedure, according to Industry Accepted Method 022 based on AACC Method 10-10.03, latest edition, provides an optimized bread-making method for evaluating bread wheat flour quality and a variety of dough ingredients by a straight-dough method in which all ingredients are incorporated in the initial mixing step.

Keys for the evaluation of the 100g Baking test:

- 0 - Excellent
- 1 - Very Good
- 2 - Good
- 3 - Questionable
- 4 - Poor
- 5 - Very Poor
- 6 - Extremely Poor

Please note:

This 100 g Baking test evaluation does not give an indication of the baking quality of the flour, but refers to the relationship between the protein content and the bread volume.

#### MYCOTOXIN ANALYSES

Mycotoxins, produced by moulds or fungi, are natural contaminants of food and feedstuffs with serious implications for public health and economics, in particular with relation to the international food trade.

During 2010 SAGL implemented a multi-mycotoxin screening method using UPLC - MS/MS. 30 of the 372 wheat crop samples were tested for Aflatoxin G<sub>1</sub>; B<sub>1</sub>; G<sub>2</sub>; B<sub>2</sub>, Fumonisin B<sub>1</sub> and B<sub>2</sub>, Deoxynivalenol, T2 - Toxin, Zearalenone and Ochratoxin A.

## Quantity of imported wheat for the 2009/2010 season (Previous season)

During the 2009/2010 season, 1 277 972 tons of wheat were imported for RSA. The biggest quantity was imported from Germany, namely 809 934 tons, followed by USA with 173 030 tons, Brazil with 123 944 tons, Canada with 72 911 tons, Australia with 55 312 tons, Ukraine with 41 230 tons and Lithuania with 1 611 tons (SAGIS).

For grading as well as dough and baking quality of the imported wheat, please refer to pages 63 to 72. No samples were received for analysis from Ukraine and Lithuania.

## Imported wheat (1 October 2009 - 30 September 2010) (Previous season)

The quality of all wheat imported into South Africa is also monitored by the SAGL. The range of analyses done on the local crop for the purpose of this survey are also done on the imported wheat. These results may only be made available at the end of each season.

Pages 63 to 72 of this report contain summaries of wheat imported from specific countries during the 2009/2010 season. This imported wheat quality is compared to a summary of the local crop quality for the same season.

The quality of the Australian and Canadian flour milled from wheat imported during the 2009/2010 season were better than the local wheat flour quality. The German and American flour were not as good as the quality of the local flour of the 2009/2010 season, while the Brazilian flour were noticeably weaker than the local flour.

### Average quality data of imported wheat during the 2009/2010 season (previous season)

	Australia	Brazil	Canada	Germany	USA	RSA
Protein, % (12 % mb)	12.23	12.28	14.09	11.18	11.43	11.68
Hl <sub>m</sub> , kg/hl	79.9	76.9	82.4	78.7	80.7	79.9
Screenings, %	1.87	3.05	2.11	1.56	2.50	1.63
Number of samples	5	10	5	70	10	480
Extraction, %	72.2	69.8	74.4	74.2	73.0	74.7
Flour colour, KJ	-2.6	0.5	-1.9	-1.3	-1.0	-2.2
<i>Farinogram</i>						
Water absorption, %	61.0	64.0	63.3	59.2	56.7	61.0
Development time, %	2.7	1.8	5.4	1.9	2.0	3.5
<i>Alveogram</i>						
Strength, cm <sup>2</sup>	40.6	26.2	52.2	33.2	31.7	35.5
P/L	1.80	1.95	0.69	1.63	1.03	1.17
<i>Extensogram</i>						
Strength, cm <sup>2</sup>	101	50	120	84	98	83
<i>Mixogram (Bühler)</i>						
Peak time, min	3.4	2.7	2.9	3.5	4.6	2.6
<i>100 g Baking test</i>						
Volume, cm <sup>3</sup>	780	723	988	757	782	843
Evaluation	3	4	1	2	1	0
Number of samples	5	10	5	70	10	100

# 2009/2010 IMPORTED WHEAT QUALITY - AUSTRALIA (1 Oct 2009 to 30 Sep 2010)

## 2009/2010 Imported Wheat Quality Versus 2009/2010 RSA Wheat Quality

Country of origin		Australia						RSA Crop Average							
Class and Grade bread wheat		B1	B2	B3	B4	UT	COW	Average	B1	B2	B3	B4	UT	COW	Average
No. of samples		5	-	-	-	-	-	5	159	138	79	30	16	58	480
<b>WHEAT GRADING</b>															
Protein (12% mb), %	12.23	-	-	-	-	-	-	12.23	12.84	11.50	10.62	9.64	11.29	11.51	11.68
Moisture, %	10.0	-	-	-	-	-	-	10.0	11.5	11.3	11.3	10.8	11.0	11.0	11.3
Falling number, sec	510	-	-	-	-	-	-	510	382	375	359	339	240	365	367
1000 Kernel mass (13% mb), g	35.2	-	-	-	-	-	-	35.2	38.7	39.5	40.4	40.2	36.6	38.1	39.2
HlM (dirty), kg/hl	79.9	-	-	-	-	-	-	79.9	80.8	80.0	79.5	79.7	75.2	79.0	79.9
Screenings (<1.8mm), %	1.87	-	-	-	-	-	-	1.87	1.36	1.30	1.47	1.44	3.36	3.03	1.63
Gravel, stones, turf and glass, %	0.00	-	-	-	-	-	-	0.00	0.00	0.00	0.01	0.00	0.00	0.00	0.00
Foreign matter, %	0.01	-	-	-	-	-	-	0.01	0.05	0.05	0.09	0.09	0.12	0.11	0.07
Other grain & unthreshed ears, %	0.14	-	-	-	-	-	-	0.14	0.25	0.26	0.30	0.35	0.64	0.65	0.33
Heat damaged kernels, %	0.00	-	-	-	-	-	-	0.00	0.00	0.00	0.00	0.00	0.05	0.00	0.00
Immature kernels, %	0.00	-	-	-	-	-	-	0.00	0.07	0.05	0.04	0.03	0.01	0.06	0.05
Insect damaged kernels, %	0.00	-	-	-	-	-	-	0.00	0.19	0.26	0.29	0.27	1.10	0.52	0.30
Heavily frost damaged kernels, %	0.00	-	-	-	-	-	-	0.00	0.00	0.00	0.00	0.00	0.13	0.00	0.01
Sprouted kernels, %	0.02	-	-	-	-	-	-	0.02	0.09	0.11	0.16	0.14	2.73	0.36	0.23
Total damaged kernels, %	0.02	-	-	-	-	-	-	0.02	0.36	0.43	0.50	0.44	3.96	0.96	0.60
Combined deviations, %	2.04	-	-	-	-	-	-	2.04	2.02	2.04	2.34	2.32	8.07	4.76	2.63
Field fungi, %	0.23	-	-	-	-	-	-	0.23	0.19	0.19	0.21	0.15	0.54	0.16	0.20
Storage fungi, %	0.00	-	-	-	-	-	-	0.00	0.01	0.01	0.01	0.00	0.03	0.01	0.01
Ergot, %	0.00	-	-	-	-	-	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Noxious seeds ( <i>Crotalaria spp.</i> , <i>Datura spp.</i> )	0	-	-	-	-	-	-	0	0	0	0	0	0	0	0
Noxious seeds ( <i>Argemone mexicana</i> )	0	-	-	-	-	-	-	0	0	0	0	0	0	0	0
Live insects	No	-	-	-	-	-	-	No	No	No	No	No	No	No	No
Undesirable odour	No	-	-	-	-	-	-	No	No	No	No	No	No	No	No
		B1	B2	B3	B4	UT	COW	Average	B1	B2	B3	B4	UT	COW	Average
No. of samples		5	-	-	-	-	-	5	26	28	20	10	5	11	100
BÜHLER EXTRACTION, %		72.2	-	-	-	-	-	72.2	74.8	75.1	74.8	73.8	74.1	74.9	74.7
<b>FLOUR</b>															
Colour, KJ	-2.6	-	-	-	-	-	-	-2.6	-2.2	-2.3	-2.3	-2.3	-1.7	-2.2	-2.2
Protein (12% mb), %	11.2	-	-	-	-	-	-	11.2	11.7	10.6	9.6	8.6	10.4	10.6	10.5
Wet Gluten (14% mb), %	29.6	-	-	-	-	-	-	29.6	32.1	29.0	25.9	22.2	28.3	29.1	28.6
Dry Gluten (14% mb), %	10.3	-	-	-	-	-	-	10.3	11.3	10.1	9.0	7.9	9.6	10.2	10.0
<b>100g BAKING TEST</b>															
Baking water absorption, %	61.1	-	-	-	-	-	-	61.1	61.5	60.3	59.3	58.2	59.5	60.4	60.2
Loaf volume, cm <sup>3</sup>	780	-	-	-	-	-	-	780	902	853	803	727	868	847	843
Evaluation	3	-	-	-	-	-	-	3	1	0	0	0	0	0	0
<b>FARINOGRAM</b>															
Water absorption, %	61.0	-	-	-	-	-	-	61.0	62.3	61.3	60.5	59.3	58.3	60.6	61.0
Development time, min	2.7	-	-	-	-	-	-	2.7	4.8	3.4	2.9	2.0	2.7	3.7	3.5
Stability, mm	8.6	-	-	-	-	-	-	8.6	8.4	6.8	5.7	3.8	4.9	6.9	6.6
Mixing tolerance index, BU	25	-	-	-	-	-	-	25	39	44	52	71	71	48	49

## 2009/2010 Imported Wheat Quality Versus 2009/2010 RSA Wheat Quality

Country of origin	Australia							RSA Crop Average						
	B1	B2	B3	B4	UT	COW	Average	B1	B2	B3	B4	UT	COW	Average
No. of samples	5	-	-	-	-	-	5	26	28	20	10	5	11	100
<b>ALVEOGRAM</b>														
Strength (S) , cm <sup>2</sup>	40.6	-	-	-	-	-	40.6	42.4	36.5	32.7	25.0	26.8	35.4	35.5
Stability (P), mm	113	-	-	-	-	-	113	93	90	90	87	61	83	89
Distensibility (L), mm	63	-	-	-	-	-	63	97	86	75	58	101	90	85
P/L	1.80	-	-	-	-	-	1.80	1.03	1.13	1.31	1.80	0.61	0.97	1.17
<b>EXTENSOGRAM</b>														
Strength, cm <sup>2</sup>	101	-	-	-	-	-	101	96	85	74	61	71	87	83
Max. height, BU	457	-	-	-	-	-	457	355	344	327	304	295	345	337
Extensibility, mm	153	-	-	-	-	-	153	187	174	158	139	171	173	170
<b>MIXOGRAM</b>														
Peak time, min	3.4	-	-	-	-	-	3.4	2.6	2.6	2.7	2.8	2.8	2.5	2.6
Absorption, %	61.1	-	-	-	-	-	61.1	61.8	60.4	59.4	58.5	60.3	60.6	60.4
<b>MYCOTOXINS</b>														
Aflatoxin, ppb [max.value]	<2 [<2]							1.00 [4.00]						
Deoxynivalenol, ppm [max. value]	0.00 [0.00]							0.05 [0.48]						
Ochratoxin A, ppb [max. value]	0.00 [0.00]							0.17 [1.00]						
No. of samples	1							30						

# 2009/2010 IMPORTED WHEAT QUALITY - BRAZIL (1 Oct 2009 to 30 Sep 2010)

## 2009/2010 Imported Wheat Quality Versus 2009/2010 RSA Wheat Quality

Country of origin		Brazil						RSA Crop Average							
Class and Grade bread wheat		B1	B2	B3	B4	UT	COW	Average	B1	B2	B3	B4	UT	COW	Average
No. of samples		-	-	-	-	-	10	10	159	138	79	30	16	58	480
<b>WHEAT GRADING</b>															
Protein (12% mb), %	-	-	-	-	-	12.28	12.28	12.84	11.50	10.62	9.64	11.29	11.51	11.68	
Moisture, %	-	-	-	-	-	12.2	12.2	11.5	11.3	11.3	10.8	11.0	11.0	11.3	
Falling number, sec	-	-	-	-	-	274	274	382	375	359	339	240	365	367	
1000 Kernel mass (13% mb), g	-	-	-	-	-	31.7	31.7	38.7	39.5	40.4	40.2	36.6	38.1	39.2	
HlM (dirty), kg/hl	-	-	-	-	-	76.9	76.9	80.8	80.0	79.5	79.7	75.2	79.0	79.9	
Screenings (<1.8mm), %	-	-	-	-	-	3.05	3.05	1.36	1.30	1.47	1.44	3.36	3.03	1.63	
Gravel, stones, turf and glass, %	-	-	-	-	-	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.00	0.00	
Foreign matter, %	-	-	-	-	-	0.29	0.29	0.05	0.05	0.09	0.09	0.12	0.11	0.07	
Other grain & unthreshed ears, %	-	-	-	-	-	0.15	0.15	0.25	0.26	0.30	0.35	0.64	0.65	0.33	
Heat damaged kernels, %	-	-	-	-	-	0.38	0.38	0.00	0.00	0.00	0.00	0.05	0.00	0.00	
Immature kernels, %	-	-	-	-	-	0.01	0.01	0.07	0.05	0.04	0.03	0.01	0.06	0.05	
Insect damaged kernels, %	-	-	-	-	-	0.32	0.32	0.19	0.26	0.29	0.27	1.10	0.52	0.30	
Heavily frost damaged kernels, %	-	-	-	-	-	0.00	0.00	0.00	0.00	0.00	0.00	0.13	0.00	0.01	
Sprouted kernels, %	-	-	-	-	-	2.99	2.99	0.09	0.11	0.16	0.14	2.73	0.36	0.23	
Total damaged kernels, %	-	-	-	-	-	3.70	3.70	0.36	0.43	0.50	0.44	3.96	0.96	0.60	
Combined deviations, %	-	-	-	-	-	7.19	7.19	2.02	2.04	2.34	2.32	8.07	4.76	2.63	
Field fungi, %	-	-	-	-	-	0.98	0.98	0.19	0.19	0.21	0.15	0.54	0.16	0.20	
Storage fungi, %	-	-	-	-	-	0.11	0.11	0.01	0.01	0.01	0.00	0.03	0.01	0.01	
Ergot, %	-	-	-	-	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
Noxious seeds ( <i>Crotalaria spp.</i> , <i>Datura spp.</i> )	-	-	-	-	-	0	0	0	0	0	0	0	0	0	
Noxious seeds ( <i>Argemone mexicana</i> )	-	-	-	-	-	0	0	0	0	0	0	0	0	0	
Live insects	-	-	-	-	-	Yes	Yes	No	No	No	No	No	No	No	
Undesirable odour	-	-	-	-	-	Yes	Yes	No	No	No	No	No	No	No	
	B1	B2	B3	B4	UT	COW	Average	B1	B2	B3	B4	UT	COW	Average	
<b>No. of samples</b>	-	-	-	-	-	10	10	26	28	20	10	5	11	100	
<b>BÜHLER EXTRACTION, %</b>	-	-	-	-	-	69.8	69.8	74.8	75.1	74.8	73.8	74.1	74.9	74.7	
<b>FLOUR</b>															
Colour, KJ	-	-	-	-	-	0.5	0.5	-2.2	-2.3	-2.3	-2.3	-1.7	-2.2	-2.2	
Protein (12% mb), %	-	-	-	-	-	10.6	10.6	11.7	10.6	9.6	8.6	10.4	10.6	10.5	
Wet Gluten (14% mb), %	-	-	-	-	-	28.5	28.5	32.1	29.0	25.9	22.2	28.3	29.1	28.6	
Dry Gluten (14% mb), %	-	-	-	-	-	9.5	9.5	11.3	10.1	9.0	7.9	9.6	10.2	10.0	
<b>100g BAKING TEST</b>															
Baking water absorption, %	-	-	-	-	-	60.6	60.6	61.5	60.3	59.3	58.2	59.5	60.4	60.2	
Loaf volume, cm <sup>3</sup>	-	-	-	-	-	723	723	902	853	803	727	868	847	843	
Evaluation	-	-	-	-	-	4	4	1	0	0	0	0	0	0	
<b>FARINOGRAM</b>															
Water absorption, %	-	-	-	-	-	64.0	64.0	62.3	61.3	60.5	59.3	58.3	60.6	61.0	
Development time, min	-	-	-	-	-	1.8	1.8	4.8	3.4	2.9	2.0	2.7	3.7	3.5	
Stability, mm	-	-	-	-	-	2.7	2.7	8.4	6.8	5.7	3.8	4.9	6.9	6.6	
Mixing tolerance index, BU	-	-	-	-	-	80	80	39	44	52	71	71	48	49	

## 2009/2010 Imported Wheat Quality Versus 2009/2010 RSA Wheat Quality

Country of origin	Brazil							RSA Crop Average						
Class and Grade bread wheat	B1	B2	B3	B4	UT	COW	Average	B1	B2	B3	B4	UT	COW	Average
No. of samples	-	-	-	-	-	10	10	26	28	20	10	5	11	100
<b>ALVEOGRAM</b>														
Strength (S) , cm <sup>2</sup>	-	-	-	-	-	26.2	26.2	42.4	36.5	32.7	25.0	26.8	35.4	35.5
Stability (P), mm	-	-	-	-	-	95	95	93	90	90	87	61	83	89
Distensibility (L), mm	-	-	-	-	-	50	50	97	86	75	58	101	90	85
P/L	-	-	-	-	-	1.95	1.95	1.03	1.13	1.31	1.80	0.61	0.97	1.17
<b>EXTENSOGRAM</b>														
Strength, cm <sup>2</sup>	-	-	-	-	-	50	50	96	85	74	61	71	87	83
Max. height, BU	-	-	-	-	-	253	253	355	344	327	304	295	345	337
Extensibility, mm	-	-	-	-	-	141	141	187	174	158	139	171	173	170
<b>MIXOGRAM</b>														
Peak time, min	-	-	-	-	-	2.7	2.7	2.6	2.6	2.7	2.8	2.8	2.5	2.6
Absorption, %	-	-	-	-	-	60.5	60.5	61.8	60.4	59.4	58.5	60.3	60.6	60.4
<b>MYCOTOXINS</b>														
Aflatoxin, ppb [max.value]	<2 [<2]							1.00 [4.00]						
Deoxynivalenol, ppm [max. value]	2.90 [3.90]							0.05 [0.48]						
Ochratoxin A, ppb [max. value]	0.00 [0.00]							0.17 [1.00]						
<b>No. of samples</b>	2							30						

# 2009/2010 IMPORTED WHEAT QUALITY - CANADA (1 Oct 2009 to 30 Sep 2010)

## 2009/2010 Imported Wheat Quality Versus 2009/2010 RSA Season

Country of origin		Canada							RSA Crop Average						
Class and Grade bread wheat		B1	B2	B3	B4	UT	COW	Average	B1	B2	B3	B4	UT	COW	Average
No. of samples		5	-	-	-	-	-	5	159	138	79	30	16	58	480
<b>WHEAT GRADING</b>															
Protein (12% mb), %	14.09	-	-	-	-	-	-	14.09	12.84	11.50	10.62	9.64	11.29	11.51	11.68
Moisture, %	12.2	-	-	-	-	-	-	12.2	11.5	11.3	11.3	10.8	11.0	11.0	11.3
Falling number, sec	365	-	-	-	-	-	-	365	382	375	359	339	240	365	367
1000 Kernel mass (13% mb), g	34.6	-	-	-	-	-	-	34.6	38.7	39.5	40.4	40.2	36.6	38.1	39.2
HlM (dirty), kg/hl	82.4	-	-	-	-	-	-	82.4	80.8	80.0	79.5	79.7	75.2	79.0	79.9
Screenings (<1.8mm), %	2.11	-	-	-	-	-	-	2.11	1.36	1.30	1.47	1.44	3.36	3.03	1.63
Gravel, stones, turf and glass, %	0.00	-	-	-	-	-	-	0.00	0.00	0.00	0.01	0.00	0.00	0.00	0.00
Foreign matter, %	0.01	-	-	-	-	-	-	0.01	0.05	0.05	0.09	0.09	0.12	0.11	0.07
Other grain & unthreshed ears, %	0.15	-	-	-	-	-	-	0.15	0.25	0.26	0.30	0.35	0.64	0.65	0.33
Heat damaged kernels, %	0.00	-	-	-	-	-	-	0.00	0.00	0.00	0.00	0.00	0.05	0.00	0.00
Immature kernels, %	0.00	-	-	-	-	-	-	0.00	0.07	0.05	0.04	0.03	0.01	0.06	0.05
Insect damaged kernels, %	0.02	-	-	-	-	-	-	0.02	0.19	0.26	0.29	0.27	1.10	0.52	0.30
Heavily frost damaged kernels, %	0.00	-	-	-	-	-	-	0.00	0.00	0.00	0.00	0.00	0.13	0.00	0.01
Sprouted kernels, %	0.02	-	-	-	-	-	-	0.02	0.09	0.11	0.16	0.14	2.73	0.36	0.23
Total damaged kernels, %	0.04	-	-	-	-	-	-	0.04	0.36	0.43	0.50	0.44	3.96	0.96	0.60
Combined deviations, %	2.31	-	-	-	-	-	-	2.31	2.02	2.04	2.34	2.32	8.07	4.76	2.63
Field fungi, %	0.36	-	-	-	-	-	-	0.36	0.19	0.19	0.21	0.15	0.54	0.16	0.20
Storage fungi, %	0.05	-	-	-	-	-	-	0.05	0.01	0.01	0.01	0.00	0.03	0.01	0.01
Ergot, %	0.01	-	-	-	-	-	-	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Noxious seeds ( <i>Crotalaria spp.</i> , <i>Datura spp.</i> )	0	-	-	-	-	-	-	0	0	0	0	0	0	0	0
Noxious seeds ( <i>Argemone mexicana</i> )	0	-	-	-	-	-	-	0	0	0	0	0	0	0	0
Live insects	No	-	-	-	-	-	-	No	No	No	No	No	No	No	No
Undesirable odour	No	-	-	-	-	-	-	No	No	No	No	No	No	No	No
		B1	B2	B3	B4	UT	COW	Average	B1	B2	B3	B4	UT	COW	Average
<b>No. of samples</b>		5	-	-	-	-	-	5	26	28	20	10	5	11	100
<b>BÜHLER EXTRACTION, %</b>		74.4	-	-	-	-	-	74.4	74.8	75.1	74.8	73.8	74.1	74.9	74.7
<b>FLOUR</b>															
Colour, KJ	-1.9	-	-	-	-	-	-	-1.9	-2.2	-2.3	-2.3	-2.3	-1.7	-2.2	-2.2
Protein (12% mb), %	13.2	-	-	-	-	-	-	13.2	11.7	10.6	9.6	8.6	10.4	10.6	10.5
Wet Gluten (14% mb), %	36.6	-	-	-	-	-	-	36.6	32.1	29.0	25.9	22.2	28.3	29.1	28.6
Dry Gluten (14% mb), %	13.0	-	-	-	-	-	-	13.0	11.3	10.1	9.0	7.9	9.6	10.2	10.0
<b>100g BAKING TEST</b>															
Baking water absorption, %	63.5	-	-	-	-	-	-	63.5	61.5	60.3	59.3	58.2	59.5	60.4	60.2
Loaf volume, cm <sup>3</sup>	988	-	-	-	-	-	-	988	902	853	803	727	868	847	843
Evaluation	1	-	-	-	-	-	-	1	1	0	0	0	0	0	0
<b>FARINOGRAM</b>															
Water absorption, %	63.3	-	-	-	-	-	-	63.3	62.3	61.3	60.5	59.3	58.3	60.6	61.0
Development time, min	5.4	-	-	-	-	-	-	5.4	4.8	3.4	2.9	2.0	2.7	3.7	3.5
Stability, mm	10.7	-	-	-	-	-	-	10.7	8.4	6.8	5.7	3.8	4.9	6.9	6.6
Mixing tolerance index, BU	31	-	-	-	-	-	-	31	39	44	52	71	71	48	49

## 2009/2010 Imported Wheat Quality Versus 2009/2010 RSA Wheat Quality

Country of origin		Canada							RSA Crop Average						
Class and Grade bread wheat		B1	B2	B3	B4	UT	COW	Average	B1	B2	B3	B4	UT	COW	Average
No. of samples		5	-	-	-	-	-	5	26	28	20	10	5	11	100
<b>ALVEOGRAM</b>															
Strength (S) , cm <sup>2</sup>	52.2	-	-	-	-	-	-	52.2	42.4	36.5	32.7	25.0	26.8	35.4	35.5
Stability (P), mm	88	-	-	-	-	-	-	88	93	90	90	87	61	83	89
Distensibility (L), mm	128	-	-	-	-	-	-	128	97	86	75	58	101	90	85
P/L	0.69	-	-	-	-	-	-	0.69	1.03	1.13	1.31	1.80	0.61	0.97	1.17
<b>EXTENSOGRAM</b>															
Strength, cm <sup>2</sup>	120	-	-	-	-	-	-	120	96	85	74	61	71	87	83
Max. height, BU	353	-	-	-	-	-	-	353	355	344	327	304	295	345	337
Extensibility, mm	238	-	-	-	-	-	-	238	187	174	158	139	171	173	170
<b>MIXOGRAM</b>															
Peak time, min	2.9	-	-	-	-	-	-	2.9	2.6	2.6	2.7	2.8	2.8	2.5	2.6
Absorption, %	63.9	-	-	-	-	-	-	63.9	61.8	60.4	59.4	58.5	60.3	60.6	60.4
<b>MYCOTOXINS</b>															
Aflatoxin, ppb [max.value]	<2 [<2]							1.00 [4.00]							
Deoxynivalenol, ppm [max. value]	1.40 [1.40]							0.05 [0.48]							
Ochratoxin A, ppb [max. value]	1.00 [1.00]							0.17 [1.00]							
No. of samples	1							30							

# 2009/2010 IMPORTED WHEAT QUALITY - GERMANY (1 Oct 2009 to 30 Sep 2010)

## 2009/2010 Imported Wheat Quality Versus 2009/2010 RSA Season

Country of origin		Germany							RSA Crop Average						
Class and Grade bread wheat		B1	B2	B3	B4	UT	COW	Average	B1	B2	B3	B4	UT	COW	Average
No. of samples		-	39	19	-	12	-	70	159	138	79	30	16	58	480
<b>WHEAT GRADING</b>															
Protein (12% mb), %	-	11.24	10.91	-	11.42	-	11.18	12.84	11.50	10.62	9.64	11.29	11.51	11.68	
Moisture, %	-	12.3	12.6	-	12.5	-	12.4	11.5	11.3	11.3	10.8	11.0	11.0	11.3	
Falling number, sec	-	383	356	-	399	-	378	382	375	359	339	240	365	367	
1000 Kernel mass (13% mb), g	-	43.0	44.3	-	41.7	-	43.1	38.7	39.5	40.4	40.2	36.6	38.1	39.2	
HlM (dirty), kg/hl	-	79.2	78.7	-	77.2	-	78.7	80.8	80.0	79.5	79.7	75.2	79.0	79.9	
Screenings (<1,8mm), %	-	1.60	1.40	-	1.67	-	1.56	1.36	1.30	1.47	1.44	3.36	3.03	1.63	
Gravel, stones, turf and glass, %	-	0.00	0.00	-	0.00	-	0.00	0.00	0.00	0.01	0.00	0.00	0.00	0.00	
Foreign matter, %	-	0.06	0.05	-	0.08	-	0.06	0.05	0.05	0.09	0.09	0.12	0.11	0.07	
Other grain & unthreshed ears, %	-	0.40	0.50	-	1.50	-	0.62	0.25	0.26	0.30	0.35	0.64	0.65	0.33	
Heat damaged kernels, %	-	0.01	0.01	-	0.00	-	0.01	0.00	0.00	0.00	0.00	0.05	0.00	0.00	
Immature kernels, %	-	0.02	0.00	-	0.02	-	0.01	0.07	0.05	0.04	0.03	0.01	0.06	0.05	
Insect damaged kernels, %	-	0.02	0.03	-	0.02	-	0.02	0.19	0.26	0.29	0.27	1.10	0.52	0.30	
Heavily frost damaged kernels, %	-	0.07	0.03	-	0.04	-	0.05	0.00	0.00	0.00	0.00	0.13	0.00	0.01	
Sprouted kernels, %	-	0.04	0.02	-	0.03	-	0.03	0.09	0.11	0.16	0.14	2.73	0.36	0.23	
Total damaged kernels, %	-	0.08	0.07	-	0.06	-	0.07	0.36	0.43	0.50	0.44	3.96	0.96	0.60	
Combined deviations, %	-	2.14	2.02	-	3.31	-	2.31	2.02	2.04	2.34	2.32	8.07	4.76	2.63	
Field fungi, %	-	0.12	0.09	-	0.09	-	0.10	0.19	0.19	0.21	0.15	0.54	0.16	0.20	
Storage fungi, %	-	0.01	0.01	-	0.01	-	0.01	0.01	0.01	0.01	0.00	0.03	0.01	0.01	
Ergot, %	-	0.00	0.00	-	0.00	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
Noxious seeds ( <i>Crotalaria spp.</i> , <i>Datura spp.</i> )	-	0	0	-	0	-	0	0	0	0	0	0	0	0	
Noxious seeds ( <i>Argemone mexicana</i> )	-	0	0	-	0	-	0	0	0	0	0	0	0	0	
Live insects	-	No	No	-	No	-	No	No	No	No	No	No	No	No	
Undesirable odour	-	No	No	-	No	-	No	No	No	No	No	No	No	No	
	B1	B2	B3	B1	UT	B1	Average	B1	B2	B3	B4	UT	COW	Average	
<b>No. of samples</b>	-	39	19	-	12	-	70	26	28	20	10	5	11	100	
<b>BÜHLER EXTRACTION, %</b>	-	74.3	74.2	-	73.6	-	74.2	74.8	75.1	74.8	73.8	74.1	74.9	74.7	
<b>FLOUR</b>															
Colour, KJ	-	-1.2	-1.4	-	-1.3	-	-1.3	-2.2	-2.3	-2.3	-2.3	-1.7	-2.2	-2.2	
Protein (12% mb), %	-	10.1	9.8	-	10.3	-	10.0	11.7	10.6	9.6	8.6	10.4	10.6	10.5	
Wet Gluten (14% mb), %	-	27.2	26.5	-	27.5	-	27.0	32.1	29.0	25.9	22.2	28.3	29.1	28.6	
Dry Gluten (14% mb), %	-	9.7	9.3	-	9.9	-	9.6	11.3	10.1	9.0	7.9	9.6	10.2	10.0	
<b>100g BAKING TEST</b>															
Baking water absorption, %	-	59.8	59.5	-	60.1	-	59.7	61.5	60.3	59.3	58.2	59.5	60.4	60.2	
Loaf volume, cm <sup>3</sup>	-	759	726	-	799	-	757	902	853	803	727	868	847	843	
Evaluation	-	2	2	-	1	-	2	1	0	0	0	0	0	0	
<b>FARINOGRAM</b>															
Water absorption, %	-	59.4	59.0	-	58.9	-	59.2	62.3	61.3	60.5	59.3	58.3	60.6	61.0	
Development time, min	-	1.9	1.9	-	2.1	-	1.9	4.8	3.4	2.9	2.0	2.7	3.7	3.5	
Stability, mm	-	4.3	4.6	-	5.0	-	4.5	8.4	6.8	5.7	3.8	4.9	6.9	6.6	
Mixing tolerance index, BU	-	58	58	-	54	-	57	39	44	52	71	71	48	49	

## 2009/2010 Imported Wheat Quality Versus 2009/2010 RSA Wheat Quality

Country of origin		Germany							RSA Crop Average						
Class and Grade bread wheat		B1	B2	B3	B4	UT	COW	Average	B1	B2	B3	B4	UT	COW	Average
No. of samples		-	39	19	-	12	-	70	26	28	20	10	5	11	100
<b>ALVEOGRAM</b>															
Strength (S) , cm <sup>2</sup>	-	33.9	31.1	-	34.3	-	33.2	42.4	36.5	32.7	25.0	26.8	35.4	35.5	
Stability (P), mm	-	101	95	-	93	-	98	93	90	90	87	61	83	89	
Distensibility (L), mm	-	61	61	-	68	-	62	97	86	75	58	101	90	85	
P/L	-	1.70	1.62	-	1.39	-	1.63	1.03	1.13	1.31	1.80	0.61	0.97	1.17	
<b>EXTENSOGRAM</b>															
Strength, cm <sup>2</sup>	-	84	81	-	87	-	84	96	85	74	61	71	87	83	
Max. height, BU	-	393	379	-	385	-	387	355	344	327	304	295	345	337	
Extensibility, mm	-	147	146	-	154	-	148	187	174	158	139	171	173	170	
<b>MIXOGRAM</b>															
Peak time, min	-	3.4	3.6	-	3.4	-	3.5	2.6	2.6	2.7	2.8	2.8	2.5	2.6	
Absorption, %	-	59.9	59.6	-	60.1	-	59.9	61.8	60.4	59.4	58.5	60.3	60.6	60.4	
<b>MYCOTOXINS</b>															
Aflatoxin, ppb [max.value]	1.41 [4.00]							1.00 [4.00]							
Deoxynivalenol, ppm [max. value]	0.00 [<0.25]							0.05 [0.48]							
Ochratoxin A, ppb [max. value]	0.15 [1.00]							0.17 [1.00]							
No. of samples	34							30							

# 2009/2010 IMPORTED WHEAT QUALITY - USA (1 Oct 2009 to 30 Sep 2010)

## 2009/2010 Imported Wheat Quality Versus 2009/2010 RSA Season

Country of origin		USA							RSA Crop Average						
Class and Grade bread wheat		B1	B2	B3	B4	UT	COW	Average	B1	B2	B3	B4	UT	COW	Average
No. of samples		1	5	3	-	1	-	10	159	138	79	30	16	58	480
<b>WHEAT GRADING</b>															
Protein (12% mb), %	12.16	11.59	10.73	-	11.98	-	11.43	12.84	11.50	10.62	9.64	11.29	11.51	11.68	
Moisture, %	10.9	11.1	11.2	-	11.9	-	11.2	11.5	11.3	11.3	10.8	11.0	11.0	11.3	
Falling number, sec	536	470	440	-	499	-	471	382	375	359	339	240	365	367	
1000 Kernel mass (13% mb), g	31.1	31.3	31.7	-	28.9	-	31.2	38.7	39.5	40.4	40.2	36.6	38.1	39.2	
HlM (dirty), kg/hl	82.1	81.1	80.1	-	79.0	-	80.7	80.8	80.0	79.5	79.7	75.2	79.0	79.9	
Screenings (<1,8mm), %	2.44	2.55	2.22	-	3.15	-	2.50	1.36	1.30	1.47	1.44	3.36	3.03	1.63	
Gravel, stones, turf and glass, %	0.00	0.00	0.00	-	0.00	-	0.00	0.00	0.00	0.01	0.00	0.00	0.00	0.00	
Foreign matter, %	0.12	0.08	0.04	-	0.20	-	0.09	0.05	0.05	0.09	0.09	0.12	0.11	0.07	
Other grain & unthreshed ears, %	0.16	0.24	0.33	-	0.00	-	0.24	0.25	0.26	0.30	0.35	0.64	0.65	0.33	
Heat damaged kernels, %	0.00	0.10	0.08	-	0.00	-	0.08	0.00	0.00	0.00	0.00	0.05	0.00	0.00	
Immature kernels, %	0.00	0.00	0.00	-	0.00	-	0.00	0.07	0.05	0.04	0.03	0.01	0.06	0.05	
Insect damaged kernels, %	0.00	0.18	0.23	-	0.08	-	0.17	0.19	0.26	0.29	0.27	1.10	0.52	0.30	
Heavily frost damaged kernels, %	0.00	0.00	0.00	-	0.00	-	0.00	0.00	0.00	0.00	0.00	0.13	0.00	0.01	
Sprouted kernels, %	0.00	0.20	0.03	-	0.00	-	0.11	0.09	0.11	0.16	0.14	2.73	0.36	0.23	
Total damaged kernels, %	0.00	0.46	0.34	-	0.08	-	0.34	0.36	0.43	0.50	0.44	3.96	0.96	0.60	
Combined deviations, %	2.72	3.33	2.93	-	3.43	-	3.16	2.02	2.04	2.34	2.32	8.07	4.76	2.63	
Field fungi, %	0.08	0.14	0.37	-	0.66	-	0.26	0.19	0.19	0.21	0.15	0.54	0.16	0.20	
Storage fungi, %	0.08	0.00	0.00	-	0.08	-	0.02	0.01	0.01	0.01	0.00	0.03	0.01	0.01	
Ergot, %	0.00	0.00	0.00	-	0.01	-	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
Noxious seeds ( <i>Crotalaria spp.</i> , <i>Datura spp.</i> )	0	0	0	-	0	-	0	0	0	0	0	0	0	0	
Noxious seeds ( <i>Argemone mexicana</i> )	0	0	0	-	0	-	0	0	0	0	0	0	0	0	
Live insects	No	No	No	-	No	-	No	No	No	No	No	No	No	No	
Undesirable odour	No	No	No	-	No	-	No	No	No	No	No	No	No	No	
		B1	B2	B3	B4	UT	COW	Average	B1	B2	B3	B4	UT	COW	Average
No. of samples		1	5	3	-	1	-	10	26	28	20	10	5	11	100
BÜHLER EXTRACTION, %	73.7	72.8	72.8	-	73.6	-	73.0	74.8	75.1	74.8	73.8	74.1	74.9	74.7	
<b>FLOUR</b>															
Colour, KJ	-1.3	-1.3	-0.4	-	-1.1	-	-1.0	-2.2	-2.3	-2.3	-2.3	-1.7	-2.2	-2.2	
Protein (12% mb), %	11.0	10.4	9.5	-	10.7	-	10.2	11.7	10.6	9.6	8.6	10.4	10.6	10.5	
Wet Gluten (14% mb), %	26.7	26.2	23.3	-	26.3	-	25.4	32.1	29.0	25.9	22.2	28.3	29.1	28.6	
Dry Gluten (14% mb), %	9.7	9.4	8.1	-	9.7	-	9.1	11.3	10.1	9.0	7.9	9.6	10.2	10.0	
<b>100g BAKING TEST</b>															
Baking water absorption, %	60.9	59.2	57.3	-	58.6	-	58.8	61.5	60.3	59.3	58.2	59.5	60.4	60.2	
Loaf volume, cm <sup>3</sup>	875	773	737	-	870	-	782	902	853	803	727	868	847	843	
Evaluation	0	2	1	-	0	-	1	1	0	0	0	0	0	0	
<b>FARINOGRAM</b>															
Water absorption, %	58.8	57.1	55.7	-	55.9	-	56.7	62.3	61.3	60.5	59.3	58.3	60.6	61.0	
Development time, min	2.2	2.1	1.8	-	1.7	-	2.0	4.8	3.4	2.9	2.0	2.7	3.7	3.5	
Stability, mm	8.0	7.3	4.1	-	8.7	-	6.6	8.4	6.8	5.7	3.8	4.9	6.9	6.6	
Mixing tolerance index, BU	37	48	62	-	38	-	50	39	44	52	71	71	48	49	

## 2009/2010 Imported Wheat Quality Versus 2009/2010 RSA Wheat Quality

Country of origin	USA							RSA Crop Average						
Class and Grade bread wheat	B1	B2	B3	B4	UT	COW	Average	B1	B2	B3	B4	UT	COW	Average
No. of samples	1	5	3	-	1	-	10	26	28	20	10	5	11	100
<b>ALVEOGRAM</b>														
Strength (S) , cm <sup>2</sup>	38.2	33.1	27.8	-	29.5	-	31.7	42.4	36.5	32.7	25.0	26.8	35.4	35.5
Stability (P), mm	86	81	74	-	59	-	78	93	90	90	87	61	83	89
Distensibility (L), mm	84	75	74	-	99	-	78	97	86	75	58	101	90	85
P/L	1.02	1.08	1.08	-	0.60	-	1.03	1.03	1.13	1.31	1.80	0.61	0.97	1.17
<b>EXTENSOGRAM</b>														
Strength, cm <sup>2</sup>	110	95	93	-	112	-	98	96	85	74	61	71	87	83
Max. height, BU	435	419	417	-	430	-	421	355	344	327	304	295	345	337
Extensibility, mm	174	161	154	-	186	-	163	187	174	158	139	171	173	170
<b>MIXOGRAM</b>														
Peak time, min	3.9	4.5	4.9	-	4.8	-	4.6	2.6	2.6	2.7	2.8	2.8	2.5	2.6
Absorption, %	60.9	60.2	59.3	-	60.6	-	60.1	61.8	60.4	59.4	58.5	60.3	60.6	60.4
<b>MYCOTOXINS</b>														
Aflatoxin, ppb [max.value]	0.40 [2.00]							1.00 [4.00]						
Deoxynivalenol, ppm [max. value]	0.29 [1.10]							0.05 [0.48]						
Ochratoxin A, ppb [max. value]	0.20 [1.00]							0.17 [1.00]						
No. of samples	5							30						

## Accuracy Awards

SAGL participates in several international proficiency schemes including AACC International, BIPEA and FAPAS, as part of our quality assurance procedures to demonstrate technical competency. SAGL has received the 2009 AACC International Accuracy Award for Feed Analysis (also received in 2004) as well as the AACC International Accuracy Award for Mixograph analysis (also received in 2006, 2007 and 2008).





## CERTIFICATE OF ACCREDITATION

*In terms of section 22(2)(b) of the Accreditation for Conformity Assessment, Calibration and Good Laboratory Practice Act, 2006 (Act 19 of 2006), read with sections 23(1), (2) and (3) of the said Act, I hereby certify that:-*

**SOUTHERN AFRICAN GRAIN LABORATORY**  
Co. reg no: 1997/018518/08

Facility Accreditation Number: **T0116**

Is a South African National Accreditation System accredited Testing laboratory  
provided that all SANAS conditions and requirements are complied with

This certificate is valid as per the scope as stated in the accompanying schedule of accreditation,  
Annexure "A", bearing the above accreditation number for

### **CHEMICAL & PHYSICAL ANALYSIS**

*The facility is accredited in accordance with the recognised International Standard*

**ISO/I EC 17025:2005**

*The accreditation demonstrates technical competency for a defined scope and the operation of a  
laboratory quality management system*

While this certificate remains valid, the Accredited Facility named above is authorised to  
use the relevant SANAS accreditation symbol to issue facility reports and/or certificates

  
\_\_\_\_\_  
M P R Josias  
Chief Executive Officer

Effective Date: 01 November 2009  
Certificate Expires: 31 October 2014

## ANNEXURE A

## SCHEDULE OF ACCREDITATION

Testing Laboratory Number: T0116

<b>Permanent Address of Laboratory:</b> Southern African Grain Laboratory Grain Building 477 Witjhente Road The Willows 0040	<b>Technical Signatories</b>	Mr J Nortje (All) Ms M Hammett (Chemical) Ms M E Vorster (Physical) Mr B van der Linde (Grading) Ms A de Jager (Vitamins & Minerals) Mrs M Henning (In House Method 11) Ms H Schoeman (In House Method 24)
<b>Postal Address:</b> PostNet Suite # 391 Private Bag X 1 The Willows 0041	<b>Nominated Representative</b>	Mrs B du Pless
Tel (012) 807-4019 Fax (012) 807-4160 E-mail info@sagl.co.za	<b>Management Representative</b>	Mrs W Louw
Issue No. 15 Date of issue 17 February 2011 Expiry date 31 October 2014		
Materials/Products Tested	Types of Tests/Properties Measured, Range of Measurement	Standard Specifications, Equipment/ Techniques Used
<b>CHEMICAL</b>		
Ground barley	Moisture (Oven method)	Analytical EBC 3.2, 1997
Ground grains, semolina and flour	Moisture (Oven method)	ICC No 110/1, 1976
Whole and milled maize and soya beans	Moisture (Oven method)	AACC 44-15.02, 1999
All flours, cereal grains, oil seeds and animal feeds	Nitrogen and protein (Combustion-method)	AACC 45-30.01, 1999
Food stuffs	Dietary fibre (total)	In-House Method 12
Food Stuff and Feeds	Carbohydrates (by difference) (calculation) Energy Value (calculation) Total Digestible Nutritional Value (calculation)	SOP MC 23
Food Stuff and Feeds, Semolina and Milled Pasta	Determination of Ash	In-House Method 11
Wheat Kernels,	Moisture (oven method)	Wheat Grading Regulation No. R005 including amendments
Flours of grains, e.g. barley, triticale, maize, rye, sorghum and wheat, oilseeds, feeds, mixed feeds and foodstuffs	Crude fat (Ether extraction by Soxhlet)	In-House Method 24

Original date of accreditation: 01 November 1999

Page 1 of 2



Signature of Field Manager

Field Manager

## ANNEXURE A

Laboratory No: T0116  
 Date of issue: 17 February 2011  
 Expiry date: 31 October 2014

Materials/Products Tested	Types of Tests/Properties Measured, Range of Measurement	Standard Specifications, Equipment/ Techniques Used
<b>VITAMINS &amp; MINERALS</b>		
Grain based vitamin-fortified food and feed products and fortification mixes	Vitamin A as all trans-Retinol (Saponification)  Thiamine Mononitrate  Riboflavin (Vitamin B2)  Niacinamide  Pyridoxine Hydrochloride  Folic Acid	In-House Method 1.  In-House Method 2  In-House Method 2  In-House Method 2  In-House Method 3
Grain based fortified food and feed products and fortification mixes	Iron and Zinc (total) by AA	In-House Method 10
<b>GRADING</b>		
Maize	Defective Kernels (white maize, yellow maize)	Government Gazette 32190 (08 May 2009); Regulation R.473 including amendments
Cereals as grain: (Wheat, barley, rye and oats)	Hectolitre mass (Kern 222)	ISO 7971-3
Wheat	Screenings	Government Gazette No. 19036 (10 July 1998); Regulation No. R305 including amendments
Meal and flour of wheat, rye, barley, other grains, starch containing and malted products	Falling Number	ICC No 107/1, 1995
<b>PHYSICAL</b>		
Wheat flour	Alveograph (Rheological properties)	ICC No 121, 1992
Wheat Flour and brown bread flour	Farinograph (Rheological properties)	AACC 54-21.01, 1999 – Constant Flour Weight Procedure
Wheat flour and whole wheat flour of hard/soft/durum wheat	Mixograph (Rheological properties)	Industry Accepted Method 020 (Based on AACC 54-40.02, 1999)

Original date of accreditation: 01 November 1999

Page 2 of 2

ISSUED BY THE SOUTH AFRICAN NATIONAL ACCREDITATION SYSTEM


  
Field Manager

No. R. 1186

17 December 2010

**AGRICULTURAL PRODUCT STANDARDS ACT, 1990**  
**(ACT NO. 119 OF 1990)**

**REGULATIONS RELATING TO THE GRADING, PACKING AND MARKING OF  
 BREAD WHEAT INTENDED FOR SALE IN THE REPUBLIC OF SOUTH AFRICA**

The Minister of Agriculture, Forestry and Fisheries, acting under section 15 of the Agricultural Product Standards Act, 1990 (Act No. 119 of 1990), has

- (a) made the regulations in the Schedule;
- (b) determined that the said regulations shall come into operation on the date of publication; and
- (c) repealed the regulations published in Government Notice No's R 905 of 10 July 1998 as amended by R 1421 of 6 November 1998, R 876 of 14 September 2001, R 979 of 19 July 2002, and R 1210 of 29 August 2003.

**SCHEDULE**

**Definitions**

1 Unless the context otherwise indicates, any word or expression in these regulations to which a meaning has been assigned in the Act shall have that meaning, and

"animal rests" means dead rodents, dead birds and dung.

"bag" means a bag manufactured from - -

- (a) jute or phormium or a mixture of jute and phormium; or
- (b) polypropylene that complies with SABS specification CKS632.

"bulk container" means any vehicle or container in which bulk wheat is stored or transported.

"consignment" means --

- (a) a quantity of wheat of the same class, which belongs to the same owner, delivered at any one time under cover of the same consignment note, delivery note or receipt note, or delivered by the same vehicle or bulk container, or loaded from the same bin of a grain elevator or from a ship's hold, or
- (b) in the case where a quantity referred to in paragraph (a), is subdivided into different grades, each such quantity of each of the different grades.

"container" means a bag or bulk container;

"cultivar list" means the list of cultivars determined from time to time by the Executive Officer: Agricultural Product Standards and which is obtainable from the Executive Officer: Agricultural Product Standards, Private Bag X258, Pretoria, 0001.

"damaged wheat" means wheat --

- (a) which have been damaged by insects;
- (b) which have been distinctly discoloured (orange-brown, dark brown or black) by external heat or as a result of heating caused by internal fermentation in wheat with an excessive

moisture content, excluding wheat kernels in respect of which the discolouration is confined to the germ end;

- (c) which are immature and have a distinctly green colour; and
- (d) in which germination has proceeded to such an extent that the skin covering the embryo has been broken or the developing sprouts and/or rootlets are clearly visible.

"ergot sclerotia" means the sclerotia of the fungus *Claviceps purpurea*; and "ergot" has a corresponding meaning.

"falling number" means the time in seconds according to Hagberg-Perten as a measure of the degree of Alpha-Amylase activity in grain and flour.

"field fungi infected wheat" means wheat of which the kernels are visibly infected with fungi, and that -

- (a) clearly have greyish brush-ends that are discoloured as a whole, or where field fungi growth is present from the brush-ends into the crease;
- (b) have a dull, fetid, chalky or pinkish and shrunken appearance as a result of *Fusarium* infection.

"foreign matter" means all material excluding wheat, other grain and unthreshed ears.

"heavily frost-damaged wheat" means -

- (a) wheat which have been damaged by severe frost during the milk to soft dough stage and which is characterised by the kernel's being fairly plump but covered entirely with small blisters extending into the crease, excluding -
  - (i) kernels in which blistering is confined to the back of the kernel, and
  - (ii) immature wrinkled kernels in which wrinkling has been caused by frost while the kernels were still immature, and
- (b) kernels which have a slightly flaked-off bran coat due to frost. Provided that evidence of frost damage is present and that the bran coat had not been rubbed off due to handling.

"hectolitre mass" means the mass in kilogram per hectolitre.

"insect" in relation to wheat, means any live insect that is injurious to stored grain irrespective of the stage of development of that insect.

"other grain" means the kernel's or pieces of kernel's of barley, oats, triticale, maize, rye and sorghum;

"poisonous seeds" means the seeds or bits of seeds of plant species that may in terms of the Foodstuffs, Cosmetics and Disinfectants Act, 1972 (Act No. 54 of 1972) represent a hazard to human or animal health when consumed, including seeds of *Argemone mexicana*, *Convolvulus* spp., *Crotalaria* spp., *Datura* spp., *Ipomoea purpurea*, *Lathyrus sativus*, *Ricinus communis* or *Xanthium* spp.

"protein content" means the percentage protein in wheat on a 12% moisture basis.

"screenings" means all material that passes through the standard sieve;

"standard sieve" is a slotted sieve - -

- (a) with a flat bottom of metal sheet of 1,0 mm thickness with apertures 12,7 mm long and 1,8 mm wide with rounded ends. The spacing between the slots in the same row must be 2,43 mm wide and the spacing between the rows of slots must be 2,0 mm wide. The slots

- must be alternately orientated with a slot always opposite the solid inter segment of the next row of slots;
- (b) of which the upper surface of the sieve is smooth;
  - (c) with a round frame of suitable material with an inner diameter of between 300 mm and 310 mm maximum and at least 50 mm high;
  - (d) that fits onto a tray with a solid bottom and must be at least 20 mm above the bottom of the tray.

"stinking smut infection" means wheat that is infected with *Tilletia* spp. with the exception of wheat infected with *Tilletia indica* (kaunal bunt). Wheat is considered stinking smut infected if one or more of the following characteristics are present:

- (a) an unmistakable stinking smut odour; or
- (b) wheat kernels that are smeared with stinking smut; or
- (c) more than four stinking smut balls (or pieces of balls equal to four stinking smut balls) per 100 g of wheat;

"storage fungi infected wheat" means wheat that are visibly infected with fungi, and that show -

- (a) blue, green, blackish or yellow fungal growth anywhere on the kernel; or
- (b) visible mould beneath the bran.

"the Act" means the Agricultural Product Standards Act, 1980 (Act No. 119 of 1980).

"unthreshed ears" means ears and nits of ears of wheat, barley, milo and rye that still contain seeds that are completely covered with glumes; and

"wheat" means the kernels and pieces of kernels of the species *Triticum aestivum*.

#### **Restrictions on sale of wheat**

- 2 (1) No person shall sell a consignment of wheat in the Republic of South Africa --
- (a) unless the wheat is sold according to the classes set out in regulation 3;
  - (b) unless the wheat complies with the standards for the classes set out in regulation 4;
  - (c) unless the wheat, where applicable, complies with the grades of wheat and the standards for grades set out in regulations 5 and 6 respectively;
  - (d) unless the wheat is packed in accordance with the packing requirements set out in regulation 7;
  - (e) unless the containers or sale documents, as the case may be, are marked in accordance with the marking requirements set out in regulation 8; and
  - (f) if such wheat contains a substance that renders it unfit for human consumption or for processing into or utilisation thereof as food or feed.
- (2) The Executive Officer may grant written exemption, entirely or partially, to any person on such conditions as he or she may deem necessary from the provisions of subregulation (1).

**PART I****QUALITY STANDARDS****Classes of wheat**

3 The classes of wheat are –

- (a) Bread Wheat; and
- (b) Other Wheat.

**Standards for classes**

4 (1) Notwithstanding the provisions of sub-regulations (2) and (3), all consignments of wheat must –

- (a) be free from any toxin, chemical or other substances that renders it unsuitable for human consumption or for processing into or utilisation thereof as food or feed and may not exceed the permissible deviations regarding aflatoxin in terms of the Foodstuffs, Cosmetics and Disinfectants Act, 1972 (Act No. 54 of 1972);
  - (b) contain not more poisonous seeds or ergot sclerotia than permitted in terms of the Foodstuffs, Cosmetics and Disinfectants Act, 1972 (Act No. 54 of 1972);
  - (c) be free from organisms of phytosanitary importance as determined in terms of the Agricultural Pest Act, 1983 (Act No. 36 of 1983);
  - (d) be free from mould infected, sour and rancid other grain, foreign matter and any other matter;
  - (e) be free from any undesired odour, taste or colour not typical of undamaged and sound wheat;
  - (f) be free from animal pests,
  - (g) with the exception of Other Wheat, be free from insects;
  - (h) with the exception of Other Wheat, be free from stinking smut infection; and
  - (i) with the exception of Other Wheat, have a moisture content not exceeding 13 per cent.
- (2) A consignment shall be classified as Bread Wheat if –
- (a) the wheat in the consignment consists of at least 95 per cent (m/m) of one or more of the bread wheat cultivars specified in the cultivar list; and
  - (b) it complies with the standards for Grade 1, Grade 2, Grade 3, Grade 4 or Utility Grade set out in regulation 6.
- (3) A consignment of wheat shall be classified as Other Wheat if it does not comply with the standards for Bread Wheat.

**Grades of wheat**

5 (1) The grades for Bread Wheat shall be as follows

- (a) Grade 1;
- (b) Grade 2.

- (c) Grade 3;
  - (d) Grade 4; and
  - (e) Utility grade
- (2) No grades are determined for Other Wheat

**Standards for grades of wheat**

5 (1) Subject to the provisions of subregulations (2), (3) and (4) a consignment of wheat shall be graded as --

- (a) Grade 1 if the nature of deviation, specified in column 1 of Table 1 of the Annexure, in that consignment does not exceed the percentage specified in column 2 of the said table opposite the deviation concerned.
  - (b) Grade 2 if the nature of deviation, specified in column 1 of Table 1 of the Annexure, in that consignment does not exceed the percentage specified in column 3 of the said table opposite the deviation concerned.
  - (c) Grade 3 if the nature of deviation, specified in column 1 of Table 1 of the Annexure, in that consignment does not exceed the percentage specified in column 4 of the said table opposite the deviation concerned
  - (d) Grade 4 if the nature of deviation, specified in column 1 of Table 1 of the Annexure, in that consignment does not exceed the percentage specified in column 5 of the said table opposite the deviation concerned and
  - (e) Utility Grade if the nature of deviation, specified in column 1 of Table 1 of the Annexure, in that consignment does not exceed the percentage specified in column 6 of the said table opposite the deviation concerned
- (2) The minimum hectolitre masses for the different grades are as follows.
- (a) Grade 1 - 77 kg,
  - (b) Grade 2 - 76 kg,
  - (c) Grade 3 - 74 kg,
  - (d) Grade 4 - 72 kg, and
  - (e) Utility Grade - 70 kg
- (3) (a) Grade 1, Grade 2 and Grade 3 shall have a minimum falling number value of not less than 250 seconds
- (b) Grade 4 shall have a minimum falling number value of not less than 200 seconds
- (c) Utility Grade shall have a minimum falling number value of not less than 150 seconds
- (d) Notwithstanding the provision of paragraph (a), wheat shall be deemed to comply with the requirements of the paragraph concerned if it deviates with not more than 30 seconds lower than the minimum prescribed for Grade 1, Grade 2 and Grade 3, as the case may be

(4) The minimum protein content (on a 12 per cent moisture basis) for the different grades shall be as follows:

- (a) Grade 1 - 12 per cent
- (b) Grade 2 - 11 per cent.
- (c) Grade 3 - 10 per cent
- (d) Grade 4 - 9 per cent; and
- (e) Utility Grade - 8 per cent.

## PART II

### PACKING AND MARKING REQUIREMENTS

#### *Packing requirements*

7 Wheat of different grades shall be packed in different containers, or stored separately.

#### *Marking requirements*

8 (1) Every container or the accompanying sale documents of a consignment of wheat shall be marked or endorsed by means of appropriate symbols specified in subregulation (2), with --

- (a) the class of the wheat, and
  - (b) the grade
- (2) The symbols referred to in subregulation (1) shall appear in the order of class and grade
- (3) The symbols used to indicate the different --
- (a) classes shall be --
    - (i) B in the case of Bread Wheat; and
    - (ii) O in the case of Other Wheat.
  - (b) grades shall be --
    - (i) 1 in the case of Grade 1;
    - (ii) 2 in the case of Grade 2;
    - (iii) 3 in the case of Grade 3;
    - (iv) 4 in the case of Grade 4; and
    - (v) UT in the case of Utility Grade.

## PART III

### SAMPLING

#### *Taking of sample*

9 (1) A sample of a consignment of wheat shall --

- (a) in the case of wheat delivered in bags and subject to regulation 10, be obtained by sampling at least ten per cent of the bags, chosen from that consignment at random, with a bag probe. Provided that at least 25 bags in a consignment shall be sampled and where a consignment consists of less than 25 bags, all the bags in that consignment shall be sampled, and
- (b) in the case of wheat delivered in bulk and subject to regulation 10, be obtained by sampling that consignment throughout the whole depth of the layer, in at least six different places, chosen at random in that bulk quantity, with a bulk sampling apparatus
- (2) The collective sample obtained in subregulation (1)(a) or (b) shall --
  - (a) have a total mass of at least 10 kg, and
  - (b) be thoroughly mixed by means of dividing before further examination
- (3) If it is suspected that the sample referred to in sub regulation (1)(a) is not representative of that consignment, an additional five per cent of the remaining bags, chosen from that consignment at random, shall be emptied into a suitable bulk container and sampled in the manner contemplated in subregulation (1)(b)
- (4) If it is suspected that the sample referred to in sub regulation (1)(b) is not representative of that consignment, an additional representative sample shall be obtained by using an alternative sampling pattern, apparatus or method
- (5) A sample taken in terms of these regulations shall be deemed to be representative of the consignment from which it was taken

#### ***Sampling if contents differ***

10 (1) If, after an examination of the wheat taken from different bags in a consignment in terms of regulation 9(1)(a), it appears that the contents of those bags differ substantially --

- (a) the bags concerned shall be placed separately;
- (b) all the bags in the consignment concerned shall be sampled with a bag probe in order to do such separation, and
- (c) each group of bags with similar contents in that consignment shall for the purposes of these regulations be deemed to be a separate consignment

(2) If, after the discharge of a consignment of wheat in bulk has commenced, it is suspected that the consignment could be of a class or grade other than that determined by means of the initial sampling, the discharge shall immediately be stopped and the part of the consignment remaining in the bulk container as well as the wheat already in the hopper shall be sampled anew with a bulk sampling apparatus or by catching at least 20 samples, by means of a suitable container, at regular intervals throughout the whole offloading period from the stream of wheat flowing in bulk.

#### ***Working sample***

11 A working sample is obtained by dividing the representative sample of the consignment according to the ICC (International Association for Cereal Science and Technology) 101/1 method.

**PART IV****DETERMINATION OF OTHER SUBSTANCES*****Determination of undesirable odours and harmful substances***

12 A consignment of wheat or a sample of a consignment of wheat shall be sensorially assessed or chemically analysed in order to determine --

- (a) whether it contains a substance that renders the wheat unfit for human consumption or for processing into or for utilisation as food or feed; and
- (b) whether it has a musty, sour, rancid or other undesirable odour. Provided that a working sample of unscreened wheat that is ground in a grain mill to a fine meal may be used for the determination concerned.

**PART V****DETERMINATION OF CLASS, HECTOLITRE MASS,  
MOISTURE CONTENT, PROTEIN CONTENT AND FALLING NUMBER*****Determination of class***

13 The class of a consignment of wheat shall be determined as follows

- (a) Obtain a working sample of at least 500 g and screen the working sample in the manner prescribed in regulation 18.
- (b) Take at least 100 g of the screened wheat and remove all other grain, un threshed ears and foreign matter by hand.
- (c) Obtain a working sample of at least 25 g each after all other grain, un threshed ears and foreign matter have been removed and separate the different cultivars.
- (d) Determine the combined mass of all of the cultivars that belongs according to the cultivar list to the same class and express the mass thus determined as a percentage of the mass of the working sample.
- (e) Such percentage represents the percentage of all the cultivars that belongs according to the cultivar list to the same class in the consignment.

***Determination of the hectolitre mass***

14 The hectolitre mass of a consignment of unscreened wheat may be determined by any suitable instrument. Provided that the instrument complies with and has been calibrated to, the specifications detailed in ISO (International Organization for Standardization) 7971-3.

***Determination of moisture content***

15. The moisture content of a consignment wheat may be determined by any suitable method. Provided that the results thus obtained is in accordance with the maximum permissible deviation for a class 1 moisture meter as detailed in ISO ( International Organization for Standardization) 7700/1 based on the results of the 72 hour 103°C oven dried method [AACC (American Association of Cereal Chemists) Method 44-15A].

***Determination of protein content***

16 The percentage of protein of a consignment of wheat may be determined according to any suitable method. Provided that --

- (a) the determination shall be conducted on a sample which had been sifted using a screen with the same apertures as the standard sieve and from which other grain, un threshed ears and foreign matter had been removed by hand, and
- (b) the results thus obtained are in accordance ( $\pm 0.3$  per cent) with the results obtained by the Cumas Combustion Analysis Method (AACC (American Association of Cereal Chemists) Method 46-30)

**Determination of falling number in wheat**

17. (1) The falling number of a consignment of wheat may be determined according to any suitable method. Provided that --

- (a) the determination shall be conducted on a sample which had been sifted using a screen with the same apertures as the standard sieve and from which other grain, un threshed ears and foreign matter had been removed by hand, and
- (b) the results thus obtained are in accordance ( $\pm 5$  percent) with the results obtained by the ICC (International Association for Cereal Science and Technology) 107/1 method

(2) If the falling number of a consignment of wheat is determined according to the ICC (International Association for Cereal Science and Technology) 107/1 method --

- (a) the sampling in the mentioned method shall be replaced with the manner prescribed in regulation 9, and
- (b) only the altitude corrected value shall be used

**PART VI**

**DETERMINATION OF PERCENTAGE DEVIATIONS**

**Determination of percentage screenings**

18. (1) The percentage screenings in a consignment of wheat shall be determined as follows:
- (a) Obtain a working sample of at least 500 g
  - (b) Place the sample on the standard sieve and screen the sample by moving the sieve 50 strokes to and fro, alternately away from and towards the operator of the sieve, in the same direction as the long axes of the slots of the sieve. Move the sieve which rests on a table or other suitable smooth surface, 250 mm to 460 mm away from and towards the operator with each stroke. The prescribed 50 strokes must be completed within 50 to 60 seconds. Provided that the screening process may also be performed in some or other container or an automatic sieving apparatus
  - (c) Determine the mass of the material that has passed through the sieve and express it as a percentage of the mass of the working sample.
  - (d) Such percentage represents the percentage screenings in the consignment.

**Determination of the percentage heavily frost-damaged wheat**

19. The percentage heavily frost-damaged wheat in a consignment of wheat shall be determined as follows:

- (a) Obtain a working sample of at least 25 g of a screened sample.

- (b) Remove all heavily frost-damaged kernels by hand and determine the mass thereof.
- (c) Express the mass thus determined as a percentage of the mass of the working sample.
- (d) Such percentage represents the percentage heavily frost-damaged wheat in the consignment concerned.

**Determination of the percentages other grain and unthreshed ears**

20 The percentage other grain and unthreshed ears in a consignment of wheat shall be determined as follows:

- (a) Obtain a working sample of at least 50 g from a screened sample.
- (b) Remove all other grain and unthreshed ears by hand and determine the mass thereof.
- (c) Express the mass thus determined as a percentage of the mass of the working sample.
- (d) Such percentage represents the percentage other grain and unthreshed ears in the consignment concerned.

**Determination of the percentage foreign matter**

21 The percentage foreign matter in a consignment of wheat is determined as follows:

- (a) Obtain a working sample of at least 100 g from a screened sample.
- (b) Remove all foreign matter by hand and determine the mass thereof.
- (c) Express the mass thus determined as a percentage of the mass of the working sample.
- (d) Such percentage represents the percentage foreign matter in the consignment concerned.

**Determination of the percentage damaged wheat**

22 The percentage damaged wheat in a consignment of wheat shall be determined as follows:

- (a) Obtain a working sample of at least 25 g of a screened sample.
- (b) Remove all damaged kernels by hand and determine the mass thereof.
- (c) Express the mass thus determined as a percentage of the mass of the working sample.
- (d) Such percentage represents the percentage damaged wheat in the consignment concerned.

**Determination of the percentage heat-damaged wheat**

23 The percentage heat-damaged wheat in a consignment of wheat shall be determined as follows:

- (a) Obtain a working sample of at least 100 g from a screened sample.
- (b) Remove all heat-damaged kernels by hand and determine the mass thereof. Kernels from an additional working sample may also be sensorially assessed (by smelling and tasting the kernel(s)) to confirm suspicion of heat damage.
- (c) Express the mass thus determined as a percentage of the mass of the working sample.

- (d) Such percentage represents the percentage heat-damaged wheat in the consignment concerned.

**Determination of percentage field fungi infected wheat**

24 The percentage field fungi infected wheat in a consignment of wheat shall be determined as follows:

- (a) Obtain a working sample of at least 25 g from a screened sample.
- (b) Remove all field fungi infected kernels by hand and determine the mass thereof.
- (c) Express the mass thus determined as a percentage of the mass of the working sample.
- (d) Such percentage represents the percentage of field fungi infected wheat in the consignment concerned.

**Determination of percentage storage fungi infected wheat**

25 The percentage storage fungi infected wheat in a consignment of wheat shall be determined as follows:

- (a) Obtain a working sample of at least 100 g from a screened sample.
- (b) Remove all storage fungi infected kernels by hand and determine the mass thereof.
- (c) Express the mass thus obtained as a percentage of the mass of the working sample.
- (d) Such percentage represents the percentage storage fungi infected wheat in the consignment concerned.

**PART VII**

**Offence and penalties**

26 Any person who contravenes or fails to comply with any provision of these regulations shall be guilty of an offence and upon conviction be liable to a fine of not exceeding R50 000 or to imprisonment for a period not exceeding two years, or to both that fine or imprisonment.

## ANNEXURE/AANHANGSEL

TABLE 1/TABEL 1

STANDARDS FOR GRADES OF BREAD WHEAT/  
STANDAARDE VIR GRADE VAN BROODKORING

Nature of deviation/ Aard van afwyking	Maximum percentage permissible deviation (m/m)/ Maksimum persentasie toelaatbare afwyking (m/m)					Utility Grade/ Utiliteit- graad
	Grade 1/ Graad 1	Grade 2/ Graad 2	Grade 3/ Graad 3	Grade 4/ Graad 4		
(a) Heavily frost-damaged kernels/ Erg rybeskadigde korrels	3 5	4 5	5 5	6 5	7 10	
(b) Field fungi-infected kernels/Land- swambesmette korrels	2	2	2	2	2	
(c) Storage fungi-infected kernels/ Opbergingswambesmette korrels	0,5	0,5	0,5	0,5	0,5	
(d) Screenings/Sisels	3	3	3	4	10	
(e) Other grain and unthreshed ears/ Ander graan en ongedorsle are	1	1	1	1	4	
(f) Gravel stones, turf and glass/ Gruis, koppies, turf en glas	0,5	0,5	0,5	0,5	0,5	
(g) Foreign matter including gravel, stones, turf and glass: Provided that such deviations are individually within the limits specified in item (f)/ Vreemde voorwerpe met insluiting van gruis, koppies, turf en glas: Met dien verstande dat sodanige afwykings individueel binne die perke's in item (f) aan- gegee	1	1	1	1	3	
(h) Heat-damaged kernels/Hittes- skadigde korrels	0,5	0,5	0,5	0,5	0,5	

<u>Nature of deviation/ Aard van afwyking</u>	<u>Maximum percentage permissible deviation (m/m)</u> <u>Maksimum persentasie toelaatbare afwyking (m/m)</u>					<u>Utility Grade/ Utiliteit- graad</u>
	<u>Grade 1/ Graad 1</u>	<u>Grade 2/ Graad 2</u>	<u>Grade 3/ Graad 3</u>	<u>Grade 4/ Graad 4</u>		
(ii) Damaged kernels, including heat-damaged kernels. Provided that such deviations are individually within the limit specified in item (h) and provided further that the minimum falling number value prescribed in regulation 6(3) for the grade concerned is at least complied with/Beskadigde korrels met inbegrip van hittebeskadigde korrels. Met dien verstande dat sodanige afwyking individueel binne die perke is in item (h) aangegee en met dien verstande voorts dat minstens aan die minimum valgetalwaarde in regulasie 6(3) vir die betrokke graad voldoen word.	1 3 2	4 2	5 2	6 2	7 5	
(j) Deviations in items (d), (e), (g) and (i) collectively. Provided that such deviations are individually within the limits of the said items/Afwykkings in items (d), (e), (g) en (i) gesamentlik. Met dien verstande dat sodanige afwykkings individueel binne die perke van genoemde items is.	5	5	6	5	10	

