

Amino Acid Profile

Amino acids are the building blocks of proteins and approximately 22 amino acids are commonly distributed among the proteins of all biological materials. Of these, 18 can be found in cereal grain proteins. Amino acids are organic compounds containing basic amine (-NH₂) and acidic carboxyl (-COOH) functional groups, in addition to a side chain (R group) specific to each amino acid.

Amino acid composition is an important feature in determining the nutritional value of wheat for human and animal diets. Amino acids are considered crucial to good health, contributing considerably to the health of the human nervous system, hormone production as well as muscular structure and are needed by vital organs and for cellular structure.

The classification of amino acids is based on different features, one being whether the amino acid can be acquired through the diet. According to this, three types are identified: essential, conditionally essential and non-essential amino acids. Classification as essential or non-essential, does however not reflect their actual importance, since all of them are necessary for human health. Essential amino acids are considered “essential” as they cannot be synthesized by the body and must be obtained from the diet. The nine essential amino acids are phenylalanine, valine, threonine, tryptophan, methionine, leucine, isoleucine, lysine and histidine. Arginine, cysteine, glycine, glutamine, proline and tyrosine, are considered conditionally essential in the human diet, meaning their synthesis can be limited under special pathophysiological conditions. Alanine, aspartic acid, asparagine, glutamic acid and serine are non-essential amino acids, meaning they can be synthesized by the body.^(1,2)

Whole meal and white flours from different classes and varieties of wheat grown in three different countries (USA, USSR and Australia) show generally similar amino acid compositions, with high contents of glutamic acid (including glutamine) and proline, very low tryptophan, and relatively low contents of lysine and threonine. Compared with the range of variation in the protein contents of these samples, the variation in their amino acid compositions resulting either from genetic differences or growing conditions such as fertilizer level is rather limited. The reason being that the amino acid compositions of the major endosperm proteins, representing close to 80% of the total wheat proteins, are very similar. Differences in the expression of levels of individual protein genes therefore do not usually result in significant differences in the amino acid compositions of the samples.

Significant variation in amino acid composition may however occur in extreme cases. Research showed that lysine and threonine were for example higher in yellow-berry kernels compared to normal kernels, while glutamic acid (including glutamine) was significantly lower. Extreme differences in fertilization conditions may also result in significant variation in the compositions of whole grain flour or specific protein fractions. It has been found that the proportions of glutamine, proline and phenylalanine in wheat grain and flour all increased with increased levels of nitrogen fertilization, whereas threonine, serine, glycine, alanine, valine and sulphur amino acids decreased.

In a study where wheat was grown under even more extreme fertilization conditions, with variation in nitrogen and sulphur, the grain amino acid composition changed significantly. Less than half of the amounts of cysteine and methionine were present in grain grown with no sulphur and high nitrogen levels compared to grain grown with adequate levels of sulphur.

Proteins are not distributed uniformly throughout a wheat kernel with variation occurring in both the protein content and composition. Even though the starchy endosperm's protein concentration is only a third of that in the germ and less than half of that in the aleurone layer, the starchy endosperm proteins still represent close to three quarters of the total grain protein. The starchy endosperm is characterized by high levels of glutamine and proline and low levels of basic amino acids, while the aleurone and germ contain significantly less proline and glutamine, with high levels of arginine and asparagine in the aleurone layer and germ, respectively. Since the various morphological parts of the wheat kernel differ in protein content and composition, milling extraction rates affect the content and composition of flour.

A large number of flour fractions or mill streams are produced during commercial flour milling which are recombined to provide flours with specific processing characteristics. As a result of the irregular distribution of components within the wheat kernel, these flour streams also vary in their composition and functional properties.

The amino acid compositions of flours differ from those of the grains from which they were milled in containing less lysine, arginine, aspartate (+ asparagine), glycine and alanine but more glutamate (+ glutamine) and proline. Analyses of manually dissected pericarp, testa, aleurone, starchy endosperm and germ suggest that these differences in composition result from differences in the distribution of protein classes throughout the wheat kernel, for example the proportions of basic amino acids increase and the nitrogen content decreases from the outside towards the centre of the endosperm.⁽³⁾

Due to the fact that cystine consists of two cysteine molecules, joined by a disulfide (S-S) bond, cysteine and cystine are interchangeable in wheat. The ratio of cysteine to cystine is dependent on the degree of oxidation in a dough. Addition of an oxidizing agent, such as ascorbic acid, will increase the amount of cystine at the expense of cysteine. This has a “strengthening” effect on the gluten by increasing its elasticity.⁽²⁾

The results of the samples analysed by SAGL and reported as g amino acid/100 g sample, are provided in Table 7 on pages 68 and 69. The values obtained for all amino acids on these samples, were within the normal range reported for wheat in literature, deficient in certain essential amino acids, such as tryptophan, lysine, threonine, methionine and histidine, but high in glutamic acid and proline, which is not essential.

Due to the fact that protein and amino acid composition of wheat varies with crop varieties, application of fertilizers, irrigation practices, soil composition and climatic conditions, the amino acid content showed a wide variation between samples. The only exception being tryptophan, ranging from 0.14 to 0.17 g/100 g this season and between 0.13 to 0.19 g/100 g the previous season. Tryptophan is nutritionally important since it is a precursor for important metabolites such as serotonin and nicotinamide. The World Health Organisation’s (WHO) recommended daily dose for tryptophan is 4 mg/kg/day.⁽⁴⁾

Lysine values varied between 0.33 and 0.44 g/100 g, comprising \pm 2.4 – 3.2 % of the total amino acid content. Lysine is the precursor for carnitine and is required for the structural modification of collagen together with the amino acids glycine and proline.⁽¹⁾ The WHO recommended daily dose for lysine is 30 mg/kg/day. Threonine’s WHO recommended daily dose is 15 mg/kg/day⁽⁴⁾ and ranged from 0.35 to 0.52 g/100 g this season and from 0.32 to 0.54 g/100 g the previous season. Threonine supports digestive function, the immune system, liver and cardiovascular function as well as the central nervous system.⁽⁵⁾

Methionine values were higher this season than during the previous season, ranging from 0.22 to 0.37 g/100 g, compared to 0.18 and 0.26 g/100 g previously. The WHO daily recommendation is 15 mg/kg/day for the sulphur containing amino acids in total.⁽⁴⁾ The main functions of methionine include building of various protein molecules and the synthesis of the equally important sulphurous amino acid, cysteine.⁽⁵⁾

The values for histidine varied between 0.26 and 0.39 g/100 g and comprises \pm 2.1 – 2.3 % of the total amino acid content. Histidine is involved in the formation of proteins, it influences several of the metabolic reactions in the body and assists in regulating blood pH values. The results also showed that the samples were high in the essential amino acid leucine, with values ranging from 0.43 to 1.20 g/100 g. Phenylalanine values varied between 0.53 and 0.88 g/100 g.

The results showed that the samples were rich in glutamic acid and proline, together contributing \pm 42% of the total amino acid content. Glutamic acid contributes to the health of the immune and digestive systems, as well as energy production. Proline is a non-essential amino acid manufactured mainly from ornithine, glutamine and glutamate in the liver and is one of the principal amino acids required by the body for building collagen.⁽⁵⁾

References:

1. Richard D. Semba, Michelle Shardell, Fayrouz A. Sakr Ashour, Ruin Moaddel, Indi Trehan, Kenneth M. Maleta, M. Isabel Ordiz, Klaus Kraemer, Mohammed A. Khadeer, Luigi Ferrucci, Mark J. Manary. (2016). Child Stunting is Associated with Low Circulating Essential Amino Acids. *EbioMedicine* (6), page 246-252.
2. Williams, P. 2011. *A Practical Introduction to Cereal Chemistry*. First Choice Books, page 4-19.
3. Khan, K. and Shrewry, P.P. Eds. 2009. *Wheat Chemistry and Technology*. 4th Ed: Chapter 8.
4. World Health Organization. Protein and amino acids requirements in human nutrition: report of a joint FAO/WHO/UNU expert consultation, Teck. Rep. Series no 935, World Health Organization, Geneva, Switzerland 2007.
5. <http://aminoacidstudies.org>