

Please see a comparison of the moisture, crude protein and crude fat results between the crop survey and ARC Grain Crops sunflower cultivar trials' samples in Table 4.

<b>Table 4: Comparison between the moisture, crude protein and crude fat results of the sunflower crop quality and ARC cultivar trial samples of the 2020/21 season</b>			
<b>Analysis</b>	<b>Moisture, % (5hr, 105°C)</b>	<b>Crude Protein, % (as is)</b>	<b>Crude Fat, % (as is)</b>
<b>Sunflower Crop Quality Survey results</b>			
<b>Ave</b>	5.0	15.02	39.5
<b>Min</b>	3.9	11.41	33.3
<b>Max</b>	6.9	18.04	46.1
<b>Stdev</b>	0.53	1.22	1.89
<b>N</b>	<b>157</b>	<b>157</b>	<b>157</b>
<b>ARC Grains Crops Cultivar trial sample results</b>			
<b>Ave</b>	5.2	16.82	42.5
<b>Min</b>	2.3	10.65	32.8
<b>Max</b>	6.3	23.33	54.3
<b>Stdev</b>	0.69	3.93	4.24
<b>N</b>	<b>147</b>	<b>147</b>	<b>147</b>
<b>% Difference between crop and cultivar samples</b>	<b>-0.2</b>	<b>-1.80</b>	<b>-3.0</b>

See Table 5 on page 23 for a summary of the RSA Sunflower Crop Quality averages of the 2020/21 season compared to those of the 2019/20 season.

Please also see pages 24 to 30 for the average sunflower quality per region.

